



Cumulative Contents

Vol. 151-249

Vol. 151/1, 1985

Changes in serum osteocalcin associated with parathyroid hormone infusion in X-linked hypophosphatemic rickets Cole D.E.C., Gundberg C.M.	1
Comparison of ¹⁵ N analysis by optical emission spectrometry and mass spectrometry for clinical studies during total parenteral nutrition Ragon A., Reynier J.P., Guiraud G., Francois G.	9
Improved method of thyroid peroxidase extraction from the human thyroid gland Sugawara M., Thayer C.L., Kita T., Kuma K.	17
Comparison of enzymatic characteristics of creatine kinase BB from human healthy and tumor stomach tissues Kanemitsu F., Okigaki T.	23
Salivary cortisol for the evaluation of Cushing's syndrome Luthold W.W., Marcondes J.A.M., Wajchenberg B.L.	33
The role of urea in albumin dimerisation in nephrotic urines Birtwistle R.J., Hardwicke J.	41
Serum and interstitial fluid apolipoprotein E levels in the healthy and in hyperlipoproteinemia type III as studied by radioimmunoassay Gustafson S., Vessby B., Ostlund-Lindqvist A.M.	49
Enzyme linked immunosorbent assay for lactoferrin. Plasma and tissue measurements Bezwdoda W.R., Baynes R.D., Khan Q., Mansoor N.	61
Determination of choline-containing phospholipids in human bile and serum by a new enzyme sensor Campanella L., Mascini M., Palleschi G., Tomassetti M.	71
Colorimetric measurement of serum magnesium with an IL micro centrifugal analyzer Datta P., Graham G.A.	85
A quantitative IgE radioimmunoassay in dried blood spots suitable for neonatal screening of atopy Adriaenssens K.M., Philips E.S., Colfs B.	91

Vol. 151/2, 1985

Serum bile acid composition in patients with cystic fibrosis Setchell K.D.R., Smethurst P., Giunta A.M., Colombo C.	101
Release pattern of CK-MB and mitochondrial CK following myocardial ischaemia Sylvén C., Kallner A., Henze A. et al.	111
Heterogeneity of ¹²⁵ I-labeled human thyroglobulin preparations Spencer C.A., Platler B.W., Guttler R.B., Nicoloff J.T.	121
Urinary kallikrein: Assay validation and physiological variability Ash K.O., Smith J.B., Lynch M. et al.	133

Glycosaminoglycan excretion in random samples of urine	
Huang K.-C., Sukegawa K., Orii T.	141
Screening test for urinary glycosaminoglycans and differentiation of various mucopolysaccharidoses	
Huang K.-C., Sukegawa K., Orii T.	147
Immunological study of tissue polypeptide antigen (TPA) — demonstration of keratin-like sites and blood group antigen-like sites on TPA molecules	
Ochi Y., Ura Y., Hamazu M. et al.	157
Erythrocyte malondialdehyde release in vitro: A functional measure of vitamin E status	
Cynamon H.A., Isenberg J.N., Nguyen C.H.	169
Direct automated assay method for serum or urine levels of ketone bodies	
Harano Y., Ohtsuki M., Ida M. et al.	177
Basal level of human growth hormone (hGH) in normal serum	
Hashida S., Nakagawa K., Ishikawa E., Ohtaki S.	185
A simple method for quantitative determination of lipoprotein-X (LP-X) in human serum	
Panteghini M., Malchiodi A., Bonora R.	187

Vol. 151/3, 1985

Free drug measurements: Methodology and clinical significance	
Kwong T.C.	193
Deficiency of acyl-CoA: Dihydroxyacetone phosphate acyltransferase in thrombocytes of Zellweger patients: A simple postnatal diagnostic test	
Wanders R.J.A., Van Wieringh G., Schrakamp G. et al.	217
Circulating prostatic acid phosphatase-immunoglobulin complexes in Sjögren's syndrome	
Virji M.A.	223
Determination of zinc and copper in urine using Zeeman effect flame atomic absorption spectroscopy	
Liska S.K., Kerkay J., Pearson K.H.	231
Determination of zinc in whole blood, plasma and serum using Zeeman effect flame atomic absorption spectroscopy	
Liska S.K., Kerkay J., Pearson K.H.	237
A study of the microheterogeneity of transferrin in cirrhotic patients	
Poupon R.E., Papoz L., Sarmini H., Elinck R.	245
Thermal transition studies of a mature xanthelasma by differential scanning calorimetry	
Winder A.F., Bruckdorfer K.R., Fielder A.R.	253
Urinary steroid profiles in pregnancy by capillary column gas chromatography	
Hahnel E., Wilkinson S.P., Hahnel R.	259
Age and sex dependency of the biochemical indices of bone remodelling	
Stepan J.J., Tesarova A., Havranek T. et al.	273
Estimation of VLDL cholesterol in hyperlipidemia	
Wilson P.W.F., Zech L.A., Gregg R.E. et al.	285
Benzoylalanine: Detection and identification of an alanine conjugate with benzoic acid in hyperammonemic patients treated with sodium benzoate	
Shinka T., Inoue Y., Kuhara T. et al.	293
Interference by sulfonylurea drugs with the Jaffe method for creatinine	
Roach N.A., Kroll M.H., Elin R.J.	301
Determination of seminal fructose using D-fructose dehydrogenase	
Nakashima K., Takei H., Adachi O. et al.	307
A rapid, simple method for the micro-estimation of purine nucleoside phosphorylase activity in peripheral blood lymphocytes	
Renouf J.A., Thong Y.H., Chalmers A.H.	311
A non-competitive enzyme-linked immunosorbent assay for measuring human plasma apolipoprotein B levels	
James R.W., Ferrer M., Pometta D.	317

Vol. 152/1-2, 1985

A simple spectrophotometric method for quantitative fecal carbohydrate measurement Ameen V.Z., Powell G.K.	3
Failure of ultracentrifugation as a means of separating plasma free insulin from immunoglobulin fraction prior to radioimmunoassay Escobar L.J., White N.H., Santiago J.V., Gingerich R.L.	11
Comparison of the peptide and saccharide moieties of γ -glutamyltransferase isolated from neoplastic and non-neoplastic human liver tissue Tsuchida S., Yamazaki T., Camba E.M. et al.	17
Immunoelectrophoresis of lipoprotein lipase from cultured skin fibroblasts in patients with combined α -ketoacid dehydrogenase deficiency Otulakowski G., Nyhan W., Sweetman L., Robinson B.H.	27
Platelet-associated proteins in human breast cyst fluids Miller W.R., Dawes J.	37
Effect of nifedipine in hypothermic cardioplegia: A phosphorus-31 nuclear magnetic resonance study Bernard M., Menasche P., Fontanarava E. et al.	43
Quantitative determination of phosphatidylglycerol in amniotic fluid by enzymatic assay Farquharson J., Jamieson E.C., Paton R.D. et al.	55
Liquid chromatographic determination of inhibition of monoamine oxidase activity in platelet rich plasma of depressed patients treated with phenelzine Gupta R.N., Steiner M.	63
Changes in human plasma α -d-mannosidase activity during pregnancy Hirani S., Winchester B.	71
Level of plasma fibronectin in patients with peripheral vascular disease Dionysiou-Asteriou A., Karavassilis J.	79
An enzyme-amplified monoclonal immunoenzymometric assay for prostatic acid phosphatase Moss D.W., Self C.H., Whitaker K.B. et al.	85
Copper dependent control of the enzymic and phagocyte induced degradation of some biopolymers, a possible link to ischemic inflammation Hartmann H.J., Gartner A., Weser U.	95
Characterisation of high affinity binding sites of androgens in primary hepatocellular carcinoma Wilkinson M.L., Iqbal M.J., Williams R.	105
Bile acid therapies applied to patients suffering from cerebrotendinous xanthomatosis Koopman B.J., Wolthers B.G., Van der Molen J.C., Waterreus R.J.	115
Semi-automated determination of chromium in whole blood and serum by Zeeman electrothermal atomic absorption spectrophotometry Schermaier A.J., O'Connor L.H., Pearson K.H.	123
Mass screening for glucose-6-phosphate dehydrogenase deficiency: Improved fluorescent spot test Solem E., Pirzer C., Siegel M. et al.	135
Comparison of a cytosol radioreceptor assay with a radioimmunoassay for 1,25-dihydroxyvitamin D in serum or plasma De Leenheer A.P., Bauwens R.M.	143
The concentration of blood components related to fuel metabolism during prolonged fasting in children Lamers K.J.B., Doesburg W.H., Gabreels F.J.M. et al.	155
Serum ferritin in type I diabetes Carenini A., Ronchi D., Garbetta F. et al.	165
Decreased serum β -d-mannosidase activity in diabetic patients, in comparison with other glycosidases Bernard M., Brochet C., Percheron F.	171
A qualitative agar gel immunoprecipitin (IP) test for detection of fecal occult human hemoglobin Kim Y.D., Nolan J.M., Malkin A. et al.	175
Determination of free and total 3-methoxy-4-hydroxyphenylethylene glycol in human plasma by high-performance liquid chromatography with electrochemical detection Semba J.I., Watanabe A., Takahashi R.	185

Age- and sex-related differences in urinary excretion of 6 β -hydroxycortisol in humans Nakamura J., Yakata M.	193
Effect of differences in saturation sensitivity of phospholipid stains on clinical predictivity of L/S ratios Spillman T., Cotton D.B., Gonik B.	199
Determination of glutamine and glutamate in plasma of men and women by ion exchange chromatography Bos K.D., Slump P.	205
Evaluation of a commercial reagent for precipitating human serum low-density lipoprotein Shaikh M., Miller N.E.	213
Mevalonic aciduria: An inborn error of cholesterol biosynthesis? Berger R., Smit G.P.A., Schierbeek H. et al.	219
A radiochemical microassay of S-adenosylhomocysteine hydrolase in lymphoid cells Dinjens W.N.M., Chiba P., Van Laarhoven J.P.R.M. et al.	223
Determination of individual human faecal bile acids by gas-liquid chromatography after enzymatic deconjugation and simultaneous solvolysis and methylation using dimethoxypropane Van Faassen A., Nagengast F.M., Hectors M. et al.	231

Vol. 152/3, 1985

Sympathoadrenal activity and plasma glucose effects on plasma dopamine-beta-hydroxylase levels in rats Munoz J.A., Garcia-Estan J., Salom M.G. et al.	243
Octanoylglucuronide excretion in patients with a defective oxidation of medium-chain fatty acids Duran M., Ketting D., Van Vossen R. et al.	253
Plasma α_1 -antichymotrypsin in liver disease Lindmark B.E., Eriksson S.G.	261
Concentrations of succinylacetone after homogentisate and tyrosine loading in healthy individuals with low fumarylacetoacetase activity Kvittingen E.A., Leonard J.V., Pettit B.R., King G.S.	271
Effect of alcohol intake on high and low density lipoprotein metabolism in healthy volunteers Malmendier C.L., Delcroix C.	281
Platelet monoamine oxidase: Specific activity and turnover number in schizophrenics and their families Summers K.M., Andrew B., Gillespie C. et al.	289
Serum unconjugated bile acids: Qualitative and quantitative profiles in ileal resection and bacterial overgrowth Setchell K.D.R., Harrison D.L., Gilbert J.M., Mupthy G.M.	297
A microassay for acid β -galactosidase activity toward asialofetuin Mutoh T., Naoi M., Sobue I. et al.	307
Kinetic studies on neurotoxin-binding inhibitory antibodies in myasthenia gravis Ito A., Furukawa S., Satoyoshi E., Sakurai H.	315
Limitations of new thyroid function tests in pregnancy Gow S.M., Kellett H.A., Seth J. et al.	325
The determination of plasma oxalate concentrations using an enzyme/bioluminescent assay Parkinson I.S., Kealey T., Laker M.F.	335
The determination of osteopetrotic phenotypes by selective inactivation of red cell carbonic anhydrase isoenzymes Conroy C.W., Maren T.H.	347
An improved method for the simultaneous analysis of normal and modified urinary nucleosides and nucleobases by high performance liquid chromatography Sander G., Wieland J., Topp H. et al.	355
The effect of laser deterioration of nephelometer performance Salkie M.L.	363

Vol. 153/1, 1985

Study of a myoglobin test in patients hospitalized for suspected myocardial infarction Mainard F., Massoubre B., Le Marec H., Madec Y.	1
Raised C-reactive protein levels in serum from smokers Das I.	9
Crude spectrin extraction from reticulocyte-rich blood samples Marik T., Kodicek M., Brabec V.	15
Differential investigation of the capacity of succinate oxidation in human skeletal muscle Fischer J.C., Ruitenbeek W., Berden J.A. et al.	23
Simple enzymatic method for plasma branched-chain amino acids Burrin J.M., Paterson J.L., Hall G.M.	37
Improved high performance liquid chromatographic determination of thiamine diphosphate in erythrocytes Baines M.	43
Liquid chromatographic analysis of 4-hydroxy-3-methoxymandelic acid (VMA) and 4-hydroxy-3-methoxyphenylacetic acid (HVA) in urine Fenech C., Guyon F., Dauchy P.	49
Determination of urinary lysozyme during aminoglycoside therapy — Appraisal of lysozyme inhibition caused by gentamicin and tobramycin Morsky P., Ylitalo P., Koivula T.	55
Sensitive determination of cerebrospinal fluid pyruvate, lactate and glucose concentrations Bozimowski D., Artiss J.D., Zak B.	63

Vol. 153/2, 1985

Induction of cytochromes P-450 in pancreatic disease: Consequence, coincidence or cause? Acheson D.W.K., Rose P., Houston J.B., Braganza J.M.	73
Rapid method for measuring physiological concentrations of 25-hydroxyvitamin D levels in blood serum Proszynska K., Lukaszewicz J., Jarocewicz N., Lorenc R.S.	85
A sensitive sandwich enzyme immunoassay for human myoglobin using Fab'-horseradish peroxidase conjugate: Methods and results in normal subjects and patients with various diseases Nishida Y., Kawai H., Nishino H.	93
The effect of [¹²⁵ I]thyroglobulin tracer heterogeneity on serum Tg RIA measurement Spencer C.A., Platler B.W., Nicoloff J.T.	105
Vitamin E sufficiency in children with cholestasis: A comparison between erythrocyte peroxide hemolysis and serum α -tocopherol Clark J.H., Nagamori K.E., Ellett M.L., Fitzgerald J.F.	117
Fast separation of the three main plasma lipoprotein classes by ultracentrifugation using vertical rotor and multiple discontinuous density gradient Sclavons M.M., Cordonnier C.M., Mailleux P.M. et al.	125
Discrimination of phenylketonurics from persistent hyperphenylalaninemia patients using a simple phenylalanine loading test Blitzer M.G., Bailey-Wilson J.E., Shapira E.	137
Serum angiotensin converting enzyme activity in patients with psoriasis Ryder K.W., Epinette W.W., Jay S.J. et al.	143

Vol. 153/3, 1985

Detection of Bence-Jones protein by isoelectricfocussing of unconcentrated urine followed by nitrocellulose blotting and immunoperoxidase staining Norden A.G.W., Fulcher L.M., Flynn F.V.	149
---	-----

Microviscosity of tracheal aspirates in newborns with respiratory distress	
Motta C., Touillon A.M., Labbe A., Dastugue B.	157
Some aspects of the glycoprotein and glycopolypeptide content of human gastric mucus	
Ene M.D., Clamp J.R.	165
The effect of proteolytic degradation of plasma fibronectin on the responses of functional and immunometric assays for intact fibronectin	
Cotton G., Brown R.A.	173
Creatine kinase: Purification, characterization, and tissue distribution of the MM subtypes	
Mickelson J.K., Carlson C.J., Kaysen G.A., Rapaport E.	181
An automated determination of urinary 4-hydroxy-3-methoxymandelic acid	
Kodama K., Nakata T., Yamanaka K., Aoyama M.	191
Assay of platelet monoamine oxidase in whole blood	
Van Kempen G.M.J., Van Brussel J.L., Pennings Ed. J.M.	197
Increased plasma glycosidase and protease activity in uraemia: Possible role in the aetiology of the anaemia of chronic renal failure	
Shannon J.S., Lappin T.R.J., Elder G.E. et al.	203
Pancreatic secretory trypsin inhibitor-like immunoreactivity in pancreatectomized patients	
Halila H., Huhtala M.-L., Schroder T. et al.	209
Purification and determination of the modifying protein responsible for the post-synthetic modification of creatine kinase (EC 2.7.3.2) and enolase (EC 4.2.1.11)	
Van Landeghem A.A.J., Soons J.B.J., Wever R.A. et al.	217
A reproducible procedure for measuring sodium transport in cultured human fibroblasts from normal and obese donors	
Spurlock G., Morgan K., Mir M.A.	225
Normal pyruvate oxidation in Friedreich ataxia and Charcot-Marie-Tooth disease fibroblasts	
Wilson W.G.	233
A simple procedure for the routine determination of aspartate aminotransferase and alanine aminotransferase with pyridoxal phosphate	
Javier Gella F., Olivella T., Cruz Pastor M. et al.	241
Plasma vitamin A analysis by liquid chromatography: Probuco interference	
Shephard G.S., Hough B.J., Van Stuijvenberg M.E., Labadarios D.	249
The determination of methaemoglobin levels by first derivative spectroscopy	
Heales S.J.R., Bennett J.	253

Vol. 154/1, 1986

Distribution of serum zinc between albumin and α_2 -macroglobulin in patients with different zinc metabolic disorders	
Kiellerich S., Christiansen C.	1
A simple quantitative micro-method of arginase assay in blood spots dried on filter paper	
Konarska L., Tomaszewski L.	7
Ribonuclease from cytosolic fraction of human erythrocytes	
Czajkowska B., Naskalski J.W., Sznajd J.	19
Quantitation of apolipoprotein B by polyclonal and monoclonal antibodies	
Riesen W.F., Sturzenegger E., Imhof C., Mordasini R.	29
Gilbert's syndrome: Diagnosis by typical serum bilirubin pattern	
Sieg A., Stiehl A., Raedsch R. et al.	41
Measurement of plasma oxalate in healthy subjects and in patients with chronic renal failure using immobilised oxalate oxidase	
Kasidas G.P., Rose G.A.	49
Inhibitory activity of whole urine: A comparison of urines from stone formers and healthy subjects	
Ryall R.L., Hibberd C.M., Mazzachi B., Marshall V.R.	59
Serum angiotensin converting enzyme activity in cigarette smokers	
Haboubi N.A.A., Bignell A.H.C., Haboubi N.Y.	69

Vol. 154/2, 1986

Reduced trypsin binding capacity of α_2 -macroglobulin in diabetes Roberts R.C., Hall P.K., Nikolai T.F., McKenzie A.K.	85
Impaired amylase activities caused by binding of abnormal immunoglobulin A in patients with macroamylasemia Adachi K., Suzuki K., Ohno Y., Sato B.	103
Development of a radioimmunoassay for human deoxyribonuclease I Miyauchi K., Ogawa M., Shibata T. et al.	115
Urinary pseudouridine as a tumor marker in patients with small cell lung cancer Tamura S., Fujii J., Nakano T. et al.	125
Discrimination of gamma-glutamyltranspeptidase from normal and carcinomatous pancreas Yamaguchi N., Kawai K., Ashihara T.	133
Clarifying lipemic samples with an air-driven ultracentrifuge for determination of high-density lipoprotein cholesterol Narahari Reddy M., Fitzsimons E.J.	141
Prenatal diagnosis of GM ₂ gangliosidosis using a fluorogenic sulfated substrate Inui K., Wenger D.A., Furukawa M. et al.	145

Vol. 154/3, 1986

Diagnosis and characterization of GM 2 gangliosidosis type II (Sandhoff disease) by analysis of the accumulating N-acetyl-glucosaminyl oligosaccharides with high performance liquid chromatography Warner T.G., De Kramer R.D., Applegarth D., Mock A.K.	151
Aminoterminal procollagen III peptide elevation in alcoholics who are selenium and vitamin E deficient Lu W., Bantok I., Desai S. et al.	165
The analysis of data from breast cancer estrogen and progesterone receptor assays: Scatchard plots are inferior to direct fitting by computer Cooke R.R., McIntosh J.E.A.	171
Source of increased serum aspartate and alanine aminotransferase: cycloheximide effect on carbon tetrachloride hepatotoxicity Pappas Jr. N.J.	181
Lipid peroxides in blood plasma and enzymatic antioxidative defence of erythrocytes in Down's syndrome Kedziora J., Bartosz G., Gromadzinska J. et al.	191
Use of selective media for distinguishing variant forms of hypoxanthine phosphoribosyl transferase Page T.M., Broock R.L., Nyhan W.L., Hernandez N.L.	195
Highly sensitive enzyme immunoassay for human brain aldolase C Haimoto H., Kato K.	203
Reexamination of the conditions for processing and storing blood for plasma renin assay Matsunaga M., Suzuki Y., Nakagawa K. et al.	213
Transglutaminase in erythrocytes of cystic fibrosis patients Kamoun P.P., Droin V.	219
Determination of lithium in blood serum — A comparison between molecular absorption spectrophotometry and emission flame spectrometry Popov P., Otruba V., Sommer L.	223
Detection of lactyl lactate in the urine of patients receiving Hicaliq infusion Maeda K., Ohki T., Hosoi M. et al.	227
Interference by monoclonal proteins in the o-phthalaldehyde method for blood urea nitrogen Pierce G.F., Garrett N.C., Koenig J. et al.	233
Solvolysis of mono- and disulfated bile acids by ethereal HCl, prepared by a simple technique Dommes P., Breuer N.F., Goebell H.	237

The effect of storage conditions on the stability of carbamazepine and carbamazepine-10,11-epoxide in plasma Mendez-Alvarez E., Soto-Otero R., Sierra-Marcuno G.	243
Bilirubin interference in the colorimetric assay of plasma vitamin E Desai I.D., Martinez F.E.	247

Vol. 155/1, 1986

Isoforms of creatine kinase MM and MB in acute myocardial infarction: A clinical evaluation Panteghini M., Cuccia C., Malchiodi A. et al.	1
Urinary neopterin, a marker of clinical activity in patients with Crohn's disease Prior Ch., Bollbach R., Fuchs D. et al.	11
Differential measurement of monomeric and polymeric IgA by one-directional immunoelectrodif-fusion Meillet D., Tallet F., Gobert J.G. et al.	23
Purification of vitamin D binding protein from human plasma using high performance liquid chromatography Taylor G.A., Mazhindu H.N., Findlay J.B.C., Peacock M.	31
Changes in renal enzyme activities following the administration of gentamicin to unilateral nephrectomized rats Avidan G., Aladjem M., Israeli B., Bogin E.	43
Pregnancy-associated alpha ₂ -glycoprotein (α_2 -PAG): Development of a sensitive enzyme-linked immunoassay and comparison of serum concentrations in adults and children Gerrie L.M., Armstrong S.S., Horne C.H.W.	51
Preparation of ¹²⁵ I-labelled arginine vasopressin for radioimmunoassay Sadler W.A., Wright C.P., Livesey J.H.	61
Measurement of beta-2-microglobulin in serum and plasma by an enzyme-linked immunosorbent assay (ELISA) Bjerrum O.W., Birgens H.S.	69
Zinc in hair and urine of paediatric patients Van Wouwe J.P., De Wolff F.A., Van Gelderen H.H.	77
Bromocresol purple method for serum albumin gives falsely low values in patients with renal insuffi-ciency Maguire G.A., Price C.P.	83
Reliability of immunoglobulin determination by two methods Mencke M.W., Weicker H., Wolf G.K. et al.	89
Assay of free and total fatty acids (as 2-nitrophenylhydrazides) by high performance liquid chromatography Miwa H., Yamamoto M., Nishida T.	95

Vol. 155/2, 1986

Involvement of thiol proteases in galactosialidosis Takeda E., Kuroda Y., Toshima K. et al.	109
Occurrence of short aliphatic diols in human blood: Identification by gas chromatography-mass spectrometry Sisfontes L., Nyborg G., Jones A.W., Blomstrand R.	117
Patients with coeliac disease have a characteristic gliadin antibody pattern Friis S.U., Noren O., Sjostrom H., Gudmand-Hoyer E.	133
Determination in human cerebrospinal fluid of glial fibrillary acidic protein, S-100 and myelin basic protein as indices of non-specific or specific central nervous tissue pathology Noppe M., Crols R., Andries D., Lowenthal A.	143
A turbidimetric immunochemical method for determination of serum β_2 -microglobulin using a centrifugal analyzer Evrin P.E., Jansson N.O., Bergdahl A.	151

Pulsatile pattern of thyrotropin-release in normal men Brabant G., Ranft U., Ocran K. et al.	159
CA 125 (ovarian tumour-associated antigen) in ascitic liver diseases Bergmann J.F., Beaugrand M., Labadie H. et al.	163
Analysis of volatile free fatty acids in human urine by capillary column gas chromatography/mass spectrometry Maltby D.A., Millington D.S.	167
Rapid detection of argininosuccinic aciduria and citrullinuria by fast atom bombardment and tandem mass spectrometry Millington D.S., Maltby D.A., Roe C.R.	173
Zinc concentrations in mononuclear and polymorphonuclear leukocytes Purcell S.K., Hambidge K.M., Jacobs M.A.	179
Comparison of zinc contents in human serum and plasma Chen Wei-jie, Zhao Cheng-yi, Zheng Tian-li	185
Deficient beta-ketothiolase activity in leukocytes from a patient with 2-methylacetoacetic aciduria Hiyama K., Sakura N., Matsumoto T., Kuhara T.	189

Vol. 155/3, 1986

Increase of plasma and red cell cholesterol sulfate levels in induced hypothyroidism in man Van Doormaal J.J., Muskiet F.A.J., Jansen G. et al.	195
Peroxidase activity and estradiol receptors in human breast cancer Gilabert B.P., Zamora V.A., Oriola A.P. et al.	201
An enzyme mediated, colorimetric method for the measurement of salicylate Chubb S.A.P., Campbell R.S., Ramsay J.R. et al.	209
A double antibody radioimmunoassay for human α_1 -microglobulin Itoh Y., Nishino H., Enomoto H., Kawai T.	221
Glycosaminoglycans of skin and urine in pseudoxanthoma elasticum: Evidence for chondroitin 6-sulfate alteration Longas M.O., Wiach P., Lebwohl M.G., Fleischmajer R.	227
Sweat sulfate concentrations are decreased in cystic fibrosis Cole D.E.C., Landry D.A., Boucher M.J., Gillespie C.T.	237
Rapid diagnosis of vitamin D-dependent rickets type II by use of phytohemagglutinin-stimulated lymphocytes Takeda E., Kuroda Y., Saijo T. et al.	245
Determination of mitochondrial aspartate aminotransferase in serum Hirano K., Matsuda K., Adachi T. et al.	251
Estimation of NADH oxidation in human skeletal muscle mitochondria Fischer J.C., Ruitenbeek W., Trijbels J.M.F. et al.	263
The relationship between plasma triglycerides, cholesterol, total lipids and lipid peroxidation products during human atherosclerosis Ledwozyw A., Michalak J., Stepień A., Kadziolka A.	275
Affinity electrophoresis of human serum alkaline phosphatase isoenzymes in agarose gel containing lectin Onica D., Sundblad L., Waldenlind L.	285
An O_2 -based enzyme electrode for whole blood lactate measurement under continuous flow conditions Weaver M.R., Vadgama P.M.	295
Fatty acyl composition of the major lipid classes of HDL ₂ and HDL ₃ of human serum Devitt M.M., Headon D.R.	309
The effects of arginine, dextran and Haemacel infusions on urinary albumin, β_2 -microglobulin and N-acetyl- β -D-glucosaminidase O'Reilly St. D.J., Parry E.S., Whicher J.T.	319
Tissue origin of serum placental-like alkaline phosphatase in cigarette smokers Williams G.H., McLaughlin P.J., Johnson P.M.	329

Effect of ultraviolet light on creatinine measurement in jaundiced specimens Nisbet J.A., D'Souza R.	335
---	-----

Vol. 156/1, 1986

Oncogenes	
Garrett C.T.	1
Ganglioside loading of cultured fibroblasts: A provocative method for the diagnosis of the G(M2) gangliosidoses	
Charrow J., Binns H.J.	41
Development of a sensitive enzyme immunoassay for human epidermal growth factor (Urogastrone)	
Kurobe M., Tokida N., Furukawa S. et al.	51
Energy metabolism in adenosine deaminase-inhibited human erythrocytes	
Buc H.A., Thuillier L., Hamet M. et al.	61
Determination of oxalate in urine by flow injection analysis	
Gaetani E., Laureri C.F., Vitto M. et al.	71
Diuresis-dependent excretion of multiple forms of renal brush-border enzymes in urine	
Jung K., Schulze G.	77
The analysis of thiodiglycolic acid by selected ion monitoring	
Pettit B.R.	85
HPLC assay of phenylalanine and tyrosine in blood spots on filter paper	
Roesel R.A., Blankenship P.R., Hommes F.A.	91
A simple enzymatic fluorimetric method for the determination of triglycerides in 10 µl of serum	
Gleeson M., Maughan R.J.	97
Blood plasma fluoride in haemodialysed patients	
Chaleil D., Simon P., Tessier B. et al.	105
Determination of silicon, calcium, magnesium and phosphorus in urine using inductively-coupled plasma emission spectrometry and a matrix-matching technique	
Tanaka T., Hayashi Y.	109
Measurement of radioactivity of pyruvate and other 2-oxo-acids labelled at the 1-position	
Van Laack H.L.J.M., Janssen A.J.M., Ruitenbeek W. et al.	115

Vol. 156/2, 1986

Pancreatic secretory trypsin inhibitor from human amniotic fluid and fetal and neonatal urine: Concentrations and physicochemical characterization	
Kolho K.-L., Huhtala M.-L., Jalanko H. et al.	123
Quantification of human serum apolipoprotein A-I by zone immunoelectrophoresis assay and a procedure for the preparation of an A-I standard	
Holmquist L., Vesterberg O.	131
Quantitation of CK-MB release: Diagnostic utility in coronary artery bypass grafting	
Devine J.E., Wiens R.D., Halstead J.M., Codd J.E.	145
On-line clarification for the measurement of serum glucose in hyperlipidemic specimens	
Artiss J.D., Strandbergh D.R., Sharma A., Zak B.	151
Myocardial chondroitin sulfates D and E in a case of acute carbon monoxide poisoning	
Ohishi H., Binette J.P., Schmid K.	157
Probe of the (Ca ²⁺ + Mg ²⁺)-ATPase in erythrocyte membranes of cystic fibrosis patients	
Hunsinger R.N., Cheung H.C.	165
Sulfogalactocerebroside and bis-(monoacylglyceryl)-phosphate as activators of spleen glucocerebrosidase	
Prencz E.M., Garrett K.O., Panitch H. et al.	179
Deficient activities and proteins of peroxisomal β -oxidation enzymes in infants with Zellweger syndrome	
Suzuki Y., Orii T., Mori M. et al.	191

The measurement of glycosylated albumin by reduction of alkaline nitro-blue tetrazolium Walker S.W., Howie A.F., Smith A.F.	197
Determination of volatile fatty acids by gas chromatography on a capillary and a megabore column Duisterwinkel F.J., Wolthers B.G., Van Der Slik W., Dankert J.	207
Use of pooled human serum in the standardization process of the serum fructosamine determination for the estimation of glycosylated serum proteins Smid E., Ferencz A., Fodor M.	215
Reduced imprecision of the radioenzymatic assay of plasma catecholamines by improving the stability of the internal standards Hoffmann J.J.M.L., Willemsen J.J., Lenders J.W.M., Benraad Th. J.	221
What information should be provided in the package inserts of commercial reference materials in clinical chemistry? Broughton P.M.G.	227

Vol. 156/3, 1986

Enzymatic assessment of myocardial injury after infarction or cardiac surgery. Is isoenzyme analysis required? Hermens Th. W., Willems G.M., Davids H.A. et al.	235
Quality control of azure B preparations by liquid chromatography and standardization with azure B tetrafluoroborate Nelis H.J.C.F., De Leenheer A.P.	247
Sensitive quantitation of carcinoembryonic antigen in cerebrospinal fluid and its barrier-dependent differentiation Reiber H., Jacobi C., Felgenhauer K.	259
Assessment of the AminoStat-FLM immunoagglutination test for phosphatidylglycerol in amniotic fluid Farquharson J., Jamieson E.C., Berry E. et al.	271
Diagnosis of inherited adenylosuccinase deficiency by thin-layer chromatography of urinary imidazoles and by automated cation exchange column chromatography of purines De Bree P.K., Wadman S.K., Duran M., Fabry de Jonge H.	279
ELISA for human proinsulin Hartling S.G., Dinesen B., Kappelgard A.-M. et al.	289
Changes in fatty acid profiles of plasma, erythrocytes and polymorphonuclear leukocytes in induced hypothyroidism in man: Indirect evidence for altered $\Delta 6$ desaturase activity Van Doormaal J.J., Muskiet F.A.J., Martini I.A., Doorenbos H.	299
Increase in blood plasma levels of boron and strontium in hemodialyzed patients Mauras Y., Ang K.S., Simon P. et al.	315
Plasma and red cell lipids in alcoholics with macrocytosis Clemens M.R., Kessler W., Schied H.W. et al.	321
Glycated haemoglobin and glycated plasma protein, a comparative study Ross I.S., Mitchell A.F., Stroud R.E.	329
An accurate, simple and rapid test for detecting elevated levels of C-reactive protein in serum by agglutination of fat emulsion Hulman G., Fuller M., Pearson H.J., Bell P.R.F.	337

Vol. 157/1, 1986

Clinical applications of gene probes in human genetic disease, malignancy, and infectious disease Lowe J.B.	1
Erythrocyte cation transport activities as a function of cell age Hentschel W.M., Wu L.L., Tobin G.O. et al.	33

A case of rheumatoid arthritis with various enzyme-immunoglobulin complexes	
Maekawa M., Sudo K., Kanno T.	45
Serum creatine kinase isoenzyme MB concentration after endomyocardial biopsy	
Chapelle J.P., El Allaf D., El Allaf M. et al.	55
Oligomerization of pregnancy-specific β_1 -glycoprotein (SP ₁) at physiologic pH and ionic strength	
Rosen S.W., Calvert I., Lee N. et al.	65
Fluorometric determination of nanogram quantities of protein in small samples: Application to calcium-transport adenosine triphosphatase	
Bridges M.A., McErlane K.M., Kwong E. et al.	73
The influence of serum and serum proteins on calcium oxalate crystal growth and aggregation	
Edyvane K.A., Ryall R.L., Marshall V.R.	81
Isolation and characterisation of lactoferrin separated from human whey by adsorption chromatography using Cibacron Blue F3G-A linked affinity adsorbent	
Bezwoza W.R., Mansoor N.	89
A procedure for large scale purification of human plasma amyloid P component	
Ohkubo I., Sahashi W., Namikawa C. et al.	95
Colorimetric assay for carboxypeptidase N in serum	
Hendriks D., Van Sande M., Scharpe S.	103
Lysosomal enzymes in chorionic villi, cultured amniocytes, and cultured skin fibroblasts	
Evans M.I., Moore C., Kolodny E.H. et al.	109
Estimation of aluminoxamine and ferrioxamine in plasma by high performance liquid chromatography	
Hughes H., Hagen L.E., Cameron E.C., Sutton R.A.L.	115
An improved spectrophotometric assay of urinary metanephries	
Crawford G.A., Gallery E.D.M., Gyory A.Z.	121

Vol. 157/2, 1986

IgD plasma cell leukemia associated with pyroglobulinemia and pyroglobulinuria	
Gastearena J., Fernandez F.J., Orue M.T. et al.	133
Guanidino compounds in uraemic dialysed patients	
De Deyn P., Marescau B., Lornoy W. et al.	143
Evaluation and comparison of cathodic trypsin-like immunoreactivity, pancreatic lipase and pancreatic isomylase in the diagnosis of acute pancreatitis in 849 consecutive patients with acute abdominal pain	
Moller-Petersen J., Klaerke M., Dati F.	151
Gold staining in cellulose acetate membranes	
Righetti P.G., Casero P., Del Campo G.B.	167
Estimation of vitamin D ₃ and 25-hydroxyvitamin D ₃ in muscle and adipose tissue of rats and man	
Lawson D.E.M., Douglas J., Lean M., Sedrani S.	175
Internalization of exogenous gangliosides in cultured skin fibroblasts for the diagnosis of mucopolipidosis IV	
Zeigler M., Bach G.	183
Enzyme electrode for the measurement of lactate in undiluted blood	
Mullen W.H., Churchouse S.J., Keedy F.H., Vadgama P.M.	191
Serum calcium status in health and disease: A comparison of measured and derived parameters	
White T.F., Farndon J.R., Conceicao S.C. et al.	199
Comparison of sampling methods for the analysis of methane in expired air	
Peled Y., Gilat T.	215

Vol. 157/3, 1986

Quantitation of trans-vitamin K1 in small serum samples by off-line multidimensional liquid chromatography	
Van Haard P.M.M., Engel R., Pietersma-de Bruyn A.L.J.M.	221

Enzyme immunoassay of serum cortisol using a new transferable needle lid technique Dhar T.K., Voss E., Schoneshofer M.	231
Preliminary report on a case of apolipoproteins C1 and CII deficiency Dumon M.-F., Clerc M.	239
Separation of apolipoprotein B subfractions by high performance gel permeation chromatography Schwandt P., Richter W.O., Weisweiler P.	249
Metabolic conversion of L-[U- ¹⁴ C]phenylalanine to respiratory ¹⁴ CO ₂ in healthy subjects, phenylketonuria heterozygotes and classic phenylketonurics Lehmann W.D., Fischer R., Heinrich H.C. et al.	253
ELISA of 6-beta-hydroxycortisol in human urine: Diurnal variations and effects of antiepileptic therapy Zhiri A., Wellman-Bednawska M., Siest G.	267
Capillary gas chromatographic profiling of urinary, plasma and erythrocyte sugars and polyols as their trimethylsilyl derivatives, preceded by a simple and rapid prepurification method Jansen G., Muskiet F.A.J., Schierbeek H. et al.	277
Diagnostic value of microalbuminuria in pre-eclampsia Irgens-Moller L., Hemmingsen L., Holm J.	295
A non-random distribution of transferrin iron in fresh human sera Van Eijk H.G., Van Noort W.L.	299
The reliability of the use of para toluene sulfonic acid for simultaneous hydrolysis and quantitation of both N-acetyl-glucosamine and amino acids in human transferrins Van Eijk H.G., Van Noort W.L.	305
Multiwavelength analyses of derivative spectra: An application for the determination of urinary porphyrins Walters M., Thornton D.J.	311

Vol. 158/1, 1986

Normal human serum contains two forms of acetylcholinesterase Sorensen K., Brodbeck U., Rasmussen A.G., Norgaard-Pedersen B.	1
High molecular weight angiotensinogen: A pregnancy associated protein Tewksbury D.A., Tryon E.S., Burrill R.E., Dart R.A.	7
A map of urine proteins based on one-dimensional SDS-polyacrylamide gel electrophoresis and Western blotting using one microliter of unconcentrated urine Kshirsagar B., Wiggins R.C.	13
A modified radioimmunoassay for arylsulfatase A in human serum and urine Laidler P.M., Silberring J., Van Etten R.L.	23
Measurement of total HDL, HDL ₂ and HDL ₃ by dextran-sulfate-MgCl ₂ precipitation technique in human serum Talameh Y., Wei R., Naito H.	33
Changes in plasma osteocalcin concentration following treatment with stanozolol Couch M., Preston F.E., Malia R.G. et al.	43
An improved approach to thyroid function testing in patients with non-thyroidal illness Gow S.M., Elder A., Caldwell G. et al.	49
Development and evaluation of a simple, direct, solid-phase radioimmunoassay of serum cortisol from readily available reagents McConway M.G., Chapman R.S.	59
Lipid associated tissue ferritin Cham B.E., Roeser P., Nikles A.C., Ridgway K.M.	71
Physiological concentrations of human epidermal growth factor in biological fluids: Use of a sensitive enzyme immunoassay Joh R., Itoh M., Katsumi K. et al.	81
Acid stable trypsin inhibitor in bile Yamamoto T., Sumi H., Maruyama M. et al.	91
Highly sensitive enzyme immunoassay for human creatine kinase MM and MB isozymes Kato K., Shimizu A.	99

Excretion of octopamine metabolites in neuroblastoma	
Couch M.W., Greer D.M., Williams C.M.	109
Improved assay of the enzymes of glutathione synthesis: γ -glutamylcysteine synthetase and glutathione synthetase	
Beutler E., Gelbart T.	115

Vol. 158/2, 1986

Determination of ionised calcium by ion selective electrode is not independent of albumin concentration	
Freaney R., Egan T., McKenna M.J. et al.	129
Determination of urinary estradiol using an enzymatic method during the menstrual cycle. Comparison with a method using isotope dilution-mass spectrometry (ID-MS)	
Patricot M.C., Mathian B., Serpente S., Revol A.	139
Identification of lignans and phytoestrogens in urine of chimpanzees	
Adlercreutz H., Musey P.I., Fotsis T. et al.	147
Deficient activity of glucocerebrosidase in urine from patients with type I Gaucher disease	
Aerts J.M.F.G., Donker-Koopman W.E., Koot M. et al.	155
Comparison of a tumour-derived form of intestinal alkaline phosphatase with foetal and adult intestinal alkaline phosphatases	
Moss D.W., Parmar C.R., Whitaker K.B.	165
Assay of urinary 5-hydroxyindole-3-acetic acid: Two methodologies compared	
Baars J.D., Van Haard P.M.M., Lombarts A.J.P.F.	173
Detection of carriers of cerebrotendinous xanthomatosis	
Koopman B.J., Waterreus R.J., Van Den Brekel H.W.C., Wolthers B.G.	179
The effect of sotalol on plasma lipoproteins and apolipoproteins	
Northcote R.J., Packard C.J., Ballantyne D.	187
Tissue homogenization using a micro-dismembrator for the measurement of enzyme activities	
Peters G.J., Laurensse E.J., Leyva A., Pinedo H.M.	193
An improved source receptor for 1,25-dihydroxyvitamin D3 assay	
Blayau M., Leray G., Prodhomme C. et al.	199

Vol. 158/3, 1986

Chemotherapeutic agent cisplatin monitoring in biological fluids by means of inductively-coupled plasma emission spectrometry (ICP-AES)	
Dominici C., Alimonti A., Caroli S. et al.	207
N-Acetylglucosaminiduria in a child with a progressive cerebral atrophy	
Kvittingen E.A., Guldal G., Borsting S. et al.	217
Crystal inhibition: The effects of polyanions on calcium oxalate crystal growth	
Fellstrom B., Danielson B.G., Ljunghall S., Wikstrom B.	229
Errors in measurements of CO ₂ with the use of drying agents	
Elia M., McDonald T., Crisp A.	237
Sarcoplasmic reticulum Ca ²⁺ -ATPase and acylphosphatase activities in muscle biopsies from patients with Duchenne muscular dystrophy	
Landi N., Nassi P., Liguri G. et al.	245
Mitochondrial myopathy: Tissue-specific expression of a defect in ubiquinol-cytochrome c reductase	
Darley-Usmar V.M., Watanabe M., Uchiyama Y. et al.	253
Cholesterol efflux from cultured skin fibroblasts in 12 patients clinically heterozygous for familial hypercholesterolemia	
Esteve O., Baudet M.-F., Jacotot B.	263
Hepatic acyl-CoA:cholesterol acyltransferase. Development of a standard assay and determination in patients with cholesterol gallstones	
Smith J.L., De Jersey J., Pillary S.P., Hardie I.R.	271

Rapid assay of pyruvate with a centrifugal analyser using a commercially available test kit Van Erven A., Huisman R., Vermunt K.	283
Enzyme immunoassay for ceruloplasmin: Application to cancer patient serum DiSilvestro R.A., David E.A.	287
Radioimmunoassay for nonenzymatically glycated protein in human serum Nakayama H., Taneda S., Manda N. et al.	293

Vol. 159/1, 1986

Peroxisomal β -oxidation of palmitoyl-CoA in human liver homogenates and its deficiency in the cerebro-hepato-renal (Zellweger) syndrome Wanders R.J.A., Van Roermund C.W.T., De Vries C.T. et al.	1
1,6-Anhydro- β -D-glucopyranose (β -glucosan), a constituent of human urine Dorland L., Wadman S.K., Fabry de Jonge H., Ketting D.	11
An immunoglobulin G inhibiting lactate dehydrogenase activity Hirano T., Matsuzaki H., Miura M. et al.	17
Purification and characterization of pancreatic secretory trypsin inhibitor in human gastric mucosa Shibata T., Ogawa M., Matsuda K. et al.	27
Microheterogeneity of Kasahara isozyme Hada T., Imanishi H., Yamamoto T. et al.	37
Radioimmunoassay for atrial natriuretic peptide: Method and results in normal subjects and patients with various diseases Nishiuchi T., Saito H., Yamasaki Y., Saito S.	45
Absence of argininosuccinate lyase protein in the liver of two patients with argininosuccinic aciduria Kobayashi K., Itakura Y., Saheki T. et al.	59
The nature of gastric secretions in a patient with Zollinger-Ellison syndrome without peptic ulceration Clamp J.R., Fairclough P.D.	69
Glutathione-linked enzyme activities in red cell aging Imanishi H., Nakai T., Abe T., Takino T.	73
A simple method to diagnose adrenoleukodystrophy using a dried blood spot on filter paper Nishio H., Kodama S., Yokoyama S. et al.	77
Placental alkaline phosphatase in an omental tumor Miura M., Matsuzaki H., Sakagishi Y., Sekine T.	83

Vol. 159/2, 1986

Characteristics of antisera to antigenic forms of 1,25-dihydroxycholecalciferol Brown W.B., Peacock M.	111
Analysis of conjugated and unconjugated bile acids in serum and jejunal fluid of normal subjects Tangerman A., Van Schaik A., Van der Hoek E.W.	123
The subcellular distribution and levels of calmodulin in jejunal biopsies from control subjects and patients with coeliac disease Macpherson A.J.S., Bjarnason I., Peters T.J.	133
Measurement of human growth hormone (hGH) using a rapid immunochemiluminometric assay Weeks I., Woodhead J.S.	139
Lipoprotein and apoprotein levels in postmenopausal women during treatment with norethisterone Farish E., Fletcher C.D., Hart D.M. et al.	147
Spin-echo and 2-dimensional ^1H nuclear magnetic resonance studies on urinary metabolites from patients with 2-methylacetocetyl CoA thiolase deficiency Williams S.R., Iles R.A., Chalmers R.A.	153

Direct demonstration of an inhibitor of sodium, potassium dependent adenosine triphosphatase (Na ⁺ ,K ⁺ -ATPase) in plasma from normotensive and hypertensive subjects	163
Roulston J.E., Traill K., Wathen C.G.	
Determination of aromatic-L-amino acid decarboxylase in human plasma	173
Boomsma F., Van der Hoorn F.A.J., Schalekamp M.A.D.H.	
A new solid-phase assay for biotin and biocytin and its application to the study of patients with biotinidase deficiency	185
Ping Wai Chan, Bartlett K.	
Measurement of magnesium in serum and urine with a random access analyzer by use of a modified xylidyl blue-I procedure	197
Ratge D., Kohse K.P., Wiasser H.	
An enzyme-linked immunosorbent assay (ELISA) for urinary free cortisol	205
Lewis J.G., Manley L., Townsend J.C., Elder P.A.	
Isolation of platelets from human blood by free-flow electrophoresis	211
Wilson R.B.J., Graham J.M.	

Vol. 159/3, 1986

Creatine kinase and lactate dehydrogenase isoenzymes in stage D prostatic carcinoma	219
Strom S., Bendz R.	
Determination of pipelicolic acid in urine and plasma by isotope dilution mass fragmentography	229
Van Den Berg G.A., Breukelman H., Elzinga H. et al.	
Enzyme-linked immunosorbent assay (ELISA) for calcium-binding protein of human small intestine. A quantitative method for measuring calcium-binding protein in small intestinal biopsies	239
Staun M.	
Carnitine deficiency in haemodialysed patients	249
Rodriguez-Segade S., Del la Pena C.A., Paz et al. J.M.	
Enzyme-linked immunosorbent assay for amitriptyline and other antidepressants using a monoclonal antibody	257
Denis H., Marullo S., Hoebeke J., Strosberg A.D.	
Quantitation of human serum apolipoprotein B by enzyme immunoassay in excess antigen	269
Fievet C., Dedieu A., Kandoussi A. et al.	
β -N-Acetyl-D-glucosaminidase isoenzymes from human amnionic membranes	279
Orlacchio A., Emiliani C., Di Renzo G.C., Cosmi E.V.	
Citrate interference with the determination of acetylcarnitine: A method for its elimination	291
Cooper M.B., Forte C.A., Jones D.A.	

Vol. 160/1, 1986

Analysis of plasma lipoproteins by ultracentrifugation in a new fixed angle rotor: Evaluation of a phosphotungstic acid/MgCl ₂ precipitation and a quantitative lipoprotein electrophoresis assay	1
Marz W., Gross W.	
Clinical significance and characteristics of creatine kinase-immunoglobulin complexes in sera from patients with malignant tumors	19
Kanemitsu F.	
Sandwich electroimmunofixation (SEIF) for the assessment of isoenzyme specificities of immunoglobulins isolated from enzyme-immunoglobulin complexes	27
Kanemitsu F., Okigaki T.	
A sandwich immunoassay for human prolyl 4-hydroxylase using monoclonal antibody	37
Yoshida S., Bai Y., Muragaki Y. et al.	
Unusual trihydroxy bile acids in the urine of healthy humans	47
Nakashima T., Sano A., Seto Y. et al.	

Functional activities and nonenzymatic glycosylation of plasma proteinase inhibitors in diabetes Hall P., Tryon E., Nikolai T.F., Roberts R.C.	55
5-Amino-2-nitrobenzoic acid as a reference material for the determination of γ -GT activity Yamada S., Sekiguchi M., Kawano K.	63

Vol. 160/2, 1986

Dependence of urine composition on the age and sex of healthy subjects Hesse A., Classen A., Knoll M. et al.	79
Trends in urolithiasis in various ethnic groups and by age, in Israel Zaidman J.L., Eidelman A., Pinto N. et al.	87
Strategies and methods for the analytical investigation of urinary calculi Uldall A.	93
2,8-Dihydroxyadenine lithiasis Simmonds H.A.	103
Medical investigation and treatment of urinary stones: A search for new ideas Rose G.A.	109
A stable liquid control material for urinary enzyme assays Maruhn D., Wehling K., Metz U.	117
Fluorometric measurement of urinary N-acetyl- β -D-glucosaminidase and its correlation to uremia Linko-Lopponen S.	119
Identification of inhibitors of urinary alanine aminopeptidase Mattenheimer H., Frolke W., Grotzsch H., Simane Z.	129
Urinary enzyme analysis in renal allograft transplantation Kotanko P., Keiler R., Knabl L. et al.	137
Serum and urinary enzyme activities in renal artery embolism Donadio C., Auner I., Giordani R. et al.	145
Urinary enzyme excretion after a single dose of phenacetin and paracetamol (acetaminophen) during antidiuresis and during water diuresis Metz U., Graben N., Maruhn D., Bock K.D.	151
Increased levels of urinary adenosine deaminase binding protein in children treated with cisplatin or methotrexate Goren M.P., Wright R.K., Horowitz M.E.	157
Value of enzyme determinations in urine for the diagnosis of nephrotoxicity in rats Hofmeister R., Bhargava A.S., Gunzel P.	163
Kidney-derived urinary antigens assayed with monoclonal antibodies for the detection of renal damage Falkenberg F.W., Mai U., Puppe C. et al.	171
Biological monitoring of aluminium in renal patients De Wolff F.A., Van der Voet G.B.	183
Allopurinol in renal failure and the tumour lysis syndrome Simmonds H.A., Cameron J.S., Morris G.S., Davies P.M.	189
α (2U)-Globulin: Measurement in rat kidney and relationship to hyaline droplets Stonard M.D., Phillips P.G.N., Foster J.R. et al.	197
NADH-dependent prostaglandin E_2 -9-ketoreductase activity and prostaglandin synthesis in the Brattleboro rat kidney: Effects of the antidiuretic hormone Rathaus M., Werber M.M., Podjarny E. et al.	205
Biochemical changes in liver, kidney and blood associated with common bile duct ligation Israeli B.-A., Bogin E.	211

Vol. 160/3, 1986

In vitro studies upon the release of γ -glutamyltransferase from human liver Wenham P.R., Horn D.B., Smith A.F.	223
---	-----

The physicochemical properties and the process of pyrogel formation of IgD pyroglobulin	235
Ohno T., Furukawa H., Kanoh T., Uchino H.	
Hydrolysis of conjugated steroids by β -glucuronidase from Ampullaria and application to the determination of urinary 17-hydroxycorticosteroids	245
Tsukada T., Ise M., Yoshino M.	
Quantitative analysis of the mitochondrial cytochrome P-450-linked monooxygenase system: NADPH-hepatoredoxin reductase, hepatoredoxin, and cytochrome P-450(S27) in livers of patients with cerebrotendinous xanthomatosis	255
Miki H., Takeuchi H., Yamada A. et al.	
Renal effects of dichloroacetate in vivo	265
Kodama H., Yamaguchi S., Okabe I. et al.	
A rapid quantitative capillary tube enzyme immunoassay for human chorionic gonadotropin in urine	273
Nagainis P.A., Nakagawa C.H., Baron S.L. et al.	
Calcium activated neutral proteases (milli- and micro-CANP) and endogenous CANP inhibitor of muscle in Duchenne muscular dystrophy (DMD)	281
Reddy P.A., Anandavalli T.E., Anandaraj M.P.J.S.	
Plasma fibronectin in liver cirrhosis and its diagnostic value	289
Gabrielli G.B., Casaril M., Bonazzi L. et al.	
Mitochondrial aspartate aminotransferase linked to immunoglobulin G of the α - λ type: report of a case	297
Moriyama T., Ashiie T., Kikuri K. et al.	
Degradation of vitamin B ₆ standard solutions	307
Shephard G.S., Labadarios D.	
Diagnostic value of an immunoenzymetric assay compared with an enzymic immunoinhibition procedure for creatine kinase isoenzyme MB	313
Moerman J., Nellens P., Beckers F., Sierens L.	
Stability of creatinine, urea and uric acid in urine stored under various conditions	319
Soliman S.A., Abdel-Hay M.H., Sulaiman M.I., Tayeb O.S.	

Vol. 161/1, 1986

Determination of serum neuron-specific enolase by differential immunocapture	1
Viallard J.-L., Ven Murthy M.R., Betail G., Dastugue B.	
Measurement of 5-aminolaevulinic acid by reversed phase HPLC and fluorescence detection	11
Minder E.I.	
Impaired hepatocellular integrity during general anaesthesia, as assessed by measurement of plasma glutathione S-transferase	19
Hussey A.J., Howie J., Allan L.G. et al.	
Assay of blood acetaldehyde by HPLC with fluorescence detection of its 2-diphenylacetyl-1,3-indandione-1-azine derivative	29
Rideout J.M., Lim C.K., Peters T.J.	
Serum levels of immunoreactive PSTI in acute abdominal disorders, with special reference to a possible extrapancreatic PSTI production	37
Lasson A., Borgstrom A., Ohlsson K.	
Comparison of radioimmunoassay and immunoradiometric assay for serum prostatic acid phosphatase	47
Sturgeon C.M., Hussey A.J., Beynon L. et al.	
Human inter- α trypsin inhibitor and immunologically related inhibitors investigated by quantitative immunoelectrophoresis. I. Method and reference material	59
Byrjalsen I., Hansen-Nord G., Odum L.	
Hepatic damage in the rat following administration of thyroxine or triiodothyronine, assessed by measurement of plasma glutathione S-transferase YaYa concentrations	69
Beckett G.J., Hunter J.E., Hayes J.D.	

Very long chain fatty acids in genetic peroxisomal disease fibroblasts: Differences between the cerebro-hepato-renal (Zellweger) syndrome and adrenoleukodystrophy variants	
Molzer B., Korschinsky M., Bernheimer H. et al.	81
Effect of ileal exclusion on kinetics of very low density lipoprotein triglycerides in familial hypercholesterolemia	
Koivisto P.V.I., Miettinen T.A.	91
The effect of lead on tetrahydrobiopterin metabolism. A possible mechanism for neurotoxicity	
Eggar C., Hamon C.G.B., Morar C. et al.	103
Determination of L-asparaginase activity in serum by thin-layer chromatography: Application to the treatment of acute lymphoblastic leukemia	
Clausen N., Christensen E.	111

Vol. 161/2, 1986

Multiple forms of acid and alkaline phosphatases: Genetics, expression and tissue-specific modification	
Moss D.W.	123
Serum type III procollagen peptide as a non-invasive marker of liver damage during infancy and childhood in extrahepatic biliary atresia, idiopathic hepatitis of infancy and $\alpha 1$ antitrypsin deficiency	
Trivedi P., Cheeseman P., Portmann B., Mowat A.P.	137
Non-specific binding of insulin in an equilibrium binding assay of insulin antibodies	
Storms F.E.M.G., Lutterman J.A., Ross A.H. et al.	147
Epoxycarboxylic aciduria resulting from the ingestion of castor oil	
Hagenfeldt L., Blomquist L., Midtvedt T.	157
Apolipoprotein profile in healthy males and its relation to maximum aerobic capacity (MAC)	
Berg A., Frey I., Keul J.	165
Methylmalonic aciduria and propionic acidemia studied by proton nuclear magnetic resonance spectroscopy	
Ilea R.A., Chalmers R.A., Hind A.J.	173
Precise measurement of glycated serum albumin by column affinity chromatography and immunoturbidimetry	
Reed P., Bhatnagar D., Dhar H., Winocour P.H.	191
Characteristics of ferritins in human milk secretions: Similarities to serum and tissue isoforms	
Arosio P., Ponzone A., Ferrero R. et al.	201
Assay of human pancreatic lipase in biological fluids using a non-competitive enzyme immunoassay	
Carrere J., Galabert C., Thouvenot J.P., Figarella C.	209
Rapid diagnosis of Zellweger syndrome and infantile Refsum's disease by fast atom bombardment-mass spectrometry of urine bile salts	
Lawson A.M., Madigan M.J., Shortland D., Clayton P.T.	221
Effect of fenopropen on thyroid function tests	
Tillman J., Leask J.T.S., Campbell J. et al.	233
Autoantibodies to thyroid hormones: Association with falsely low hormone levels measured by 'Amerlex' assay	
Heyma P., Harrison L.C.	239
The influence of freezing and thawing on platelet-associated vasopressin during osmotic stimulation of the neurohypophysis	
Baylis P.H., Thompson C.J., Burd J.M.	243

Vol. 161/3, 1986

Serum concentrations of laminin and aminoterminal propeptide of type III procollagen in relation to the portal venous pressure of fibrotic liver diseases	
Gressner A.M., Tittor W., Negwer A., Pick-Kober K.-H.	249

Cerebrospinal fluid lactate dehydrogenase activities in patients with central nervous system metastases	
Van Zanten A.P., Twijnstra A., Hart A.A.M., Ongerboer De Visser B.W.	259
Substrate specificity of alkaline phosphatase from human polymorphonuclear leukocytes	
Stinson R.A., Gainer A.L., Chai J., Chan J.A.	283
Effect of parathyroid hormone and uremia on erythrocyte deformability	
Bogin E., Earon Y., Blum M.	293
Membrane lipid peroxidation in erythrocytes of the newborn	
Jain S.K.	301
Lysosomal enzyme activities in fresh frozen chorionic villi and in cultured trophoblasts	
Ben-Yoseph Y., Evans M.I., Bottoms S.F. et al.	307
Quantitation of serum lactate dehydrogenase-5 with monoclonal antibodies	
Vaidya H.C., Dietzler D.N., Ladenson J.H.	315
Fatty acyl-CoA dehydrogenase enzymes in human skeletal muscle	
Carroll J.E., McGuire B.S., Hall C.L.	327
A preliminary study of urinary transferrin as a marker for prostatic cancer	
Fernandez C., Rifai N., Wenger A.S. et al.	335
Serum adenosine deaminase activity in the differentiation of respiratory diseases in children	
Yasuhara A., Nakamura M., Shuto H., Kobayashi Y.	341
Discrimination of heterozygotes for phenylketonuria, persistent hyperphenylalaninemia and controls by phenylalanine loading	
Blitzer M.G., Bailey-Wilson J.E., Shapira E.	347

Vol. 162/1, 1987

Carrier detection in Duchenne muscular dystrophy: A review of current issues and approaches	
Grüemer H.-D., Prior T.	1
A novel alkaline phosphatase isozyme in human adipose tissue	
Okochi T., Seike H., Saeki K. et al.	19
Spectrophotometric assay of xanthine oxidase with 2,2'-azino-di(3-ethylbenzthiazoline-6-sulphonate) (ABTS) as chromogen	
Majkic-Singh N., Bogavac L., Kalimanovska V. et al.	29
Atypical LD isoenzymes with five diffuse bands and evidence for two types of binding	
Chappell D.J., Buttery J.E., Pannall P.R.	37
Quantitation of serum bile acids by isotope dilution with ¹³ C-labelled homologs	
Stellaard F., Paumgartner G.	45
Competitive inhibitor binding assay (CIBA) of captopril and other ACE inhibitors	
Gronhagen-Riska C., Tikkanen L., Fyhrquist F.	53
Purification of porphobilinogen deaminase from human erythrocytes by fast protein liquid chromatography	
De Rooij F.W.M., Hamer C.M., Wilson J.H.P.	61
Development of carboxypeptidase K (creatine kinase conversion factor) in man	
Edwards R.J., Watts D.C.	69
Extended application of RP-HPLC for characterization of human hemoglobin variants	
Baudin V., Wajcman H.	75
Enzyme-linked immunosorbent assay of serum pepsinogen I	
Huang S.C., Miki K., Hirano K. et al.	85
Effect of the partial beta-agonist prenalterol on plasma renin activity in patients with left ventricular failure	
Wathen C.G., Mackay I.G., Glover D.R., Roulston J.E.	97
Determination of neuron specific enolase in amniotic fluid and maternal serum for the prenatal diagnosis of fetal neural tube defects	
Sorensen K., Brodbeck U., Norgaard-Pedersen B.	101
Identification and elimination of an ethylene glycol determination artifact	
Smith N.B.	105

Vol. 162/2, 1987

Biosensor for lactate determination in biological fluids. I. Construction and properties of the biosensor	
Racek J., Musil J.	129
Are 'precipitated LDL' really low density lipoproteins?	
Mainard F., Madec Y.	141
Diagnostic sensitivity of fasting and postprandial serum bile acids determined by different methods	
Mannes G.A., Stellaard F., Paumgartner G.	147
Serum lipids in hyper- and hypothyroidism before and after treatment	
Friis T., Pedersen L.R.	153
The effect of assay conditions on the measurement of anti-elastase function in lung secretions	
Morrison H.M., Kramps J.A., Afford S.C. et al.	165
Development and validation of an improved radioimmunoassay for serotonin in platelet-rich plasma	
Gow I.F., Corrie J.E.T., Williams B.C., Edwards C.R.W.	175
Human inter- α -trypsin inhibitor and immunologically related inhibitors investigated by quantitative immunoelectrophoresis. II. Pathological conditions	
Odum L., Hansen-Nord G., Byrjalsen I.	189
An enzyme-linked immunosorbent assay (ELISA) for plasma progesterone: Immobilised antigen approach	
Elder P.A., Yeo K.H.J., Lewis J.G., Clifford J.K.	199
Interference in the measurement of red cell glycogen by glycogen from white cells	
Collins W., Brown I., Taylor W.H.	207
Human erythrocyte transketolase: No evidence for variants	
Kaufmann A., Uhlhaas S., Friedl W., Propping P.	215
The effect of combined fenofibrate and cholestyramine therapy on low-density lipoprotein kinetics in familial hypercholesterolemia patients	
Malmendier C.L., Delcroix C., Lontie J.-F.	221
Human growth hormone (hGH) in urine and its correlation to serum hGH examined by a highly sensitive sandwich enzyme immunoassay	
Hashida S., Ishikawa E., Kato Y. et al.	229
Inhibition by 5-sulphosalicylic acid of the glutathione reductase recycling assay for glutathione analysis	
Allen K.G.D., Arthur J.R.	237

Vol. 162/3, 1987

An evaluation of the low-pH enzymatic assay of urinary D-glucaric acid, and its use as a marker of enzyme induction in exocrine pancreatic disease	
Sandle L.N., Braganza J.M.	245
On the homogeneity of stools with respect to bile acid composition and normal day-to-day variations: A detailed qualitative and quantitative study using capillary column gas chromatography-mass spectrometry	
Setchell K.D.R., Ives J.A., Cashmore G.C., Lawson A.M.	257
A method for measuring testosterone-estradiol-binding globulin - new analysis of experimental data and comparison with two commercially available kits	
Lecureuil N., Trochet J., Crouzat-Reynes G. et al.	277
The fat meal test of malabsorption: Some further experience	
Steen G., Holthuis N.	291
Deficient cholesterol side chain oxidation in patients without peroxisomes (Zellweger syndrome): Evidence for the involvement of peroxisomes in bile acid synthesis in man	
Wanders R.J.A., Schutgens R.B.H., Heymans H.S.A.	295
Development of a method to monitor low molecular mass hydrocarbons in exhaled breath of man: Preliminary evaluation of its interest for detecting a lipoperoxidation process in vivo	
Hotz P., Hoet P., Lauwerys R., Buchet J.-P.	303

Lectin-affinity chromatography of serum gamma-glutamyltransferase in liver disease Delanghe J.R., De Buyzere M.L., De Scheerder I.K. et al.	311
Human erythrocyte γ -glutamyltransferase in liver diseases Daniel D.S., Ramakrishna B.S., Balasubramanian K.A.	319
Uromuroid (Tamm-Horsfall glycoprotein) forms different polymeric arrangements on a filter surface under different physicochemical conditions Wiggins R.C.	329
Radioimmunoassay of glandular kallikrein in human plasma after partial purification by immunoaffinity column Nishimura K., Inoue Y., Kokubu T.	341

Vol. 163/1, 1987

Coeliac disease: A critical review of aetiology and pathogenesis Davidson A.G.F., Bridges M.A.	1
Enzyme-linked immunosorbent assay (ELISA) for IgG antibodies to thyroglobulin Shimojo N., Kohno Y., Tarutani O. et al.	41
Urinary protein/creatinine ratio as an indicator of allograft function following live related donor renal transplantation Krishna K.S., Pandey A.P., Kirubakaran M.G., Kanagasabapathy A.S.	51
Enzyme linked immunosorbent assay for the measurement of nonenzymatically glucosylated proteins in serum and in tissues Go R., Quinn T.J., Gonen B.	63
Total and free disopyramide by fluorescence polarization immunoassay and relationship between free fraction and alpha-1 acid glycoprotein Chen B.H., Taylor E.H., Pappas A.A.	75
Determination of trimethoprim in pediatric samples by high-performance liquid chromatography Hoppu K., Arjomaa P.	81
Binding to cellulose of IgG from patients with interstitial lung diseases Dewair M., Baur X.	87
Diagnostic significance of urinary enzymes: Development of a high performance liquid chromatographic method for the measurement of urinary lysozyme Severini G., Aliberti L.M.	97
Enzyme immunoassay of total insulin in human serum containing anti-insulin antibodies Kohno T., Ishikawa E., Sugiyama S. et al.	105
Apolipoprotein and lipoprotein cholesterol concentrations in trained and sedentary males Rifai N., King M.E., De Meersman R. et al.	113

Vol. 163/2, 1987

The relative extent of glycation of haemoglobin and albumin Olufemi S., Talwar D., Robb D.A.	125
The effect of 2,4-dihydropyridine-N-oxide, a new orally active iron chelator, on iron excretion in mice Kontoghiorghes G.J.	137
Recognition of cervical neoplasia by the estimation of a free-radical reaction product (octadeca-9,11-dienoic acid) in biopsy material Griffin J.F.A., Wickens D.G., Tay S.K. et al.	143
Recognition of cervical neoplasia by the estimation of a free-radical reaction product (octadeca-9,11-dienoic acid) in exfoliated cells Tay S.K., Singer A., Griffin J.F.A. et al.	149
Evaluation of gravimetric assays of the H ₂ O concentration in human serum and urine De Jong Th. G.M., Huizenga J.R., Gips C.H.	153

The increase in zincaemia provoked by starvation does not influence the intestinal absorption of zinc	
Couturier E., Praet J.P., Vis H.L., Devroede P.	165
Effect of temperature and sample preparation on performance of ion-moderated partition chromatography of organic acids in biological fluids	
Rumsby G., Belloque J., Ersser R.S., Seakins J.W.T.	171
Microheterogeneity forms of α_1 -acid glycoprotein as indicators of rheumatoid arthritis activity	
Mackiewicz A., Pawlowaki T., Mackiewicz-Pawlowska A. et al.	185
Serum pepsinogens I and II and stomach cancer	
Stemmermann G.N., Samloff I.M., Nomura A.M.Y., Heilbrun L.K.	191
Preparative fractionation of amyloid proteins on a microgram scale by high-performance liquid chromatography and polyacrylamide gel electrophoresis	
Kaplan B., Pras M.	199
A simple immunotechnique for the determination of serum concentration of apolipoprotein E	
Rifai N., Silverman L.M.	207
Stability of enzymes of lysosomal origin in human cerebrospinal fluid	
Goi G., Fabi A., Lombardo A. et al.	215
m-Aminophenylboronate affinity ligands distinguish between nonenzymically glycosylated proteins and glycoproteins	
Gould B.J., Hall P.M.	225
Electrophoretic differentiation of acquired angioedema from hereditary angioedema	
Papadopoulos N.M., Frank M.M.	231

Vol. 163/3, 1987

ELISA for determination of albumin in the nanogram range: Assay in cerebrospinal fluid and comparison with radial immunodiffusion	
Schwerer B., Bach M., Bernheimer H.	237
Quantification of human apolipoprotein E in plasma and lipoprotein subfractions by a non-competitive enzyme immunoassay	
Koffigan M., Kora I., Clavey V. et al.	245
A non-competitive enzyme immunoassay of human α -lactalbumin in biological fluids and tissue extracts	
Simickova M., Lang B.A., Semotan K., Kocent A.	257
Phosphatidylethanolamine methyltransferase and cAMP, cGMP phosphodiesterases in lymphocytes and monocytes in sarcoidosis	
Pacheco Y., Dubois M., Prigent A.F. et al.	267
Partial characterization of human ileal alkaline phosphatase: Differences between human ileal and duodenal enzymes	
Miura M., Matsuzaki H., Sakagishi Y., Komoda T.	279
Evidence for endotoxin binding capacity of human Gc-globulin and transferrin	
Berger D., Beger H.G.	289
Human hypoxanthine-guanine phosphoribosyltransferase. Development of a spectrophotometric assay and its use in detection and characterization of mutant forms	
Keough D.T., McConachie L.A., Gordon R.B. et al.	301
Direct assay for testosterone in saliva: Relationship with a direct serum free testosterone assay	
Johnson S.G., Joplin G.F., Burrin J.M.	309
Protein transfer at the blood cerebrospinal fluid barrier and the quantitation of the humoral immune response within the central nervous system	
Reiber H., Felgenhauer K.	319
Measurement by bioluminescence technique of erythrocyte membrane Na^+, K^+ -ATPase activity in hypertensive patients	
Lechi C., Corradini P., Polignano R. et al.	329
Detection of desialylated forms of human chorionic gonadotropin	
Imamura S., Armstrong E.G., Birken S. et al.	339

Vol. 164/1, 1987

Evaluation of iron-deficiency anemia by an iron supplementation trial in children living at a 2800-m altitude Estrella R., Hercberg S., Maggy G. et al.	1
Urinary thiosulphate as an indicator of exposure to hydrogen sulphide vapour Kangas J., Savolainen H.	7
A circulating complex between ASAT and IgG in serum in an apparently healthy woman Fex G., Berntorp K.	11
24-h Catecholamine excretion: Relationships with age and weight Jenner D.A., Harrison G.A., Prior I.A.M. et al.	17
Immunological comparison of biotinidase in serum from normal and biotinidase-deficient individuals Wolf B., Miller J.B., Hymes J. et al.	27
Systemic membrane defect and the inhibition of lymphocyte capping in Duchenne Muscular Dystrophy Goldsmith B.M., Gruemer H.-D.	33
Determination of creatine kinase isozymes in sera and tissues of patients with various lung carcinomas Usui A., Fujita K., Imaizumi M. et al.	47
Human pheochromocytoma phenol sulfotransferase: Biochemical properties and activities of thermolabile and thermostable forms Sinaheimer E.G., Anderson R.J.	55
Enzyme-linked immunosorbent assay (ELISA) for steroid hormones with polyclonal and monoclonal antibodies: An assay for urinary aldosterone Hanquez C., Urios P., Desfossez B. et al.	71
Physical state abnormality of erythrocyte membrane in myotonic dystrophy: A spin label study Thomas M.A., Lamas E., Daveloose D. et al.	83
Free and Apo B-associated Lpa-specific protein in human serum Gries A., Nimpf J., Nimpf M. et al.	93
Effects of exogenous β -galactosidase on cultured fibroblasts with β -galactosidase deficiency Nakao Y., Kozutsumi Y., Fukui S. et al.	101

Vol. 164/2, 1987

Liver-specific F antigen in the serum of patients with liver diseases and the detection of early stage hepatocellular carcinoma Mori Y., Iwama S., Mori T. et al.	127
Relationships between the lipid content and the rheological properties of airway secretions in cystic fibrosis Galabert C., Jacquot J., Zahm J.M., Puchelle E.	139
Relationship between albumin and fructosamine concentration in diabetic and non-diabetic sera Johnson R.N., Metcalf P.A., Baker J.R.	151
Plasma amino-acids analysis: Effects of delayed samples preparation and of storage Schaefer A., Piquard F., Haberey P.	163
A simple chromogenic assay for arylsulfatase A Lee-Vaupel M., Conzelmann E.	171
Determination of erythrocyte glucose 1,6-bisphosphate — A comparison of two methods using a centrifugal analyzer Thorburn D.R., Kuchel P.W.	181
Improved between-laboratory agreement for specific protein assays in serum following introduction of a common reference preparation (SPS-01) demonstrated in an external quality assessment scheme Chambers R.E., Bullock D.G., Whicher J.T.	189

Clinical application of enzyme immunoassay in the analysis of citrullinemia Imamura Y., Kobayashi K., Yamashita T. et al.	201
An alternative way of presenting quantitative amino acid results Briddon A., Oberholzer V.G.	209
Removal of ascorbate from urine prior to assaying with a commercial oxalate kit Kasidas G.P., Rose G.A.	215
Serum angiotensin converting enzyme activity in ex-smokers Ninomiya Y., Kioi S., Arakawa M.	223
An evaluation of four methods for the detection of heterozygous cystinuria Giugliani R., Ferrari I., Greene L.J.	227
A comparison of the utility of Dowex resin and polybenzimidazole Aurorez resin in the determination of urinary porphobilinogen concentrations Galbraith R.A., Sassa S., Kappas A.	235

Vol. 164/3, 1987

Lipid peroxidation and malondialdehyde in the synovial fluid and plasma of patients with rheumatoid arthritis Wade C.R., Jackson P.G., Highton J., Van Rij A.M.	245
The molecular abnormality of albumin Niigata: 269 Asp → Gly Sugita O., Endo N., Yamada T. et al.	251
Identification of glutarylcarbitine in glutaric aciduria type I by carboxylic acid analyzer with an ODS reverse-phase column Kidouchi K., Sugiyama N., Morishita H. et al.	261
Modified phosphoribosylpyrophosphate (PRPP) radioenzymatic assay: Increased sensitivity, technical simplification and new applications Ghitis J., Waxman S.	267
5'-Nucleotide phosphodiesterase isoenzymes in human serum: Quantitative measurement and some biochemical properties Luthje J., Ogilvie A.	275
Rapid suppression of plasma alkaline phosphatase activity after renal transplantation in patients with osteodystrophy Cundy T., Kanis J.A.	285
Purification of placental protein PP12 from human amniotic fluid and its comparison with PP12 from placenta by immunological, physicochemical and somatomedin-binding properties Koistinen R., Huhtala M.-L., Stenman U.-H., Seppala M.	293
A rapid and precise assay for calmodulin and cyclic AMP phosphodiesterase on a centrifugal analyser Walker S.W., Milner S.W., Howie A.F.	305
Effects of purified fish oil on platelet lipids and function in diabetic women Tilvis R.S., Rasi V., Viinikka L. et al.	315
Reduced plasma and white blood cell selenium levels in haemodialysis patients Foote J.W., Hinks L.J., Lloyd B.	323
The effect of fatty acid chain length on the rate of arylester hydrolysis by various albumins Wolfbeis O.S., Gurakar A.	329
Isoenzymes of urinary N-acetyl-β-D-glucosaminidase (NAG) in patients with renal transplants Yuen C.-T., Corbett C.R.R., Kind P.R.N. et al.	339

Vol. 165/1, 1987

Animal models of inherited hematologic disease Kaneko J.J.	1
Apoenzyme content of serum aminotransferases in patients with a renal allograft treated with	

cyclosporine A and azathioprine	
Hafkenscheid J.C.M., Rosier J.G.M.C., Van Dijk C.M.C.E.	21
Fatty acid composition of phospholipids in bile in man: promoting effect of deoxycholate on arachidonate	
Van Berge Henegouwen G.P., Van der Werf S.D.J., Ruben A.T.	27
A new simple screening method for the diagnosis of medium chain acyl-CoA dehydrogenase deficiency	
Bhuiyan A.K.M.J., Watmough N.J., Turnbull D.M. et al.	39
The direct barbituric acid assay for nicotine metabolites in urine: A simple colorimetric test for the routine assessment of smoking status and cigarette smoke intake	
Barlow R.D., Stone R.B., Wald N.J., Puhakainen E.V.J.	45
Correlation between the cystic fibrosis protein detected by isoelectric focusing and a protein of 12 kDa detected by SDS-polyacrylamide gel electrophoresis	
Barthe C., Figarella C., Galabert C., Guy-Crotte O.	53
Thermal desorption — gas chromatography of plasma isovaleraldehyde in hepatic encephalopathy in man	
Al Mardini H., O'Brien C.J., Bartlett K. et al.	61
Conditions for a reliable application of the lysoplate method in the determination of lysozyme	
Gupta D.K., Von Figura K., Hasilik A.	73
Value of glial fibrillary acidic protein determination in amniotic fluid for prenatal diagnosis of neural tube defects	
Van Regemorter N., Gheuens J., Noppe M. et al.	83
Comparison of fat agglutination slide test and latex test for C-reactive protein	
Hulman G., Fuller M.	89
Approved IFCC methods. Reference method (1986) for pH measurement in blood	
Maas A.H.J., Weisberg H.F., Burnett R.W. et al.	97
Approved recommendation (1986) on the theory of reference values. Part 1. The concept of reference values	
Solberg H.E.	111

Vol. 165/2-3, 1987

Thyroid function in leukaemia patients after allogeneic bone marrow transplantation	
Frolich M., De Planque M.M., Goslings B.M., Meinders A.E.	127
Interference of blood clotting factors in the determination of HDL ₂ and HDL ₃ -cholesterol by the dual precipitation method	
Demacker P.N.M., Hak-Lemmers H.L.M., Van Heijst P.J.	133
Human serum sialo transferrins in diseases	
Van Eijk H.G., Van Noort W.L., De Jong G., Koster J.F.	141
Determination of polyamines and metabolites in cerebrospinal fluid by isotope dilution mass fragmentography, and a clinical application	
Van Den Berg G.A., Schaaf J.M., Nagel G.T. et al.	147
Does the degree of polymerization of haptoglobin influence the results of the radial immunodiffusion assay?	
Van Rijn H.J.M., Wisman J.P.W., Gmelig Meyling F.	155
Catalytic and immunologic activities of placental-like alkaline phosphatase in clinical studies. The value of PLAP in follow-up of ovarian cancer	
Haije W.G., Van Driel J., Van Der Burg M.E.L.	165
Therapeutic drug monitoring of amitriptyline: impact of age, smoking and contraceptives on drug and metabolite levels in bulimic women	
Edelbroek P.M., Zitman F.G., Knoppert-Van Der Klein E.A.M., et al.	177
Assay of oestrogen and progestin receptors in human meningioma cytosols using immunological methods	
Blankenstein M.A., Van Der Meulen-Dijk C., Thijssen J.H.H.	189

Urinary excretion of 2-methyl-2,3-butanediol and 2,3-pentanediol in patients with disorders of propionate and methylmalonate metabolism	
Duran M., Ketting D., Bruinvis L. et al.	197
Pregnancy and guinea-pig isoferritins — isolation and characterization of both isoferritins	
Bierings M.B., Heeren J.W.A., Van Noort W.L. et al.	205
Urinary polyamines and their metabolites during new combination chemotherapeutic treatments of high grade non-Hodgkin lymphoma	
Muskiet F.A.J., Van Imhoff G.W., Van Den Berg G.A. et al.	213
Laser diffraction ellipsometry of erythrocytes under controlled shear stress using a rotational viscosimeter	
Hardeman M.R., Goedhart P., Breederveld D.	227
Influence of gastric emptying on the PABA test	
Hoek F.J., Sanders G.T.B., Tytgat G.N.J.	235
Reinvestigations into the formation and assay of C3bBbP complexes	
Asghar S.S., Barendsen C.Y., Van Der Helm H.J.	243
Glucagon and insulin response to meals in non-obese and obese Dutch women	
Hill P., Garbaczewski L., Koppeschaar H. et al.	253
Analytical and clinical evaluation of commercial protein C assays	
Sturk A., Morrien-Salomons W.M., Huisman M.V. et al.	263
Differential diagnosis of genetic disease by DNA restriction fragment length polymorphisms	
Bolhuis P.A., Defesche J.C., Van Der Helm H.J.	271
ELISA of ceruloplasmin and alpha-2-macroglobulin in paired bronchoalveolar lavage fluid and serum samples	
Out T.A., Jansen H.M., Van Steenwijk R.P. et al.	277
Immunoglobulin D in cerebrospinal fluid	
Out T.A., Hiasche E.A., Van Walbeek H.K., Van Der Helm H.J.	289
Quantification of cholesterol nucleation promoting activity in human gallbladder bile	
Drapers J.A.G., Groen A.K., Stout J.P.J. et al.	295
Prenatal diagnosis of Zellweger syndrome by measurement of very long chain fatty acid (C26:0) β -oxidation in cultured chorionic villous fibroblasts: implications for early diagnosis of other peroxisomal disorders	
Wanders R.J.A., Van Wijland M.J.A., Van Roermund C.W.T. et al.	303
Alanine glyoxylate aminotransferase and the urinary excretion of oxalate and glycolate in hyperoxaluria type I and the Zellweger syndrome	
Wanders R.J.A., Van Roermund C.W.T., Westra R. et al.	311
X-linked adrenoleukodystrophy: defective peroxisomal oxidation of very long chain fatty acids but not of very long chain fatty acyl-CoA esters	
Wanders R.J.A., Van Roermund C.W.T., Van Wijland M.J.A. et al.	321
A sensitive radioimmunoassay of atrial natriuretic peptide in human plasma, using a tracer with an immobilized glycouril agent	
Rosmalen F.M.A., Tan A.C.I.T.L., Tan H.S., Benraad T.J.	331
A symmetric dialysis method for the determination of free testosterone in human plasma	
Swinkels L.M.J.W., Ross H.A., Benraad T.J.	341
Determination of urinary cystine for monitoring anticystinuric therapy	
Backer E.T., Van Den Ende A., Geus W.P.	351
Effect of ritodrine on thyroid hormone concentrations	
Menheere P.P.C.A., Essed G.G.M., Sels J.P.J.E.	359
β -Aminoisobutyric acid as a marker of thymine catabolism in malignancy	
Van Gennip A.H., Van Bree-Blom E.J., Abeling N.G.G.M. et al.	365
The effect of oral contraceptives on plasma-free and salivary cortisol and cortisone	
Meulenbergh P.M.M., Ross H.A., Swinkels L.M.J.W., Benraad T.J.	379

Vol. 166/1, 1987

Automated enzymatic standardized lipid analyses for plasma and lipoprotein fractions	
McNamara J.R., Schaefer E.J.	1

Application of transformation algorithm and nonparametric calculation in determining the reference intervals of some urine constituents and characteristics Soliman S.A., Abdel-Hay M.H., Tayeb O.S., Sulaiman M.I.	9
Immunochemical determination of acetylcholinesterase in amniotic fluid — An evaluation of eleven monoclonal antibodies Rasmussen A.G., Sorensen K., Selmer J. et al.	17
Agarose gel isoelectrofocusing of UDP-galactose pyrophosphorylase and galactose-1-phosphate uridylyltransferase. Developmental aspect of UDP-galactose pyrophosphorylase Shin Y.S., Niedermeier H.-P., Endres W. et al.	27
Atrial natriuretic peptide in human plasma — Comparison of radioreceptor versus radioimmunoassay Sagnella G.A., Buckley M.G., Markandu N.D., MacGregor G.A.	37
Age-related accumulation of phytanic acid in plasma from patients with the cerebro-hepato-renal (Zellweger) syndrome Wanders R.J.A., Smit W., Heymans H.S.A. et al.	45
Serum human pancreatic elastase I levels in pancreatic disease Buamah P.K., Cornell C., Cassells-Smith A.J., Skillen A.W.	57
Detection of photobilirubin in urine of jaundiced infants supporting the diagnosis of 'Bronze Baby Syndrome' Meisel P., Jahrig D., Meisel M.	61
A simple latex agglutination test for urinary albumin screening Paoli C., Bardelli F., Tarli P. et al.	67
Isoelectric focusing of the cytosolic human creatine kinase isoenzymes Nealon D.A.	73
Serum aromatic L-amino acid decarboxylase activity in patients with acute hepatitis Nishigaki I., Nagatsu T.	79
Turbidimetric measurement of microalbuminuria Harmoinen A., Vuorinen P., Jokela H.	85
Determination of aromatic and neutral aminoacids by HPLC in blood specimens collected on filter paper Schrynmackers-Pitance P., Schoos-Barbette S.	91

Vol. 166/2-3, 1987

A comparison of dopamine and homovanillic acid excretion, as prognostic indicators in malignant pheochromocytoma Tippett P.A., West R.S., McEwan A.J. et al.	123
Comparison of N-acetyl- β -glucosaminidase and albuminuria with clinical finding of microangiopathy in type I diabetes mellitus Skrha J., Perusicova J., Stolba P. et al.	135
Evidence for an increased glycation of IgG in diabetic patients Danze P.-M., Tarjoman A., Rousseaux J. et al.	143
Tubular maximum for calcium reabsorption: Lack of diagnostic usefulness in primary hyperparathyroidism and familial hypocalciuric hypercalcaemia Kent G.N., Bhagat C.I., Garcia-Webb P., Gutteridge D.H.	155
A simple enzymatic fluorimetric method for the determination of branched-chain L-amino acids in microlitre volumes of plasma Gleeson M., Maughan R.J.	163
Blood and liver selenium concentrations in patients with liver diseases Valimaki M., Alfthan G., Pikkarainen J. et al.	171
Studies of the sialylation and microheterogeneity of human serum α_1 -acid glycoprotein in health and disease Moule S.K., Peak M., Thompson S., Turner G.A.	177
Comparison of the precision of seven analytical methods for the H ₂ O concentration in human serum and urine De Jong Th. G.M., Huizenga J.R., Wolthers B.G. et al.	187

Separation of plasma carotenoids and quantitation of β -carotene using HPLC	
Gatautis V.J., Pearson K.H.	195
Purification and characterization of glutathione S-transferases in human kidney	
Tateoka N., Tsuchida S., Soma Y., Sato K.	207
A fluorometric micromethod for estimation of carnosinase in dried blood samples	
Orfanos A.P., Guthrie R., Jinks D.C.	219
Estimation and characterization of glycosylated carbonic anhydrase I in erythrocytes from patients with diabetes mellitus	
Kondo T., Murakami K., Ohtsuka Y. et al.	227
Sandwich enzyme immunoassay for human interleukin- 1β (hIL- 1β) in urine	
Tanaka K., Ishikawa E., Ohmoto Y., Hirai Y.	237
Detection of microalbuminuria in diabetic patients using a simple latex agglutination test	
Spooner R.J., Weir R.J., Frier B.M.	247
Peroxisomal very long-chain fatty acid β -oxidation in human skin fibroblasts: Activity in Zellweger syndrome and other peroxisomal disorders	
Wanders R.J.A., Van Roermund C.W.T., Van Wijland M.J.A. et al.	255
Specific assays for human alkaline phosphatase isozymes	
Hirano K., Matsumoto H., Tanaka T. et al.	265
Improved method for determination of high density lipoprotein cholesterol using a sensitive reagent and a centrifugal analyzer	
Moahides J.S.	275
Diode-array spectrophotometry for simultaneous measurement of hemoglobin pigments	
Fogh-Andersen N., Siggaard-Andersen O., Lundsgaard F.C., Wimberley P.D.	283
Spectrophotometric determination of hemoglobin pigments in neonatal blood	
Fogh-Andersen N., Siggaard-Andersen O., Lundsgaard F.C., Wimberley P.D.	291
Estimation of whole body bone resorption rate: A comparison of urinary total hydroxyproline excretion with two radioisotopic tracer methods in osteoporosis	
Deacon A.C., Hulme P., Hesp R. et al.	297
Immunoglobulin light-chain immunoblots or urine proteins from patients with tubular and Bence-Jones proteinuria	
Norden A.G.W., Fulcher L.M., Flynn F.V.	307
Hemoglobin A ₁ and serum fructosamine levels in hyperthyroidism	
Ford H.C., Lim W.C., Crooke M.J.	317
Detection of wheat gliadin contamination of gluten-free foods by a monoclonal antibody dot immunobinding assay	
Freedman A.R., Galfre G., Gal E. et al.	323
Estimation of low levels of phosphatidylglycerol in aspirates of newborns using the AmnioStat FLM agglutination test	
Egberts J., Contant J., Wijnands J.B.G.	329

Vol. 167/1, 1987

Rheumatoid factors	
Dorner R.W., Alexander Jr. R.L., Moore T.L.	1
Direct analysis of tetrahydrobiopterin in cerebrospinal fluid by high-performance liquid chromatography with redox electrochemistry: Prevention of autoxidation during storage and analysis	
Howells D.W., Hyland K.	23
The crypt-villus distribution of the 10000 M(r) intestinal calcium binding protein in human jejunum	
Staun M.	31
Lack of relationship between beta-hexosaminidase activity and retinopathy in insulin dependent diabetics	
Agardh C.-D., Agardh E., Hultberg B.	37
Immunochemical characterization and quantification of the sex steroid-binding protein (SBP) in human amniotic fluid	
Auzelle M.-P., Tardivel-Lacombe J., Domingo M., Egloff M., Degrelle H.	43

Cholinesterase activities in resected bowel specimens from children with Hirschsprung's disease Causse E., Vaysse P., Fabre J., Valdiguie P., Thouvenot J.P.	51
Relationship between effects of added albumin, initial free thyroxine value and endogenous serum-binding protein concentrations on Amerlex® free thyroxine estimations Midgley J.E.M., Winton M.R.J., Wilkins T.A.	67
Guanidino compounds in serum and cerebrospinal fluid of non-dialyzed patients with renal insufficiency De Deyn P.P., Marescau B., Cuykens J.J., Van Gorp L., Lowenthal A., De Potter W.P.	81
Underestimation of serum albumin by the bromocresol purple method and a major endogenous ligand in uremia Mabuchi H., Nakahashi H.	89
Determination of total cholesterol in serum by flow injection analysis with immobilized enzymes Fernandez-Romero J.M., Luque De Castro M.D., Valcarcel M.	97
A rapid method for determining catalase in human blood Lamoureux G., Bourdeau S., Dubois G., Charbonneau R., Gagnon M., Grad B.R.	105

Vol. 167/2, 1987

Faecal lysozyme: Determination, reference intervals and some data in gastro-intestinal disease Costongs G.M.P.J., Hemrika M.H., Engels L.G.J.B. et al.	125
CSF concentration and CSF/blood ratio of fuel related components in children after prolonged fasting Lamers K.J.B., Doesburg W.H., Gabreels F.J.M. et al.	135
Isoenzyme specificities of immunoglobulins isolated from lactate dehydrogenase-immunoglobulin complexes Kanemitsu F.	147
Development and application of an enzyme-linked immunosorbent assay for acid stable trypsin inhibitor (ASTI) in human plasma Yoshida E., Sumi H., Maruyama M. et al.	155
Quantitative detection of a variant prealbumin associated with type I familial amyloidotic polyneuropathy (Japanese type) by high performance liquid chromatography Ikegawa S., Tanase S., Araki S., Morino Y.	165
Evaluation of the average of patients: Application to endocrine assays Douville P., Cembrowski G.S., Strauss J.F.	173
Intracellular magnesium of mononuclear cells from venous blood of clinically healthy subjects Reinhart R.A., Marx Jr. J.J., Haas R.G., Desbiens N.A.	187
HDL cholesterol and the acute phase reaction following myocardial infarction and acute pancreatitis Fahie-Wilson M., Mills R., Wilson K.	197
4-Nitrophenylalkaline phosphatase inhibition in muscular dystrophy Goldenberg A.L., Sanchez J.J., Garcia A.M. et al.	211
A simple, reproducible and cheap batch method for the analysis of serum glycoproteins using a Sepharose-coupled lectins and silver staining Thompson S., Latham J.A.E., Turner G.A.	217
An improved spot-test for the detection of paraquat and diquat in biological samples Woollen B.H., Mahler J.D.	225
Analysis of 25-hydroxyvitamin D in plasma using solid phase extraction Shephard G.S., Carlini S.M., Hanekom C., Labadarios D.	231

Vol. 167/3, 1987

Serum aldolase isoenzymes in benign and malignant liver disease Royds J.A., Kennedy H.J., Little P.V., Taylor C.B., Triger D.R.	237
--	-----

Effects of severe burns on glycan microheterogeneity of four acute phase proteins Mallet B., Franc J.L., Miquel M., Arnaud C.	247
Inorganic phosphate transport across the red blood cell membrane: The effect of exposure to hyperoxia Webster N.R., Toothill C.	259
Clinical usefulness of cimetidine for the treatment of acute intermittent porphyria — a preliminary report Horie Y., Udagawa M., Hirayama C.	267
Vitamin D-binding protein synthesized by a human hepatocellular carcinoma cell line Egawa T., Yamamoto H., Kishimoto S.	273
Glycylproline dipeptidyl aminopeptidase and γ -glutamyl transpeptidase in human hepatic cancer and embryonal tissues Kojima J., Ueno Y., Kasugai H., Okuda S., Akedo H.	285
A large molecular form of glucagon-like peptide-1 (GLP-1) immunoreactivity is co-released with glucagon from pancreas by arginine in normal subjects Hirota M., Shimizu I., Ohboshi C., Nishino T., Shima K.	293
An increased apo A-IV serum concentration of patients with chronic renal failure on hemodialysis Seishima M., Muto Y.	303
A simple differential immunoprecipitation assay of urinary acid and neutral α -glucosidases for glycogenosis II Tsuiji A., Yang R.-C., Omura K., Imabayashi T., Suzuki Y.	313
Hydrolysis of phosphatidate by human placental alkaline phosphatase Sumikawa K., Saeki K., Okochi T., Adachi K., Nishimura H.	321
Macromolecules inhibit calcium oxalate crystal growth and aggregation in whole human urine Edyvane K.A., Hibberd C.M., Harnett R.M., Marshall V.R., Ryall R.L.	329
Competitive enzyme immunoassay for human serum apolipoproteins C1, C11 and C111 Fu M., Liu B., Liu W., Fang D.	339
Structurally abnormal transthyretin causing familial amyloidotic polyneuropathy in Sweden Nakazato M., Steen L., Holmgren G., Kurihara T., Matsukura S., Kangawa K., Matsuo H.	341

Vol. 168/1, 1987

Detergent induced lysis of erythrocytes in kwashiorkor Rao A., Onuora C.U., Cherian A.	1
Abnormal degradation of erythrocyte membrane proteins in hereditary spherocytosis Gaczynska M., Bartosz G., Judkiewicz L., Rosin J.	7
Immunoradiometric assay of serum thyroid stimulating hormone (TSH) as a first-line test in a mixed hospital population Ronnov-Jessen D., Skov L., Faber J.	13
Changes in magnesium, zinc and calcium in men and women after an acute myocardial infarction Speich M., Gelot S., Robinet N., Arnaud P., Nicolas G.	19
Black-white differences in serum Lp(a) lipoprotein levels Parra H.-J., Luyeye I., Bouramoue C., Demarquilly C., Fruchart J.-C.	27
Interference of an α_2 component in immunological determinations of pregnancy-specific β_1 glycoprotein in serum Pala A., Di Ruzza A., Curtilli L., Strom R., Moro M., Reverberi L., Carenza L.	33
Evaluation of RU-27967 as a ligand to determine the progesterone receptor Abendstein B., Marth C., Daxenbichler G.	47
Differential reactivity of α -fetoprotein with lectins and its usefulness in the diagnosis of various liver diseases Buamah P.K., Cornell C., Cassells-Smith A.J., Skillen A.W.	69
Increased serum lactate dehydrogenase isoenzyme 1 and macro creatine kinase type 2 in a patient with lung cancer Lentjes E.G.W.M., Backer E.T.	75

Enzyme-linked immunosorbent assay (ELISA) for human epidermal growth factor (hEGF)	
Abe Y., Sagawa T., Sakai K., Kimura S.	87
A highly sensitive enzyme immunoassay of anti-insulin antibodies in human serum	
Kohno T., Ishikawa E., Sugiyama S., Kamamo M., Kuzuya H., Imura H.	97
Effects of surfactants on human serum biotinidase	
Hayakawa K., Oizumi J.	109

Vol. 168/2, 1987

Hemoglobin J-Baltimore (β 16(A13)Gly - Asp): Interference with the assay of HbA(1c)	
Vandenesch F., Baklouti F., Francina A., Vianey-Liaud C., Bertrand A., Le Devehat C., Delaunay J.	121
Detection of glucose-6-phosphate dehydrogenase deficiency in erythrocytes: A spectrophotometric assay and a fluorescent spot test compared with a cytochemical method	
Wolf B.H.M., Weening R.S., Schutgens R.B.H., Van Noorden C.J.F., Vogels I.M.C., Nagelkerke N.J.D.	129
Measurement of fetal surfactant production by fluorescence polarization of amniotic fluid in complicated pregnancies	
Masturzo P., Salmona M., Rossi G., Luerti M.	137
Stable isotope dilution analysis of pipecolic acid in cerebrospinal fluid, plasma, urine and amniotic fluid using electron capture negative ion mass fragmentography	
Kok R.M., Kaster L., De Jong A.P.J.M., Poll-The B., Saudubray J.-M., Jakobs C.	143
The molecular nature of cholecystokinin in human plasma	
Cantor P., Rehfeld J.F.	153
Direct determination of lead in blood by differential pulse stripping voltammetry	
Nedeljkovic M., Stojadinovic L., Matovic V., Koricanac Z.	169
Histamine uptake by human platelets	
Gill D.S., Barradas M.A., Mikhailidis D.P., Dandona P.	177
Studies with an automated kinetic assay for plasma angiotensin-converting enzyme activity and its potentiation by zinc ion	
Roulston J.E., Allan D.	187
Purification and characterization of a pancreas cancer-related antigen	
Furukawa J., Shimano T., Inaji H., Saito M., Matsuura N., Monden T., Mori T.	219
Determination of acid phosphatase in biological fluids using a new substrate, 2,6-dichloro-4-nitrophenyl phosphate	
Teshima S., Hayashi Y., Ando M.	231
Development of an automated kinetic Jaffe method designed to minimise bilirubin interference in plasma creatinine assays	
Knapp M.L., Mayne P.D.	239
Fructosamine measurements in serum and plasma	
MacDonald D., Pang C.-P., Cockram C.S., Swaminathan R.	247

Vol. 168/3, 1987

Stability of radioiodinated ferritin	
Covell A.M., Cook J.D.	261
A second albumin variant in an Irish population	
Leahy D.T., McLaughlin H.	273
Plasma fibronectin in normolipidaemic and hyperlipidaemic uraemic patients treated with haemodialysis	
De Giorgio L.A., Maffucci G., Toscano G., Gironi A., Vignoli S., Seghieri G.	281
Diagnostic significance of different urinary enzymes in patients suffering from chronic renal disease	
Jung K., Schulze B.-D., Sydow K.	287

Microheterogeneity of serum alkaline phosphatase isoenzymes as revealed by isoelectric focusing Rosendahl K., Waldenlind L., Onica D.	297
Determination of erythrocyte protoporphyrin by magnetic circular dichroism Ivanetich K.M., Jeans D.R.	307
Laboratory and clinical features of self-poisoning with salbutamol and terbutaline Jarvie D.R., Thompson A.M., Dyson E.H.	313
Determination of selenium in human spermatozoa and prostasomes using base digestion and electrothermal atomic absorption spectrophotometry Suistomaa U., Saaranen M., Vanha-Perttula T.	323
Assay of sodium-lithium countertransport rate by direct measurement of erythrocyte lithium concentrations Siebers R.W.L., Maling T.J.B.	329
A modified statistical approach for the detection of outlying values in external quality control: Comparison with other techniques Thienpont L.M.R., Steyaert H.L.C., De Leenheer A.P.	337

Vol. 169/1, 1987

Biochemical tests in the diagnosis, classification, and management of patients with malignant lymphoma and leukemia Goldberg D.M., Brown D.	1
New conjugated urinary metabolites in intermediate type maple syrup urine disease Hagenfeldt L., Naglo A.S.	77
Plasma glutathione S-transferase measurements in patients with alcoholic cirrhosis Beckett G.J., Hayes P.C., Hussey A.J., Bouchier I.A.D., Hayes J.D.	85
Lactate dehydrogenase, isoenzyme patterns and cation levels in human breast gross cyst fluid Biagioni S., Mannello F., Stella F., Battistelli S., Marcheggiani F., Cerroni L., Artico M., Terzano C., Troccoli R.	91
Improved method for the quantification of aminoglutethimide in urine samples Van Tellingen O., Bonfrer J.M.G., Nooyen W.J., Bruning P.F.	99
Detection of 3 β -hydroxysteroid dehydrogenase deficiency in a newborn by means of urinary steroid analysis Wolthers B.G., De Vries I.J., Volmer M., Nagel G.T.	109
Removal of interference by antihypertensive drugs in the spectrophotometric assay of metanephrines Crawford G.A., Gallery E.D.M., Gyory A.Z.	117
A simple diagnostic method of adrenoleukodystrophy: Total fatty acid analysis of erythrocyte membranes Antoku Y., Sakai T., Tsukamoto K., Imanishi K., Ohtsuka Y., Iwashita H., Goto I.	121
Urinary bile alcohols in liver dysfunction Hiraoka T., Kihira K., Kohda T., Kosaka D., Kajiyama G., Hoshita T.	127
Effect of insulin, S-adenosylhomocysteine, phospholipase C, n-butanol and Triton X-114 on alkaline phosphatase from isolated rat adipocyte plasma membranes Sykes E., Ghag S., Epstein E., Kiechle F.L.	133
Sandwich enzyme immunoassay of total insulin in the serum containing insulin antibodies Tominaga M., Honda M., Itoh Y., Mokuda O., Ikeda T., Mashiba H.	141
Problems and promises arising from measuring salivary testosterone Navarro M.A., Villabona C.M., Bonnin M.R.	143

Vol. 169/2-3, 1987

Increased excretion of propan-1,3-diol and 3-hydroxypropionic acid apparently caused by abnormal bacterial metabolism in the gut Pollitt R.J., Fowler B., Sardharwalla I.B., Edwards M.A., Gray R.G.F.	151
---	-----

Source of elevated serum mitochondrial creatine kinase activity in patients with malignancy Okano K., Yamamoto K., Ohba Y., Matsumura K., Miyaji T.	159
Alkaline phosphatase in human milk: A new heat-stable enzyme Chuang N.N.	165
A new assay for choline-containing phospholipids in amniotic fluid by an enzyme sensor Campanella L., Tomassetti M., De Angelis G., Sammartino M.P., Cordatore M.	175
Separation of 5'-nucleotide phosphodiesterase isoenzymes of human serum by anion exchange high pressure liquid chromatography Luthje J., Ogilvie A.	183
Comparison of plasminogen activator inhibitor activity and antigen in plasma samples Urden G., Hamsten A., Wiman B.	189
Quantitation of serum carotenoid concentrations in healthy inhabitants by high-performance liquid chromatography Ito Y., Sasaki R., Minohara M., Otani M., Aoki K.	197
Chromatographic profiles of urinary isoenzymes in healthy children Matsukura H., Suzuki Y., Takai R., Okada T., Naiki S., Sakuragawa N.	209
Sulfation of minoxidil by human platelet sulfotransferase Johnson G.A., Baker C.A.	217
Radioimmunoassay of glycosylated albumin with monoclonal antibody to glucitol-lysine Ohe Y., Matsuura M., Nakajima Y., Shin S., Hashimoto F., Hirota Shima M.K.	229
An enzyme immunoassay for pepsinogen II (PGII): chronological changes in serum PGII concentrations Matsusako K., Itoh M., Yokochi K., Miyamoto T., Joh T., Takeuchi T., Moriyama A., Kato T., Tanaka R.	239
Isolation and quantitation of apolipoproteins A-I and A-II from human high-density lipoproteins by fast-protein liquid chromatography Weisweiler P., Friedl C., Ungar M.	249
Evaluation by immunoblotting of the criteria for tubular lesions diagnostic with urinary SDS-PAGE Perini J.-M., Dehon B., Marianne Th., Foulard M., Roussel Ph.	255
Improved methods for the detection of unique sequences in Southern blots of mammalian DNA by non-radioactive biotinylated DNA hybridization probes Gregersen N., Koch J., Kolvraa S., Bruun Petersen K., Bolund L.	267
Characterization of human middle ear mucus glycoprotein in chronic secretory otitis media (CSOM) FitzGerald J.E., Green G.G.R., Stafford F.W., Birchall J.P., Pearson J.P.	281
Interactions of foetal steroid binding protein with other binding proteins in human serum Forbes A., Wilkinson M.L., Jawed Iqbal M., Williams R.	299
Faecal haem assay — some practical modifications of the haemoquant assay for haemoglobin in faeces Van den Berg J.W.O., Edixhoven-Bosdijk A., Koole-Lesuis R., Wilson J.H.P.	319
Decrease of serum malondialdehyde in patients treated with chlorpromazine Bindoli A., Rigobello M.P., Cavallini L., Dalla Libera A., Galzigna L.	329

Vol. 170/1, 1987

Biochemical and functional properties of protein C and protein S Strickland D.K., Kessler C.M.	1
Protein C and protein S clinical perspectives Kessler C.M., Strickland D.K.	25
Production, purification and characterization of antibodies to 1,25-dihydroxyvitamin D raised in chicken egg yolk Bauwens R.M., Kint J.A., Devos M.P., Van Brussel K.A., De Leenheer A.P.	37
Serum concentrations of a granulocyte-derived calcium-binding protein in cystic fibrosis patients and heterozygotes Hayward C., Glass S., Van Heyningen V., Brock D.J.H.	45

Low molecular mass trypsin inhibitors in normal and osteoarthritic human articular cartilage Lindenhayn K., Haupt R., Heilmann H.-H.	57
Neural tube defect-specific acetylcholinesterase: Its properties and quantitation in the detection of anencephaly and spina bifida Bonham J.R., Dale G., Atack J.R.	69
High performance liquid chromatography of plasma aminoacids using orthophthalaldehyde derivatisation Blundell G., Brydon W.G.	79
Modification of transferrin by polysaccharides in cancer patients Narita K., Kanamaru I., Makino Y., Konno K.	85
Effect of short-term digoxin treatment on ouabain-sensitive sodium transport of erythrocytes Smith J.B., Luft F.C., Wade M.B., Fineberg N.S., Weinberger M.H.	89
Sandwich enzyme immunoassay for human interleukin-1 α produced in vitro by peripheral blood mononuclear cells Tanaka K., Ishikawa E., Ohmoto Y., Hirai Y.	97
A simple assay for serum arylesterase using a specific substrate Tanaka H., Kitamura M., Inoue Y.	105
Zinc contamination due to the collection tube De Haan K.E.C., De Groot C.J., Boxma H., Van Den Hamer C.J.A.	111

Vol. 170/2-3, 1987

Identification and partial purification of a (Na-K)ATPase stimulating serine protease from plasma of insulin-dependent diabetics Finotti P., Verbaro R.	121
Altered calmodulin activity in buccal epithelial cells from cystic fibrosis patients McPherson M.A., Tiligada E., Bradbury N.A., Goodchild M.C.	135
Relationship of branched-chain α -keto-acids and lipids in sera of hypertriglyceridemic subjects Penttilä A.M., Herranen J., Lampainen E., Voutilainen E.	143
The quantification of gamma-aminobutyric acid in the cerebrospinal fluid by a radioreceptorassay Mousah H., Jacqmin P., Lesne M.	151
Low molecular weight thiol content in glutathione synthetase-deficient human fibroblasts Debieu D., Deschavanne P.J., Malaise E.P.	161
In vivo metabolism of apolipoprotein S in humans. Comparison with apolipoprotein A-I metabolism Malmendier C.L., Lontie J.-F., Delcroix C.	169
Characterization of non-TCA-precipitable 'Lowry protein' in plasma of patients with acute renal failure Haag M., Meyer H.E., Schollmeyer P., Horl W.H.	181
A new micro-assay for human liver alanine: Glyoxylate aminotransferase Allsop J., Jennings P.R., Danpure C.J.	187
Primary hyperparathyroidism and hyperuricaemia are associated but not correlated with indicators of bone turnover Broulik P.D., Stepan J.J., Pacovsky V.	195
Detection of alpha-1-antitrypsin deficiency variants by synthetic oligonucleotide hybridization Klasen E.C., Hofker M.H., Van Paassen H.M.B., Verlaan-De Vries M., Bos J.L., Frants R.R.	201
Stability of protein and mRNA in human postmortem liver-analysis by two-dimensional gel electrophoresis Finger J.M., Mercer J.F.B., Cotton R.G.H., Danks D.M.	209
Effect of bromelain on alkaline phosphatases of intestinal and non-intestinal tissues and serum Kalimanovska V., Whitaker K.B., Moss D.W.	219
Development of a reference method for determining urinary oxalate by means of isotope dilution-mass spectrometry (ID-MS) and its usefulness in testing existing assays for urinary oxalate Koolstra W., Wolthers B.G., Hayer M., Elzinga H.	227

An improved high performance liquid chromatographic method for determining urinary oxalate, making use of an ID-MS reference method Koolstra W., Wolthers B.G., Hayer M., Rutgers H.M.	237
Determination of transcortin in serum by polyethyleneglycol enhanced immunonephelometry. A comparison with equilibrium dialysis and radioimmunoassay Eriksson C.-G., Eneroth P.	245
An automated colorimetric assay for urine nicotine metabolites: a suitable alternative to cotinine assays for the assessment of smoking status Puhaikanen E.V.J., Barlow R.D., Salonen J.T.	255
Effect of long preincubation on the two forms of human erythrocyte prolidase Myara I.	263
Semi-automated enzyme-linked immunosorbent assay (ELISA) for the quantification of apolipoprotein B using monoclonal antibodies Bojanovski M., Gregg R.E., Wilson D.M., Brewer Jr. H.B.	271
Screening for adenine and hypoxanthine phosphoribosyltransferase deficiencies in human erythrocytes by high-performance liquid chromatography Sakuma R., Nishina T., Kitamura M., Yamanaka H., Kamatani N., Nishioka K.	281
Collagenase-like (CL) peptidase activity in synovial fluid from patients with rheumatoid arthritis Ito A., Hagihara M., Nagatsu T., Iwata H., Miura T.	291
Increased urinary excretion of argininosuccinate in type II citrullinemia Saeki T., Kobayashi K., Inoue I., Matuo S., Hagihara S., Noda T.	297
Use of GC/MS/SIM for rapid determination of plasma levels of o,p'-DDD, o,p'-DDE and o,p'-DDA Inouye M., Mio T., Sumino K.	305
A new radioimmunoassay of free thyroxine using ¹²⁵ I-labelled thyroxine-protein complex uninfluenced by albumin and thyroxine-binding globulin Tsutsumi S., Ishibashi K., Miyai K., Nagase S., Ito M., Amino N., Endo Y.	315
Accumulation of phosphorylated dolichol in several tissues in ceroid-lipofuscinosis (Batten disease) Hall N.A., Patrick A.D.	323
Pharmacokinetics of desferrioxamine and of its iron and aluminium chelates in patients on haemodialysis Allain P., Chaleil D., Mauras Y., Beaudeau G., Varin M.C., Poignet J.L., Ciancioni C., Ang K.S., Cam G., Simon P.	331
Correlation between qualitative and quantitative faecal occult blood tests Robertson J.D., Maughan R.J., Davidson R.J.L.	339
Increased urinary excretion of the retinol-binding protein in insulin-dependent diabetes mellitus in the absence of microalbuminuria Holm J., Hemmingsen L., Nielsen N.V., Thomsen M.	345
Hormone-induced redistribution of protein kinase C in human pituitary adenomas Ishizuka T., Ito Y., Murayama M., Miura K., Nagao S., Nozawa Y.	351

Vol. 171/1, 1988

A rapid and sensitive enzyme immunoassay for Cu/Zn superoxide dismutase with polyclonal and monoclonal antibodies Porstmann T., Wietschke R., Schmechta H., Grunow R., Porstmann B., Bleiber R., Pergande M., Stacht S., Von Baehr R.	1
Analysis of the fatty acid composition of erythrocyte phospholipids by a base catalysed transesterification method — Prevention of formation of dimethylacetals Laryea M.D., Cieslicki P., Diekmann E., Wendel U.	11
A sensitive, direct colorimetric assay of serum iron using the chromogen, nitro-PAPS Makino T., Kiyonaga M., Kina K.	19
Quantitation and characterization of urinary glycosaminoglycans in healthy subjects and patients with mastocytosis Kettelhut B.V., Metcalfe D.D.	29

Development of fluoroimmunoassays for the specific detection of morphine in urine Colbert D.L., Gallacher G., Ayling P., Turner G.J.	37
A new method for the determination of L-DOPA and 3-O-methyldopa in plasma and cerebrospinal fluid using gas chromatography and electron capture negative ion mass spectrometry De Jong A.P.J.M., Kok R.M., Cramers C.A., Wadman S.K., Haan E.	49
Organ specific properties for human urinary alkaline phosphatases Miura M., Koyama I., Matsuzaki H., Sakagishi Y., Ikezawa H., Komoda T.	63
Red cell pyrimidine 5' nucleotidase deficiency: Determination of nucleotidase activity and nucleotide content using HPLC Adair C.G., Elder G.E., Lappin T.R.J., Bridges J.M., Nelson M.G.	75
Effect of renal insufficiency on the concentration of free retinol-binding protein in urine and serum Bernard A., Vyskocyl A., Mahieu P., Lauwerys R.	85
3-Hydroxy-3-methylglutaryl-CoA lyase in human skin fibroblast: Study of its properties and deficient activity in 3-hydroxy-3-methylglutaric aciduria patients using a simple spectrophotometric method Wanders R.J.A., Schutgens R.B.H., Zoeters P.H.M.	95
Changes in the bone and liver isoenzymes of alkaline phosphatase in postmenopausal women being treated with norethisterone Deeny M., Farish E., Tillman J., Dagen M., Hart D.M., Fletcher C.D.	103
Estimation of pyruvate dehydrogenase (E ₁) activity in human skeletal muscle; three cases with E ₁ deficiency Van Laack H.L.J.M., Ruitenbeek W., Trijbels J.M.F., Sengers R.C.A., Gabreels F.J.M., Janssen A.J.M., Kerkhof C.M.C.	109

Vol. 171/2-3, 1988

Flow cytometry in the clinical laboratory Ryan D.H., Fallon M.A., Horan P.K.	125
Combined enzymatic assay of phosphatidylglycerol and phosphatidylcholine in amniotic fluid Farquharson J., Jamieson E.C., Berry E., Sutton A., Logan R.W.	175
Separation and partial characterization of two galactosyltransferase isoforms from malignant ascitic fluid Boyle F.A., Cook N.D., Peters T.J.	187
Altered erythrocyte nucleotide patterns are characteristic of inherited disorders of purine or pyrimidine metabolism Simmonds H.A., Fairbanks L.D., Morris G.S., Webster D.R., Harley E.H.	197
GC and GC-MS procedures for simultaneous phenotyping with dextromethorphan and mephentermine Baumann P., Jonzier-Perey M.	211
Stable isotope dilution analysis of succinylacetone using electron capture negative ion mass fragmentography: An accurate approach to the pre- and neonatal diagnosis of hereditary tyrosinemia type I Jakobs C., Dorland L., Wikkink B., Kok R.M., De Jong A.P.J.M., Wadman S.K.	223
Assessment of taurine deficiency in cystic fibrosis Thompson G.N.	233
The effect of renal failure and haemodialysis on the concentration of free apolipoprotein A-1 in serum and the implications for the catabolism of high-density lipoproteins Neary R.H., Gowland E.	239
Optimization of histamine radio enzyme assay with purified histamine-N-methyltransferase Harvima R.J., Harvima I.T., Fraki J.E.	247
Newborn screening for cystic fibrosis using an enzyme linked immunosorbent assay (ELISA) technique Bowling F.G., Brown A.R.D.	257

The characterisation of molecular forms of acetylcholinesterase in Hirschsprung's disease Bonham J.R., Dale G., Scott D.J., Wagget J., Atack J.R.	263
Microheterogeneity of paraproteins. I. Diagnostic value of isoelectric focusing followed by immunoblotting Schipper H.I., Bertram G., Kaboth U.	271
Purine metabolism in normal and ITP-pyrophosphohydrolase-deficient human erythrocytes Van Waeg G., Niklasson F., Ericson A., De Verdier C.-H.	279
Effect of insulin treatment on fatty acids of plasma and erythrocyte membrane lipids in type 2 diabetes Tilvis R.S., Taskinen M.-R., Miettinen T.A.	293
Apoprotein AII in acute pancreatitis: Intriguing improvement in the prediction of fatal outcome Schneider J., Meier M., Maroake D., Kaffarnik H.	305
A simple fluorometric method for the determination of serum free carnitine Maehara M., Kinoshita S., Watanabe K.	311
Improved purification of human liver alkaline phosphatase by phenyl-Sepharose column chromatography Takata K.-I., Sumikawa K., Saeki K., Okochi T., Adachi K.	317
An alpha-fetoprotein radioimmunoassay for use in maternal serum screening for Down's syndrome: Limitations and improvements Mooney R.A., McGorty B.L., French C.A., Arvan D.A.	325

Vol. 172/1, 1988

Laboratory analyses for steroid hormone receptors, and their applications to clinical medicine Mayer T.K., Mooney R.A.	1
A sensitive kinetic assay of serum albumin based on its enzyme-like hydrolytic activity, using a new chromogenic and fluorogenic substrate Gurakar A.R., Wolfbeis O.S.	35
Fetal lung maturity tests on the 10000 x g pellet McCulloch J.C., Mendelsohn D.	47
Sialic acid concentration in erythrocyte membrane subfractions in patients with myotonic dystrophy and healthy controls Sydow O.	57
Molecular analysis of peroxisomal β -oxidation enzymes in infants with Zellweger syndrome and Zellweger-like syndrome: Further heterogeneity of the peroxisomal disorder Suzuki Y., Shimozaawa N., Orii T., Igarashi N., Kono N., Hashimoto T.	65
Re-evaluation of normal human urinary proteins fractionated by one-dimensional sodium dodecyl sulphate-polyacrylamide gel electrophoresis Shiba K.S., Kanamori K., Cho H., Furuhashi N., Harada T., Shiba A., Nakao M.	77
Neutrophil chemiluminescence induced by platelet activating factor and suppressed by C-reactive protein Tatsumi N., Hashimoto K., Okuda K., Kyougoku T.	85
Determination of cytochromes in human liver contaminated with hemoglobin Tanaka A., Maki A., Kamiyama Y., Yamaoka Y., Ozawa K., Orii Y.	93
A new method of inhibiting glycolysis in blood samples Uchida K., Matuse R., Toyoda E., Okuda S., Tomita S.	101
Concanavalin A- or phorbol ester-induced translocation of protein kinase C in thymoma cells from a patient with myasthenia gravis Ishizuka T., Ito Y., Mori H., Miura K., Nagao S., Nagata K.-I., Nozawa Y.	109
Rapid high-performance liquid chromatography for pseudouridine assay in serum and urine Amuro Y., Nakaoka H., Shimomura S., Tamura S., Hada T., Higashino K.	117
Elevated excretion of N-acetylated branched-chain amino acids in maple syrup urine disease Lehnert W., Werle E.	123

Vol. 172/2-3, 1988

An enzymatic method for the α -amylase assay which comprises a new procedure for eliminating glucose and maltose in biological fluids	
Kondo H., Shiraishi T., Nagata K., Tomita K.	131
Characterization of an adenine phosphoribosyltransferase deficiency	
Chiba P., Zwiauer K., Muller M.M.	141
Electroimmunoassay of human serum lipoproteins containing Apo A-I by monoclonal antibodies	
Ayrault-Jarrier M., Petit E., Bekaert E.D., Robin H., Pastier D., Beucier I., Polonovski J., Hervaud E., Pau B.	149
A rapid electrochemical assay of lecithin in amniotic fluid using a fluoride ion-sensitive electrode	
Diomedea L., Masturzo P., Agosti S., Ornaghi F., Salmona M.	161
Enzyme-linked immunosorbent assay for ileal human goblet cell mucin	
Robertson A.M., Dalzell H., Harding C.A., Pearce S., Ram S.P., McKay E.J., Wesley A.W.	171
Direct quantitation of serum high density lipoprotein subfractions separated by gradient gel electrophoresis	
Gambert P., Farnier M., Bouzerand C., Athias A., Lallemand C.	183
Urinary albumin excretion in healthy adult subjects: Reference values and some factors affecting their interpretation	
Watts G.F., Morris R.W., Khan K., Polak A.	191
The presence and molecular forms of cardiolipin immunoreactivity in the human and rat right atrium	
Meleagros L., Ghatei M.A., Anderson J.V., Wharton J., Taylor K.M., Krikler D.M., Meijler F.L., Polak J.M., Bloom S.R.	199
Glutathione S-transferase levels in autoimmune chronic active hepatitis: A more sensitive index of hepatocellular damage than aspartate transaminase	
Hayes P.C., Hussey A.J., Keating J., Bouchier I.A.D., Williams R., Beckett G.J., Hayes J.D.	211
Thyroid dysfunction and hypercholesterolaemia in the general population of Glasgow, Scotland	
Series J.J., Biggart E.M., O'Reilly St. D.J., Packard C.J., Shepherd J.	217
Anomeric specificity of D-glucose metabolism in human erythrocytes	
Malaise-Lagae F., Sterling I., Sener A., Malaise W.J.	223
Determination of medium chain fatty acids in serum	
Vici C.D., Bachmann C., Gradwohl M., Colombo J.P.	233
Whole blood storage solution for erythrocyte sodium and sodium-lithium countertransport rate determination	
Siebers R.W.L., Maling T.J.B.	239
The isolation and identification of indigoid pigments from urine	
Jackson A.H., Jenkins R.T., Grinstein M., Ferramola de Sancovich A.-M., Sancovich H.A.	245
Identification of the carboxypeptidase responsible for the post-synthetic modification of creatine kinase in human serum	
Hendriks D., Soons J., Scharpe S., Wevers R., Van Sande M., Holmquist B.	253
The distinction of bone and liver isoenzymes of alkaline phosphatase in serum using a monoclonal antibody	
Seabrook R.N., Bailyes E.M., Price C.P., Siddle K., Luzio J.P.	261
A monoclonal antibody capture assay for intestinal alkaline phosphatase and the measurement of this isoenzyme in pregnancy	
Bailyes E.M., Seymour P.M., Fulton I., Price C.P., Luzio J.P.	267
Fatty acid composition and kinetic behaviour of liver retinyl esters in vitamin A sufficient and deficient rats	
Periquet B., Lambert W., Bailly A., Tomatis I., Ghisolfi J., De Leenheer A.P., Thouvenot J.P.	275
Inhibition of galactosyltransferase activity by cacodylate buffer	
Boyle F.A., Cook N.D., Peters T.J.	291
Application of heparin-conjugated Sepharose for the measurement of cobalamin-saturated and unsaturated transcobalamin II	
Van Kapel J., Wouters N.M.H., Lindemans J.	297

Analysis of cell surface glycoconjugates in fibroblasts from patients with cystic fibrosis Sarras Jr. M.P., Huff J.K.	311
Presence of activator proteins for the enzymatic degradation of glucosylceramide in several human tissues Vaccaro A.M., Ciaffoni F., Mandara I., Suzuki K.	323
Role of serum pepsinogen in detecting ulcer disease Vianello F., Plebani M., Piccoli A., Tessaro P., Di Mario F., Naccarato R., Burlina A.	335

Vol. 173/1, 1988

The place of reference materials in clinical enzymology Moos D.W.	1
Standardization by reference methods: A German viewpoint Schmidt E.S., Schmidt F.W.	9
γ -Glutamyltransferase reference material: An example of international cooperation Siest G., Schiele F., Muller J., Wellman M.	19
Molecular titration as a means of calibrating enzyme reference materials Roth M., Selz L.	27
Influence of isoenzyme patterns on commutability in enzyme determinations Burlina A., Bonvicini P., Plebani M., Zaninotto M.	35
Glutamate dehydrogenase: Biochemical and clinical aspects of an interesting enzyme Schmidt E.S., Schmidt F.W.	43
Clinical biochemistry of peroxisomal disorders Kaiser E., Kramar R.	57
Esterase D: Evaluation of a potential derived gene marker for hereditary retinoblastoma Van Der Heiden C., Geurtzen A.F.M., Brink W., De Klerk J.B.C., Beemer F.A.	81
Lactate dehydrogenase M-subunit deficiency: A new type of hereditary exertional myopathy Kanno T., Sudo K., Maekawa M., Nishimura Y., Ukita M., Fukutake K.	89

Vol. 173/2, 1988

Determination of zinc in blood serum by electrothermal atomic absorption spectrometry with matrix modification Accominotti M., Pegon Y., Vallon J.-J.	99
Identification of renal calculi by computerized infrared spectroscopy Lehmann C.A., McClure G.L., Smolens I.	107
The estimation of dihydropteridine reductase in human blood cells Webber S., Hural J.A., Whitely J.M.	117
Accumulation of furancarboxylic acids in uremic serum as inhibitors of drug binding Niwa T., Takeda N., Maeda K., Shibata M., Tatematsu A.	127
Similarity of the sugar moiety of human alkaline phosphatases between the kidney cortex and duodenum, or medulla and ileum Koyama I., Arai Y., Miura M., Matsuzaki H., Sakagishi Y., Komoda T.	139
Interleukin-2-dependent T lymphocytes for the diagnosis and investigation of inherited metabolic disorders Adolph K., Kucey M.T., Hodges S.D., Carter R.J., Snyder F.F.	147
Diagnostic significance of estimation of serum apolipoprotein a along with α -fetoprotein in alcoholic cirrhosis and hepatocellular carcinoma patients Nayak S.S., Kamath S.S., Kundaje G.N., Aroor A.R.	157
In vitro glycation of dried serum Ch'ng S.L., Marinah T.A.	165
Alkaline phosphatase isoenzyme pattern in human amniotic fluid is dependent on the level of total activity. Implications in cystic fibrosis diagnosis Dictus-Vermeulen C., Ameryckx J., Gueuning C., Van Bogaert E., Graff G.L.A.	173

A new microtechnique for the analysis of the human hepatic microsomal glucose-6-phosphatase system	
Burchell A., Hume R., Burchell B.	183
Enzyme pattern and lipid peroxides in endomyocardial biopsies from patients with cardiomyopathy and myocarditis	
Lehmann I., Papies B., Parsi R.A., Romaniuk P., Schimke I., Parsi E., Konig M.-L.	193
Comparison of separation techniques in radioimmunoassays for 17-hydroxyprogesterone	
El-Gamal B.A., Landon J.	201
Monoclonal IgMx with rheumatoid factor activity and cryoprecipitability identified only by immunofixation electrophoresis after 2-mercaptoethanol treatment	
Sorensen S.	217
Agarose isoelectric focusing of arylsulfatase A producing a permanent record on slab gels.	
Menna-Perper M., Nora R., Goldstein L., Manowitz P.	225
Pterins as biochemical tests for diagnosis and management of patients with malignant lymphoma and leukemia	
Reibnegger G., Fuchs D., Hausen A., Werner E.R., Wachter H.	233
Impaired production of leukotrienes by peripheral blood polymorphonuclear leukocytes from children with chronic renal failure	
Hamasaki T., Sakano T., Kobayashi M., Sakura N., Usui T., Ueda K.	235
The relationship between serum fructosamine and albumin	
Senecal P.-E., Douville P., Simard S., Coulombe R.	239

Vol. 173/3, 1988

Minimal myocardial injury after uncomplicated coronary bypass surgery. Various sources of overestimation	
Hermens W.T., Wilems G.M., Van Der Vusse G.J.	243
Early diagnosis of clostridial gas gangrene using sialidase antibodies	
Roggentin P., Gutschker-Gdaniec G.H.M., Hobrecht R., Schauer R.	251
Urinary acylcarnitines in a patient with neonatal multiple acyl-CoA dehydrogenation deficiency, quantified by a carboxylic acid analyzer with a reversed-phase column	
Kidouchi K., Niwa Nohara T.D., Asai K., Sugiyama N., Morishita H., Kobayashi M., Wada Y.	263
Reference limits of bone and liver alkaline phosphatase isoenzymes in the serum of healthy subjects according to age and sex as determined by wheat germ lectin affinity electrophoresis	
Kuwana T., Sugita O., Yakata M.	273
A high-performance liquid chromatographic assay for acid ceramidase activity in cultured fibroblasts from patients with Farber's disease and from controls	
Mitsuo K., Kobayashi T., Shinnoh N., Goto I.	281
Reactivities of human urinary prokallikrein and kallikrein toward rabbit antibodies with special reference to enzyme immunoassay	
Irie A., Takahashi S., Katayama Y., Shibata Y., Miyake Y.	289
Random distribution of iron among the two binding sites of transferrin in patients with various hematologic disorders	
Beguín Y., Huebers H., Finch C.A.	299
Succinic acidemia: a new syndrome of organic acidemia associated with congenital lactic acidosis and decreased NADH-cytochrome c reductase activity	
Asano K., Miyamoto I., Matsushita T., Murakami Y., Minouri S., Wagatsuma T., Oshima M.	305
Pharmacokinetics of desferrioxamine and of its iron and aluminum chelates in patients on peritoneal dialysis	
Allain P., Mauras Y., Varin M.C., Ang K.S., Cam G., Simon P.	313
Biochemical investigations on prolidase and prolinsase in erythrocytes from patients with prolidase deficiency	
Kodama H., Mikasa H., Ohhashi T., Ohno T., Arata J.	317

Correlation between serum lipid peroxides and the lesion size in cerebrovascular disease Huang Z.-S., Lu F.-J., Lee T.-K.	325
The carbohydrate composition of human serum amyloid P component Hess D., Ohishi H., Skinner M., Cohen A.S., Schmid K.	331
The chemiluminescence response of polymorphonuclear leukocytes from febrile patients Miyata H., Moriguchi N., Kinoshita T.	337
Lack of association between plasma lecithin: cholesterol acyltransferase concentration and plasma sex hormone concentrations in men Nanjee M., Albers J.J., Miller N.E.	343

Vol. 174/1, 1988

Methyltransferase and phospholipase A ₂ activity in the cell membrane of neutrophils and lymphocytes from patients with Behcet's disease, systemic lupus erythematosus, and rheumatoid arthritis Niwa Y., Miyachi Y., Sakane T., Kanoh T., Taniguchi S.	1
Elimination of undesirable immunoglobulin contaminants including aggregated IgG from γ -globulin preparations by jackfruit lectin affinity chromatography Kondoh H., Kobayashi K.	15
Characterization of amylase linked immunoglobulin G to distinguish human salivary and pancreatic isoamylases Moriyama T., Takebe T., Nobuoka M., Makino M.	25
Construction and clinical validation of a sensitive and specific assay for serum mature calcitonin using monoclonal anti-peptide antibodies Motte P., Vauzelle P., Gardet P., Ghillani P., Caillou B., Parmentier C., Bohuon C., Bellet D.	35
DSC, X-ray and NMR properties of cholesterol-cholestanol-dihydrate crystals Hirsch D., Azoury R., Sarig S.	55
Applications of immunoperoxidase techniques in specificity testing of monoclonal antibodies (Mabs) against von Willebrand factor (vWf) Ingerslev J., Bukh A., Moller N.P.H., Stenbjerg S., Zeuthen J.	65
Enzyme immunoassay of cortisol in human plasma using penicillinase as label Shrivastav G.S., Kumari G.L., Rao P.N.	83
Modified dual label Schilling test for pancreatic exocrine function Leung J.W.C., Frost R.A., Burgess R., Braganza J.M., Slater D.M., Cotton P.B.	93
The Philadelphia variant of galactokinase: Impaired [1- ¹⁴ C]galactose oxidation by intact erythrocytes Soni T., Brivet M., Moatti N., Dahan N., Lemonnier A.	101
Effect of insulin on globin synthesis in human reticulocytes Ballas S.K., Tran T.H.	111
Magnesium, zinc, calcium, potassium, lipoproteins and creatine kinases on the first day after acute myocardial infarction: Statistical analysis Auger J.-L., Speich M., Arnaud P.	115

Vol. 174/2, 1988

A sensitive method to assay blood complement C \bar{I} ; Inhibitor activity Drouet C., Alibeu C., Ponard D., Arlaud G.J., Colomb M.G.	121
A comparative study on the diagnostic value of prostatic acid phosphatase (PAP) and prostatic specific antigen (PSA) in patients with carcinoma of the prostate gland Van Dieijen-Visser M.P., Delaere K.P.J., Gijzen A.H.J., Brombacher P.J.	131
Ca ²⁺ influx in normal and spherocytic red cells Johnsson R., Saris N.-E.	141
Monoclonal antibodies to human kidney gamma-glutamyltransferase Donati R., Sabolovic N., Wellman M., Artur Y., Siest G.	149

Evaluation of serum aminoterminal procollagen type III propeptide as an index of portal hypertension and esophageal varices in chronic liver diseases Gressner A.M., Tittor W., Kropf J.	163
Quality of performance of assays for serum growth hormone in the United Kingdom (UK): Evidence from the UK external quality assessment scheme, 1980-1987 Seth J., Hanning I., Bacon R.R.A., Hunter W.M.	171
Factors associated with the quality of laboratory performance in the United Kingdom external quality assessment scheme for serum growth hormone Seth J., Hanning I.	185
Plasma and biliary cholestanol related to steroid metabolism in familial hypercholesterolemia patients with and without ileal exclusion Koivisto P.V.I., Miettinen T.A.	197
Quantification of soluble serum HLA Class I antigens in healthy volunteers and AIDS patients Ferreira A., Villar M.L., Alvarez Cermeno J.C., Revilla Y., Garcia Rodriguez M.C., Fontan G., Gonzalez-Porque P.	207
Serum plant sterols and lathosterol related to cholesterol absorption in coeliac disease Vuoristo M., Tilvis R., Miettinen T.A.	213
A new and rapid immunochemiluminometric assay for the measurement of Tamm-Horsfall glycoprotein Horton J.K., Davies M., Woodhead J.S., Weeks I.	225

Vol. 174/3, 1988

Renal handling of cathodic trypsin-like immunoreactivity in man Moller-Petersen J., Ladefoged J.	241
Albumin determination in frozen urines — Underestimated results Erman A., Rabinov M., Rosenfeld J.	255
On the fluorometric assay of circulating lipoperoxides Hendriks T., Assman R.F.T.A.	263
Sepiapterin reductase in human amniotic and skin fibroblasts, chorionic villi, and various blood fractions Ferre J., Naylor E.W.	271
An altered calmodulin binding protein in cystic fibrosis — A clue to the biochemical defect Shori D.K., Bradbury N.A., Goodchild M.C., Dormer R.L., McPherson M.A.	283
Effect of chronic ethanol ingestion on alpha-mannosidase isoenzymes in rat liver Lauro P., Lechner P.S., Okolo A., Eagon P.K., Glew R.H.	291
Distribution of tocopherol among human plasma lipoproteins Ogihara T., Miki M., Kitagawa M., Mino M.	299
Enzymatic analysis of total- and HDL-cholesterol: Comparison with the standardized Liebermann-Burchard method used by the lipid research clinics program Bachorik P.S., Virgil D.G., Derby C., Widman D., McMahon R., Fulwood R.P., Ezzati T.	307
α -Amylase assay with use of a benzyl derivative of p-nitrophenyl α -maltopentaoside, BG5P Satomura S., Sakata Y., Omichi K., Ikenaka T.	315
A comparative study of the accumulated sialic acid-containing oligosaccharides from cultured human galactosialidosis and sialidosis fibroblasts Van Pelt J., Kamerling J.P., Vliegthart J.F.G., Hoogeveen A.T., Galjaard H.	325
Urinary purines, pyrimidines and nucleosides in uridine-treated orotic aciduria Mills G.C.	337
A pepstatin-immunometric assay for active renin in human plasma Higaki J., Ogihara T., Nishiura M., Murakami K., Kumahara Y.	345
Stabilization of ascorbic acid in plasma Khanduja K.L., Koul A.	351

Vol. 175/1, 1988

Serum lysosomal hydrolases in cystic fibrosis	
Krall E.A., Basu A., Gloninger M.F., Glew R.H., Humphries L.	1
Enzyme immunoassay for α -human atrial natriuretic polypeptide — Direct measurement of plasma level	
Hashida S., Ishikawa E., Nakao K., Mukoyama M., Imura H.	11
3-Hydroxyoctanoic aciduria: Identification of a new organic acid in the urine of a patients with non-ketotic hypoglycemia	
Kelley R.L., Morton D.H.	19
Studies on serum immunoreactive prolyl 4-hydroxylase in liver disease — Its elevation both in hepatocellular damage and cholestatic disease	
Nagai Y., Yoshida M.	27
Enzyme-linked immunosorbent assays for serum pepsinogens I and II using monoclonal antibodies — With data on peptic ulcer and gastric cancer	
Huang S.C., Miki K., Furihata C., Ichinose M., Shimizu A., Oka H.	37
Homogeneous immunoassay based on dihexadecyl-dimethylammonium bilayer membranes	
Ihara H., Imamura T., Kunitake T., Hirayama C.	51
Assay of stable isotope labelled urea in biological fluids by selected ion monitoring	
Nelson J.E., Ruo T.I.	59
Criteria for identifying endogenous compounds as digoxin-like immunoreactive factors in humans	
Lau B.W.C., Valdes Jr. R.	67
Urinary C ₆ -C ₁₂ dicarboxylic acylcarnitines in Reye's syndrome	
Tracey B.M., Cheng K.N., Rosankiewicz J., Stacey T.E., Chalmers R.A.	79
Oestrogen receptors, lactate dehydrogenase and cellularity in human breast cancer	
Hawkins R.A., Killen E., Teedale A.L., Sangster K., Thomson M., Steele R.J.C., Blackie R.A.S.	89
The Philadelphia variant of galactokinase in human erythrocytes: Physicochemical and catalytic properties	
Soni T., Brivet M., Moatti N., Lemonnier A.	97
Total and free quinidine by fluorescence polarization immunoassay and comparison with high performance liquid chromatography	
Chen B.H., Taylor E.H., Kennedy E., Ackerman B., Olsen K., Pappas A.A.	107
Creatine kinase BB and neuron-specific γ -enolase are enhanced in leiomyosarcoma	
Usui A., Kato K., Abe T., Hachisuka K., Yamaguchi A.	109
Measurement of platelet factor 4 and β -thromboglobulin by an enzyme-linked immunosorbent assay	
Takahashi H., Yoshino N., Shibata A.	113

Vol. 175/2, 1988

Isoelectric focusing of human neutrophil alkaline phosphatase isoenzyme in agarose gel	
A. Brisson-Lougarre, H. Vergnes, J. Grozdea, R. Biorme	121
Behaviour of L- γ -glutamyl-4-nitroanilide and L- γ -glutamyl-3-carboxy-4-nitroanilide with respect to γ -glutamyltransferases of different origin	
P.M.M. Verhoeff, F.P.A.M.N. Peters, H. Steigstra, J.C.M. Hafkenscheid	129
Clinical utility of serum fructosamine in diabetes mellitus compared with hemoglobin A _{1c}	
P. Jerntrop, G. Sundkvist, G. Fex, J.-O. Jeppsson	135
Determination of cholic acid and chenodeoxycholic acid pool sizes and fractional turnover rates by means of stable isotope dilution technique, making use of deuterated cholic acid and chenodeoxycholic acid	
B.J. Koopman, F. Kuipers, C.M.A. Bijleveld, J.C. Van der Molen, G.T. Nagel, R.J. Vonk, B.G. Wolthers	143
Tumour markers in the cerebrospinal fluid of patients with central nervous system metastases from extracranial malignancies	
A.P. van Zanten, A. Twijnstra, B.W. Ongerboer de Visser, A.A.M. Hart, W.J. Nooyen	157

Kinetic studies on nitrite and nitrate in rats by ion-pair chromatography C. Stein, H.-G. Classen, G. Schwedt	167
A more simple, rapid and sensitive fluorimetric method for the determination of isoniazid and acetylisoniazid in serum. Application for acetylator phenotyping P.C. Ioannou	175
Concentration differences in isoforms of transferrin in blood from alcoholics during abuse and abstinence S. Petré, O. Vesterberg	183
gamma-Glutamyl-transpeptidase in diabetics: a case control study G. Colledro-Mels, G. Bettale, G. Bellati, M. Guanziroli, V. Bertone, G. Angeli, G. Idéo	189
<i>Letter to the Editor</i>	
Colorimetric measurement of plasma 3-hydroxybutyrate J.F. Buttery, B.R. Chamberlain	197

Vol. 175/3, 1988

Factors influencing the production of creatinine: implications for the determination and interpretation of urinary creatinine and creatine in man N.J. Fuller, M. Elia	199
Plasma and serum zinc concentrations: effect of time between collection and separation J.L. English, K.M. Hambidge	211
Fast functional assay of protein C in whole plasma using a snake venom activator: evaluation in patients with congenital and acquired protein C deficiencies H. Takahashi, M. Hanano, W. Tatewaki, A. Shibata	217
The regional variability of enzymes in the brain: relevance to CSF enzyme determinations D.A. Armbruster, H.-D. Gruemer	227
Simultaneous ion-pair and partition liquid chromatography of acetaminophen, theophylline and salicylate with application to 500 toxicologic specimens J. Osterloh, S. Yu	239
Effects of acetaldehyde on human red cell metabolism: evidence for the formation of enzyme inhibitors L.R. Solomon	249
A sensitive time-resolved immunofluorometric assay of ferritin in serum with monoclonal antibodies M.J. Khosravi, M.A. Chan, A.C. Bellem, E.P. Diamandis	267
Lipoprotein cholesterol concentrations in patients with Behçet's disease T. Mitamura, S. Ohno, H. Ariga, T. Ohsaka, N. Iwasaki, H. Matsuda, H. Matsumiya	277
Doxorubicin-induced altered glycolytic patterns in the leukemic cell studied by proton spin echo NMR J. Reglinski, W.E. Smith, C.J. Suckling, M. Al-Kabban, M.J. Stewart, I.D. Watson	285
Plasma prolidase and prolinase activity in alcoholic liver disease B. Brosset, I. Myara, M. Fabre, A. Lemonnier	291
Development of a simple, fluoroimmunoassay for paraquat R.C. Coxon, C. Rae, G. Gallacher, J. Landon	297
In vivo and in vitro ionized calcium variations induced by acute respiratory acid base disturbances C.B. Kancir, P.H. Petersen, T. Madsen, A.S. Olesen	307
Measurement of endotoxin in cerebrospinal fluid from neonates and infants using a new endotoxin-specific chromogenic test M. Inagaki, M. Nakajima, Y. Ando, S. Takashima, K. Takeshita, S. Tanaka, H. Tamura, T. Obayashi	315
Detection and purification of an abnormal protein from plasma of patients with rheumatoid arthritis O. Oda, K. Maeda, H. Yamaga, Y. Kanamono, H. Iwata, T. Miura, M. Yoshida, S. Miyata, K. Ohta, N. Yamanaka	321

Stability of vitamin E in long-term stored serum E.W. Gunter, W.J. Driskell, P.R. Yeager	329
An enzyme antigen immunoassay for the determination of neuron-specific enolase in serum samples K. Sorensen, U. Brodbeck, E. Paus, B. Norgaard-Pedersen	337
<i>Letters to the Editor</i>	
Diagnosis of Zellweger syndrome by rectal biopsy: immunoblot of peroxisomal β -oxidation enzyme and activity of dihydroxyacetone phosphate acyltransferase in rectal mucosa N. Shimozaawa, Y. Suzuki, T. Orii, J. Hashimoto	345
The frequency of C ₃ + cholinesterase in the normal Japanese population M. Oimomi, J. Ohkawa, S. Saeki, S. Baba	349

Vol. 176/1, 1988

Circadian and circannual rhythms of several enzymes of lysosomal origin in human plasma G. Goi, A. Fabi, A. Lombardo, A.B. Burlina, G. Tettamanti, N. Montalbetti, M. Cavalleri, F. Halberg	1
Modification in Ramsay's method for correct measurement of total iron-binding capacity H. Baadenhuijsen, L.G.J. Deimann, A.P. Jansen	9
An enzymatic method for the determination of butyrobetaine via conversion to carnitine after isolation by high performance liquid chromatography A. Sandor, P.E. Minkler, S.T. Ingalls, C.S. Hoppel	17
Composition and surface properties of the bronchial liquid in adult patients with cystic fibrosis H. Gilljam, O. Anderson, Å. Ellin, B. Robertson, B. Strandvik	29
Urinary excretion of a glucose-containing tetrasaccharide. A parameter for increased degradation of glycogen J. Kumlien, M.A. Chester, B.S. Lindberg, P. Pizzo, D. Zopf, A. Lundblad	39
Modification of Concanavalin A patterns of α 1-acid glycoprotein and α 2-HS glycoprotein in alcoholic liver disease M. Jezequel, N.S. Seta, M.M. Corbic, J.M. Feger, G.M. Durand	49
Acute phase reactants in the elderly N. Milman, N. Graudal, H.C. Anderson	59
Determination of phospholipids in biological samples by an improved densitometric method on thin-layer chromatograms R. Masella, A. Cantafora	63
Evaluation of urinary oestrogen assays after the menopause and their potential for screening M.H. Rae, P.A. Mole, C.R. Paterson	71
Effect of methionine loading and endogenous hypermethioninaemia on blood mercaptans in man H. Al Mardini, J. Leonard, K. Bartlett, S. Lloyd, C.O. Record	83
Antibody immobilization on swollen polystyrene tubes for the radioimmunological determination of total human serum IgE J.E. Causse, I. Ricordel, P. Taulelle, M.T. Chateau, F. Moussel, A.M. Dupuy, B. Descomps, J.P. Yver	91
Plasma fructosamine assay in children with insulin-dependent diabetes mellitus N. Shimojo, M. Hirai, K.I. Naka, C. Yoshikawa, K. Okuda, S. Aono, G. Ishiki, S. Fuji	101
Diamine and polyamine levels in the urine of healthy adults Y. Yodiat, M. Weiser, M. Kreisel, U. Bachrach	107
I-cell disease: evidence for a mannose 6-phosphate independent pathway for translocation of lysosomal enzymes in lymphoblastoid cells A. Tsuji, K. Omura, Y. Suzuki	115

Vol. 176/2, 1988

Selection and performance of monoclonal anti-C-reactive protein in ELISA quantitative assay J.A. Martinez, J.M. Coll	123
---	-----

Use of ^{86}Rb and ^{22}Na in assaying active and cotransport activities in human erythrocytes in a biracial population	
C. Sharma, E.R. Dalferes Jr., D.S. Freedman, A. Asamoah, G.S. Berenson	133
Immunoaffinity purification of human erythrocyte prolidase	
F. Endo, A. Tanoue, T. Ogata, K. Motohara, I. Matsuda	143
IgE pyroglobulinemia in multiple myeloma	
M.L. Gallango, R. Suinaga, M. Ramirez	151
Validation of a radioreceptor assay for 1,25-dihydroxyvitamin D using selected ion monitoring GC-MS	
H. Oftebro, J.A. Falch, I. Holmberg, E. Haug	157
Chromatography of serum on Sep-pak C_{18} corrects falsely elevated vitamin D metabolic levels measured by protein binding assay	
B. Dean, M.S. Kolavcic, J.D. Wark, L.C. Harrison	169
Characterisation of a slow component of normal human serum albumin	
S.O. Brennan, P.M. George, R.J. Peach	179
A kinetic procedure for the estimation of arginine in serum using arginine kinase	
C.H. Konings	185
Changes of serum apo A-IV levels in a newborn with chylothorax treated with medium-chain triglyceride milk	
M. Scishima, A. Noma, S. Yamaguchi, T. Orii, Y. Muto	195
Serum indole and skatole in patients with various liver diseases	
Y. Suyama, C. Hirayama	203
Measurement of myo-inositol in diabetic sera by GC/MS/SIM using <i>n</i> -butylboronate derivatives	
M. Hasegawa, K. Dol, S. Baba	207
Enzyme-linked immunosorbent assay (ELISA) for human tumor necrosis factor (hTNF)	
Y. Abe, M. Miyake, T. Sagawa, S. Kimura	213
Elastase-like protease and elastolytic activities expressed in cultured dermal fibroblasts derived from lesional skin of patients with pseudoxanthoma elasticum, actinic elastosis, and cutis laxa	
E. Schwartz, F.A. Cruickshank, M.G. Lebowitz	219
<i>Letters to the Editor</i>	
Effect of serum albumin concentration on free thyroxine (Amerlex FT4) values in healthy non-pregnant and pregnant women	
S. Sakata, T. Komaki, S.-I. Shiraki, N. Kumasaki, H. Iwata	225
A pitfall in the analysis of the fatty acid composition of plasma and erythrocyte membrane lipids	
C. Popp-Snijders	227
Evaluation of sex hormone binding globulin among patients with thyroxine binding globulin excess	
J. Wortsman, B.N. Premachandra, I.K. Williams, W. Rosner	229
Separation and purification of creatine kinase BB isoenzyme and neuron-specific enolase on DEAE-Sephadex A-50 chromatography	
J.-G. Geng, H.-Z. Chen, Y.-F. Yang, C.Y. Jian, Z.-H. Qian, Q.-Q. Han, Y.-M. Yao	233
Choline supplementation increases serum alkaline phosphatase activity in rats	
J. Wuarin-Bierman, L. Wecker	237

Vol. 176/3, 1988

The prolamin antibody reactivity against hordein polypeptides in sera from patients with coeliac disease	
S.U. Friis, H. Sjöström, O. Norén, N. Rüdiger, D. Anthonsen	241
Sialic acid as a marker for differentiation between benign and malignant intracranial tumors	
E. Marth, G. Flaschka, S. Stiegler, J.R. Möse	251
Synthesis of pyrenesulfonylamido-sphingomyelin and its use as substrate for determining sphingomyelinase activity and diagnosing Niemann-Pick disease	
R. Klar, T. Levade, S. Gatt	259
Ion-exchange chromatography and clinical criteria in the screening of the aminoacidopathies	
P.R. Parvy, J.I. Bardet, D.M. Rabier, J.-M. Saudubray, P.P. Kamoun	269

Amino acid composition of peptides remaining after <i>in vitro</i> digestion of a gliadin sub-fraction with duodenal mucosa from patients with coeliac disease	
H.J. Cornell	279
Solid phase immunoassay for high molecular weight alkaline phosphatase in human sera using a specific monoclonal antibody	
J.T. Deng, P.G. Parsons	291
Determination of human erythropoietin by radioimmunoassay. Method and clinical data	
O.J. Nielsen	303
Field measurement of plasma and erythrocyte cholinesterases	
R.A. Magnotti, Jr., K. Dowling, J.P. Eberly, R.S. McConnell	315
Enhanced urinary trypsin inhibitory activity in gentamicin-induced nephrotoxicity in rats	
S. Smetana, S. Khalef, Z. Nitsan, N. Hurwitz, A. Miskin, Y. Bar-Khayim, Y. Birk	333
Measurement of cytochrome <i>c</i> oxidase activity in human liver specimens obtained by needle biopsy	
Y. Sakai, A. Tanaka, I. Ikai, Y. Yamaoka, K. Ozawa, Y. Orii	343

Vol. 177/1,2, 1988

Assessment of the role of pancreatic proteases in human abdominal aortic aneurysms and occlusive disease	
M.A. Dubick, G.C. Hunter, E. Perez-Lizano, G. Mar, M.C. Geokas	1
Effect of colchicine on collagen synthesis by liver fibroblasts in murine schistosomiasis	
M.M. Mansour, M.A. Dunn, L.A. Salah	11
Strong activity of acid-stable trypsin inhibitor in bovine thrombin for clinical use	
H. Sumi, H. Hamada, S. Maehara, H. Mihara	21
Enzyme-linked immunosorbent assay of lipoprotein(a) in serum and cord blood	
A. Abe, S. Maeda, K. Makino, M. Seishima, K. Shimokawa, A. Noma, M. Kawade	31
Elevated nucleotide pyrophosphatase activity in cultured skin fibroblasts from patients with Lowe's syndrome	
K. Horie, T. Yano, I. Funakoshi, I. Yamashima	41
Clinical relevance of increase neuron-specific enolase concentration in cerebrospinal fluid	
C. Jacobi, H. Reiber	49
Stabilization and quantitative determination of disulfiram in human plasma samples	
B. Johansson	55
Purification of acidic glutathione S-transferase from human lung, placenta and erythrocyte and the development of a specific radioimmunoassay for their measurement	
A.F. Howie, J.D. Hayes, G.J. Beckett	65
The diagnostic value of urinary transferrin compared to serum prostatic specific antigen (PSA) and prostatic acid phosphatase (PAP) in patients with prostatic cancer	
M.P. van Diejen-Visser, M.W.G. Hendriks, K.P.J. Dekae, A.H.J. Gijzen, P.J. Brombacher	77
Iron, copper and zinc concentrations in human sweat and plasma: the effect of exercise	
O.J. Aruoma, T. Reilly, D. MacLaren, B. Halliwell	81
A method to detect the <i>in vitro</i> modification of serum beta-2-microglobulin in health and disease	
M.A. Forbes, A.M. Cox, E.H. Cooper	89
Relationship between the urinary excretion of albumin and retinol-binding protein in insulin-dependent diabetics	
J. Holm, L. Hemmungen, N.V. Nielsen	101
Micromethod for determination of ATPase activities using bioluminescence technique	
M. Steiner, H. Bauer, F. Krassnigg, W.-B. Schill, H. Adam	107
Letters to the Editors	
Second antibodies covalently linked to Sepharose with 1,1'-carbonyldimidazole — effect of storage on radioimmunoassay separation efficiency	
M.G. McConway, R.S. Chapman	115
Platelet intracellular calcium in patients with Duchenne muscular dystrophy	
C. Lechi, D. Sinigaglia, P. Guzzo, G. Bonadonna, M. Zatti	119

Vol. 177/3, 1988

A simplified method for determination of uroporphyrinogen decarboxylase activity in human blood	
Adjarov D.G., Elder G.H.	123
Levels of 5'-nucleotide phosphodiesterase isoenzymes in normal and pathological sera	
Luthje J., Pickert S., Ogilvie A., Hornemann E., Siegfried W., Waldherr A., Domschke W.	131
Serum procollagen III N-terminal peptide and laminin P1 fragment concentrations in alcoholic liver disease and primary biliary cirrhosis	
Van Zanten R.A.A., Van Leeuwen R.E., Wilson J.H.P.	141
Plasma alkaline phosphatase isoenzymes in diabetes mellitus	
Tibi L., Collier A., Patrick A.W., Clarke B.F., Smith A.F.	147
Correlation between concentrations of magnesium, zinc, and potassium in plasma, erythrocytes and muscles	
Ladefoged K., Hagen K.	157
Diagnostic efficiency in discriminating liver malignancies from cirrhosis by serum gamma-glutamyltransferase isoforms	
Sacchetti L., Castaldo G., Cimino L., Budillon G., Salvatore F.	167
Fat macroglobule formation from chylomicrons and non-traumatic fat embolism	
Hulman G.	173
Dipeptidyl peptidase III and alanyl aminopeptidase in the human seminal plasma: origin and biochemical properties	
Vanha-Perttula T.	179
A specific method for the direct determination of lipoprotein cholesterol in electrophoretic patterns	
Aufenanger J., Haux P., Weber U., Kattermann R.	197

Vol. 177/3, 1988

Urinary neopterin excretion in pulmonary sarcoidosis: Correlation to clinical course of the disease	
Prior C., Frank A., Fuchs D., Hausen A., Judmaier G., Reibnegger G., Werner E.R., Wachter H.	211
Precipitation of LDL with sulfated polyanions: Three methods compared	
Siekmeier R., Marz W., Gross W.	221
Hb G-Philadelphia, or [$\alpha 68(E17)Asn \rightarrow Lys$], in North Sardinia: Detection by isoelectric focusing and identification by HPLC of tryptic peptides	
Manca L., Demuro P., Masala B.	231
Discovery of methoxyacetylcarbamide in the urine of normal adults and phenylketonuric children	
Antoshechkin A.G., Tatur V.Y., Maximova L.A.	239
Accurate and specific HPLC assay of coproporphyrinogen III oxidase activity in human peripheral leucocytes	
Guo R., Lim C.K., Peters T.J.	245
Quantitative determination of biocytin in urine of patients with biotinidase deficiency using high-performance liquid chromatography (HPLC)	
Suormala T.M., Baumgartner E.R., Bausch J., Holick W., Wick H.	253
The effect of age and sex on β -hexosaminidase in urine	
Hultberg B., Isaksson A., Berg B., Tryding N., Ekman S., Nilsson J.-E.	271
Approved recommendation (1988) on the theory of reference values. Part 3. Preparation of individuals and collection of specimens for the production of reference values	
Solberg H.E.	S3

Vol. 178/1, 1988

Apolipoprotein measurements in clinical biochemistry and their utility vis-a-vis conventional assays	
Bachorik P.S., Kwiterovich Jr. P.O.	I

Colorimetric method for assay of serum γ -glutamyltransferase activity with some L- γ -glutamyl-carboxyanilides Szewczuk A., Kuropatwa M., Lang D.	35
Serum noncholesterol sterols related to cholesterol metabolism in familial hypercholesterolemia Gylling H., Miettinen T.A.	41
Sodium-lithium countertransport activity and its sensitivity to inhibitors with erythrocyte ageing in man Carr S., Thomas T.H., Wilkinson R.	51
Determination of 3,4-dihydroxyphenylalanine (DOPA) in plasma and cerebrospinal fluid by high performance liquid chromatography with electrochemical detection Boomsma F., Van der Hoorn F.A.J., Man in 't Veld A.J., Schalekamp M.A.D.H.	59
A sensitive non-radioactive assay for pyrimidine nucleoside phosphorylase using reversed-phase high performance liquid chromatography Laurense E.J., Pinedo H.M., Peters G.J.	71
Interaction of horse-spleen ferritin with aluminium citrate Cochran M., Chawtur V.	79
Plasma acetate levels in response to intravenous fat or glucose/insulin infusions in diabetic and non-diabetic subjects Akanji A.O., Ng L., Humphreys S.	85
Thermostability of human serum biotinidase activity Hayakawa K., Ozumi J.	95
Allelic polymorphism of human placental alkaline phosphatase in three races: sugar chain heterogeneities Sato M., Furiya K., Tanaka S., Sakagishi Y., Komoda T.	101

Vol. 178/2, 1988

Correlation of two different assays for urinary kallikrein in normotensive and hypertensive subjects Maier M., Jerabek I., Reissert G., Holtzl E., Binder B.R.	127
An immunoblotluminescence assay for gamma-gamma enolase activity in human serum and cerebrospinal fluid Wevers R.A., Theunisse A.W.G., Rijksen G.	141
Serum pseudouridine as a biochemical marker in patients with hepatocellular carcinoma Amuro Y., Nakaoka H., Shimomura S., Fujikura M., Yamamoto T., Tamura S., Hada T., Higashino K.	151
Competitive enzyme-linked immunosorbent assay (ELISA) for the quantitation of apolipoprotein A-I using a monoclonal antibody Bojanovski M., Gregg R.E., Wilson D.M., Brewer Jr. H.B.	159
A 3,3',5-triiodothyronine autoantibody (IgG, lambda) in a case of Hashimoto's thyroiditis Muratsugu M., Tsuchiya Y., Makino M.	171
Cholesterol monomer activity correlates with nucleation time in model bile Chijiwa K., Kiyosawa R., Nakayama F.	181
Enzyme immunoassay of medullasin in peripheral blood Aoki Y., Yoshida M., Kominami E.	193
A sensitive two-site immunoassay for human pancreatic secretory trypsin inhibitor (PSTI) using monoclonal antibodies Kurobe M., Kono M., Yoshida N., Hayashi K.	205
Comparison of the performance and clinical utility of a carboxy-terminal assay and an intact assay for parathyroid hormone St. John A., Davies C., Riley W.J., Kent G.N., Brown R.C., Aston J.P., Weeks I., Woodhead J.S.	215
Serum osteocalcin and other indices of bone formation: An 8-decade population study in healthy men and women Worsfold M., Sharp C.A., Davie M.W.J.	225

Vol. 178/3, 1988

Betaine metabolism in human neonates and developing rats Davies S.E.C., Chalmers R.A., Randall E.W., Iles R.A.	241
Triglyceride lowering effect of MaxEPA fish lipid concentrate: a multicentre placebo controlled double blind study Miller J.P., Heath I.D., Choraria S.K., Shephard N.W., Gajendragadkar R.V., Marcus A.W., Batson G.A., Smith D.W., Saynor R.	251
Enzyme-linked immunosorbent assay for quantitation of cereal proteins toxic in coeliac disease Friis S.U.	261
Increase of erythrocyte resistance to hemolysis and modification of membrane lipids induced by hemodialysis Peuchant E., Salles C., Vallot C., Wone C., Jensen R.	271
Level of plasma fibronectin in patients with breast cancer Ruelland A., Kerbrat P., Clerc C., Legras B., Cloarec L.	283
Characterisation of galactosyltransferase isoforms by ion-exchange and lectin affinity chromatography Boyle F.A., Peters T.J.	289
Carbamylated haemoglobin in chronic renal failure Smith W.G.J., Holden M., Benton M., Brown C.B.	297
A second antibody solid-phase enzyme immunoassay of prostaglandin E ₂ in human gastric juice Orsini B., Calabro A., Renzi D., Pradelles P., Fedi P., Surrenti C.	305
Is the ex vivo rat gastric chamber model suitable for studying the gastrototoxicity of refluxed duodenal contents? Initial results using deoxycholic acid Armstrong D., Farrell M., Hanby A., Murphy G.M., Dowling R.H.	313
Detection of lysozyme and alpha ₂ -macroglobulin-lysozyme complexes by immunoblotting Moraky P.	327
Alterations in specific activity of lysosomal alpha-glucosidase in cystic fibrosis Vladutiu G.D., Kewin C.C.	337
Enzymatic antioxidative defence of erythrocytes in an Italian family with Hb Volga or $\alpha_2\beta_2$ 27 (B9) Ala \rightarrow Asp Falcioni G., Grelloni F., De Sanctis G., Pierani P., Felici L., Coppa G.V.	345
Purine metabolism in a bone-marrow transplanted adenosine deaminase deficient patient Boulieu R., Bory C., Souillet G.	349

Vol. 179/1, 1989

Evaluation of a direct assay for atrial natriuretic peptide Tan A.C.I.T.L., Rosmalen F.M.A., Hofman J.A., Kloppenborg P.W.C., Benraad Th. J.	1
Specific determination of plasma dexamethasone by HPLC and RIA - application to standard dexamethasone suppression test in psychiatric patients Lejeune-Lennin C., Bosson D., Linkowski P.	13
Clinical significance of a new isoform of serum alanine aminopeptidase: relationship with liver disease and alcohol consumption Sanderink G.-J., Artur Y., Paille F., Siest G.	23
Association between increased concentrations of free thyroxine and unsaturated free fatty acids in non-thyroidal illnesses: role of albumin Tikanoja S.H., Joutti A., Liewendahl B.K.	33
Complement activation after prolonged exercise Dufaux B., Order U.	45
Urinary excretion of 2,3-butanediol and acetoin by babies on a special care unit Mills G.A., Walker V.	51
Isolation and characterization of dipeptidyl aminopeptidase IV from human kidney cortex Wolf G.B., Scherberich J.E., Fischer P., Schoeppe W.	61

Stability of alpha ₂ -microglobulin, beta ₂ -microglobulin and retinol binding protein in urine Donaldson M.D.C., Chambers R.E., Woolridge M.W., Whicher J.T.	73
Validation of the Friedewald formula for the measurement of low density lipoprotein cholesterol in insulin-dependent diabetes mellitus Winocour P.H., Ishola M., Durrington P.N.	79
Characterization of two monoclonal antibodies to human apolipoprotein A-I that also bind to rabbit apolipoprotein A-I. Application to ELISA evaluation of rabbit serum apo A-I Leroy A., Olivier P., Vu-Dac N., Fruchart J.C.	85
The determination of plasma oxalate concentrations using an enzyme/bioluminescent assay. 2. Co-immobilisation of bioluminescent enzymes and studies of in vitro oxalogenesis Parkinson I.S., Channon S.M., Tomson C.R.V., Adonai L.R., Ward M.K., Laker M.F.	97
Variation in alpha-1-antitrypsin phenotypes associated with penicillamine therapy Whitehouse D.B., Lovegrove J.U., Hopkinson D.A.	109

Vol. 179/2, 1989

Effect of fish oil supplementation on erythrocyte lipid pattern, malondialdehyde production and glutathione-peroxidase activity in psoriasis Corrocher R., Ferrari S., De Gironcoli M., Bassi A., Olivieri O., Guarini P., Stanzial A., Barba A.L., Gregolini L.	121
Human plasma contains low molecular weight factors which stimulate active sodium transport in erythrocytes Tepper T., Huisman R.M., Jilderda J.F., Van der Hem G.K., De Zeeuw D.	133
α -Fetoprotein antibody-lectin enzyme immunoassay to characterize sugar chains for the study of liver diseases Kinoshita N., Suzuki S., Matsuda Y., Taniguchi N.	143
Severity of GM1 gangliosidosis and urinary oligosaccharide excretion Takahashi Y., Orii T.	153
Racial effect on serum creatine-kinase: Implications for estimation of heterozygosity risks for females at-risk for Duchenne dystrophy Passos-Bueno M.R., Rabbi-Bortolini E., Azevedo E., Zatz M.	163
Sandwich enzyme immunoassay for serum amyloid A protein (SAA) Yamada T., Uchiyama K., Yakata M., Gejyo F.	169
A sensitive and specific chemiluminescence method for estimating the ability of human granulocytes and monocytes to generate O ₂ ⁻ Nishida A., Kimura H., Nakano M., Goto T.	177
Isolation of apolipoprotein A-I-containing lipoproteins by immunoaffinity chromatography using a formylated derivative of cellulose gel Ohta T., Hattori S., Murakami M., Horiuchi S., Nishiyama S., Matsuda I.	183
Determination of tellurium in biological fluids by means of electrothermal vaporization-inductively coupled plasma-mass spectrometry (ETV-ICP-MS) Newman R.A., Osborn S., Siddik Z.H.	191
Insulin binding to erythrocytes and fatty acid composition of erythrocyte membrane phospholipids in healthy men Pelikanova T., Kohout M., Hilgertova J., Rase J., Farska I., Valek J.	197
Carboxyl terminal glycine extended progastrin (gastrin-G) in human gastric mucosa and gastrinomas Azuma T., Inokuchi H., Kawai K., Fujimoto S., Nakajima M., Taggart R.T., Walsh J.H.	201
Role of variant prealbumin in the pathogenesis of familial amyloidotic polyneuropathy: Fate of normal and variant prealbumin in the circulation of rats Ando Y., Ikegawa S., Miyazaki A., Inoue M., Morino Y., Araki S.	205
Serum and ascitic fluid concentrations of apolipoproteins A and B in alcoholic cirrhosis patients Nayak S.S., Bhat K.R., Nair A., Rao S., Arora A.R.	207

Vol. 179/3, 1989

Diagnosis of subtypes of GM1 gangliosidosis in vitro and in vivo — Using urinary oligosaccharides as substrates Takahashi Y., Orii T.	219
Improved competitive enzyme-linked immunoassay (ELISA) for albuminuria Neumann R.G., Cohen M.P.	229
Fibrinogen sevilla, a congenital dysfibrinogenemia characterized by an abnormal monomer aggregation and a defective plasmin lysis Fernandez F.J., Noguerol P., Sosa R., Cuesta B., Paramo J.A., Rocha E.	239
Assessment of antithyroglobulin and antimicrosomal autoantibodies in patients with autoimmune thyroid disease: Comparison of haemagglutination assay, enzyme-linked immunoassay and radioligand assay Engler H., Staub J.-J., Althaus B., Ryff-deLeche A., Gerber H.	251
Application of solid-phase antibodies to radioimmunoassay. Improved performance of Dynospheres® following affinity purification of antisera McConway M.G., Chapman R.S.	265
Relationship between hair, serum and bone aluminium in hemodialyzed patients Chappuis P., De Vernejoul M.-C., Paolaggi F., Rousselet F.	271
The subunit composition of factor XIII proteins in normal and factor XIII deficient plasma and serum analysed by line immunoelectrophoresis Kroll J.	279
Changes in the immunoreactivity of Apo A-I during storage Wang X.L., Dudman N.P.B., Blades B.L., Wilcken D.E.L.	285
Erythrocyte and plasma cholesterol exchange in sickle cell anemia Ngogang J., Mouray H., Lebreton de Vonne T., Raisonier A.	295
Microbial influences on urinary polyamine excretion Satink H.P.W.M., Hessels J., Kingma A.W., Van Den Berg G.A., Muskiet F.A.J., Halie M.R.	305
Procollagen III peptides and fibronectin in alcohol-related chronic liver disease: Correlations with morphological features and biochemical tests Gabrielli G.B., Faccioli G., Casaril M., Capra F., Bonazzi L., Falezza G., Tomba A., Barcchino F., Corrocher R.	315
Kinetics of serum glutamate dehydrogenase during short-term alcohol withdrawal Weill J., Schellenberg F., Le Goff A.-M.	323
The lysosomal β -D-N-acetylglucosaminidase isozymes in human plasma during pregnancy: Separation and quantification by a simple automated procedure Goi G., Fabi A., Lombardo A., Bairati C., Bovati L., Burlina A.B., Agosti S., Serio C., Tettamanti G.	327
Retention of antimony in hair during leishmaniasis treatment Dorea J.G., Merchan-Hamann E., Ryan D.E., Holzbecher J.	341

Vol. 180/1, 1989

Monoclonal gammopathies — Their identification and biological significance Osterland C.K.	1
Inherited occurrence of a heat stable alkaline phosphatase in the absence of malignant disease Onica D., Rosendahl K., Waldenlind L.	23
Lipid lowering actions of 7-(2-methylene butyryl)-(2H)-1,4-benzoxazin-3-(4H)-one derivatives in mice, rats and Syrian hamsters Moussavi Z., Plancke M.O., Olivier Ph., Lesieur D., Fruchart J.C., Sauzieres J.	35
Clinical significance of mucin-like high molecular weight glycoprotein originated from lung cancer as tumor marker Imokawa M., Okabe H., Ochi Y., Kajita Y.	45
Identification of an extra lactate dehydrogenase band in human erythrocytes Hirano T., Matsuzaki H., Sekine T.	59

Characteristics and partial purification of prolidase and prolinase from leukocytes of a normal human and a patient with prolidase deficiency	63
Kodama H., Ohhashi T., Ohba Ch., Ohno T., Arata J., Kubonishi I., Miyoshi I.	
Development of an RIA-based primary screen for IL-1 antagonists	73
Maloff B.L., Shaw J.E., Di Meo T.M., Fox D., Bruin E.M.	
Determination of the biological activity of antithrombin III related antigen in urine	79
Ueshima S., Yamamoto T., Okada K., Tanaka H., Matsuo O.	
Comparison of unitages for total free erythrocyte protoporphyrin (total EP) in anemia screening	87
Harada K., Omori S., Ueno T., Misumi J., Miura H.	
Direct measurement of whole blood glucose by a needle-type sensor	93
Yamasaki Y., Ueda N., Nao K., Sekiya M., Kawamori R., Shichiri M., Kamada T.	
Aspartylglucosamine excretion in heterozygous carriers of aspartylglucosaminuria	99
Mononen T.K.	
Comparative study: The antioxidant enzymes and lipid peroxidation in cord and maternal red blood cell haemolysates	103
Novak Z., Kovacs L., Pal A., Pataki L., Varga Sz. I., Matkovic B.	

Vol. 180/2, 1989

Plasma lipoproteins and adrenocortical hormones in men — Positive association of low density lipoprotein cholesterol with plasma cortisol concentration	113
Nanjee M.N., Miller N.E.	
Glycated hemoglobin in hypoglycemia	121
Niketic V., Djurdjic V., Stojkovic V., Maric S., Djordjevic P.	
Single step affinity chromatographic purification of human α -amylase from aspirated duodenal juice and its application in the measurement of pancreatic α -amylase synthesis rates in man	129
Ogden J.M., O'Keefe S.J.D., Ehlers M.R.W., Kirsch R.E., Marks I.N.	
Amniotic fluid phospholipids. Study of the cold acetone precipitation effect	141
Budria R.J., Gonzalez F.E., Sanz H.M.T., Ostari S.J.L.	
Variability in lipoprotein concentrations in serum after prolonged storage at -20°C	147
Tiedink H.G.M., Katan M.B.	
Measurement of 25-hydroxyvitamin D ₂ , 25-hydroxyvitamin D ₃ , 24,25-dihydroxyvitamin D ₂ and 25,26-dihydroxyvitamin D ₂ in a single plasma sample by mass fragmentography	157
Coldwell R.D., Trafford D.J.H., Varley M.J., Kirk D.N., Makin H.L.J.	
Metabolism of pyrazinamide and allopurinol in hereditary xanthine oxidase deficiency	169
Yamamoto T., Higashino K., Kono N., Kawachi M., Nanahoshi M., Takahashi S., Suda M., Hada T.	
Differences between human liver- and bone-type alkaline phosphatases	177
Miura M., Matsuzaki H., Bailyes E.M., Koyama I., Sakagishi Y., Sekine T., Komoda T.	

Vol. 180/3, 1989

The simultaneous determination of oxidized and reduced glutathiones in liver tissue by ion pairing reverse phase high performance liquid chromatography with a coulometric electrochemical detector	203
Harvey P.R.C., Ilson R.G., Strasberg S.M.	
Binding of glycosaminoglycan inhibitors to calcium oxalate crystals in relation to ionic strength	213
Fellstrom B., Lindsjo M., Danielson B.G., Karlsson F.A., Ljunghall S.	
Renin substrate in human amniotic fluid	221
Metsarinne K., Fyhrquist F., Rosenlof K., Autero M., Pekonen F., Rutanen E.-M.	
Effect of drugs on histamine radio-enzyme assay	231
Harvima R.J., Harvima I.T., Kajander E.O., Penttila I.M., Horsmanheimo M., Fraki J.E.	
The effect of folic acid and 5-formyltetrahydrofolic acid on folate-deficient HEP 2 cells in culture	241
Leeming R.J., Hamon C.G.B., Tomkyns J.G.	

Circadian and seasonal variations of electrolytes in aging humans	
Toutiou Y., Toutiou C., Bogdan A., Reinberg A., Motohashi Y., Auzeby A., Beck H.	245
Correction of subclinical ascorbate deficiency in patients receiving dialysis: Effects on plasma oxalate, serum cholesterol, and capillary fragility	
Tomson C.R.V., Channon S.M., Parkinson I.S., McArdle P., Qureshi M., Ward M.K., Laker M.F.	255
Interactions with hemoglobin: A source of error in measurements of transketolase activity in hemolysates	
Grudzinski A., Waltham M., Ioannoni B., Price J., Nixon P.F.	265
A multiwell lectin-binding assay using <i>Lotus tetragonolobus</i> for measuring different glycosylated forms of haptoglobin	
Thompson S., Stappenbeck R., Turner G.A.	277
A two-site immunoradiometric assay for C-reactive protein in serum	
Shapiro D., Shenkin A.	285
Detection and identification of the plant lignans lariciresinol, isolariciresinol and seco-isolariciresinol in human urine	
Bannwart C., Adlercreutz H., Wahala K., Brunow G., Hase T.	293
α -L-Fucosidase activity in normal human lymphocytes	
Roger L., Bernard M.-A., Percheron F., Foglietti M.-J.	303

Vol. 181/1, 1989

Isolation of creatine kinase BB isoenzyme with high specific activity and adequate purity for radioimmunoassay from human placenta on preparative polyacrylamide gel electrophoresis	
Geng J.-G., Cheng H.-Z., Yang Y.-F., Qian Z.-H., Jiang C.-Y.	1
Alterations of acidic phospholipids in bronchoalveolar lavage fluids of patients with pulmonary alveolar proteinosis	
Honda Y., Kataoka K., Hayashi H., Takahashi H., Suzuki A., Akino T.	11
An enzyme immunoassay of slow moving protease (SMP) in human gastric mucosa	
Matsusako K., Itoh M., Joh T., Miyamoto T., Yokochi K., Takeuchi T., Moriyama A., Kato T., Tanaka R.	19
Sensitive enzyme immunoassay for human aldolase B	
Haimoto H., Kurobe N., Hosoda S., Kato K.	27
Mechanism of inhibition of erythrocyte glutathione reductase by Mitomycin-C	
Okuda J., Hirai Y., Hayazaki T.	37
Glycan microheterogeneity of α_1 -acid glycoprotein in sera and dialysates of patients on continuous ambulatory peritoneal dialysis	
Fraeyman N.H., Van de Velde E.J., Belpaire F.M., Lameire N.H.	47
Increased phosphoinositide breakdown by phospholipase C in erythrocyte membranes from patients with cystic fibrosis	
Doughney C., McPherson M.A., Goodchild M.C., Dormer R.L.	55
Aberrant free radical activity in cystic fibrosis	
Salh B., Webb K., Guyan P.M., Day J.P., Wickens D., Griffin J., Braganza J.M., Dormandy T.L.	65
Uptake of trihalomethanes by patients during hemodialysis	
Cailleux A., Subra J.F., Riberi P., Allain P.	75
The serum activities of AP, gamma-GT, GLDH, GPT and CHE after complete biliary obstruction and choledochocaval fistula in the rat	
Putzki H., Reichert B., Heymann H.	81
Serum unconjugated bile acids in patients with small bowel bacterial overgrowth	
Bolt M.J.G., Stellaard F., Sitrin M.D., Paumgartner G.	87
Metabolism of androstenedione by placental microsomes in pregnancy hypertension	
Hahnel M.E., Martin J.D., Michael C.A., Hahnel R.	103
Reversed-phase high-performance liquid-chromatography of 2,8-dihydroxyadenine in serum and urine with electrochemical detection	
Kojima T., Nishina T., Kitamura M., Kamatani N., Nishioka K.	109

Vol. 181/2, 1989

A new case with hereditary xanthinuria: Response to exercise Landaas S., Borch K., Aagaard E.	119
Biochemical diagnosis of Hunter syndrome on Epstein-Barr virus-transformed lymphoblastoid cell lines Morabito E., Giambarrasi I., Rocchi M., Di Natale P.	125
Comparison between the chromate inhibition test and a cytochemical method for the determination of glucose-6-phosphate dehydrogenase deficiency in erythrocytes Jonges G.N., Hagen H., Van Noorden C.J.F., Weening R.S., Roos D.	135
Measurement of plasma uroporphyrin during treatment and long-term follow-up of patients with Porphyria cutanea tarda Adjarov D.G., Kerimova M.A.	143
Concentration and physicochemical characterisation of unsaturated cobalamin binding proteins in amniotic fluid Gueant J.L., Hambaba L., Vidailhet M., Schaefer C., Wahlstedt V., Nicolas J.P.	151
Patients with hereditary fructose intolerance have normal erythrocyte aldolase activity Gitzelmann R., Steinmann B., Tuchschild	163
Phosphatidylcholine storage in mucopolipidosis IV Bargal R., Bach G.	167
Carnitine deficiency associated with anticonvulsant therapy Rodriguez-Segade S., Alonso de la Pena C., Tutor J.C., Paz J.M., Fernandez M.P., Rozas I., Del Rio R.	175
Plasma and serum lactoferrin levels in cystic fibrosis. Relationship with the presence of cystic fibrosis protein Barthe C., Galabert C., Guy-Crotte O., Figarella C.	183
Determination of NADH ₂ -ferricyanide oxidoreductase (cytochrome b ₅ reductase, diaphorase) activity of human erythrocytes by an analysis of the time dependence of NADH ₂ oxidation Papandreou P., Rakitzis E.T.	189
Correlations among blood levels of renin-angiotensin-aldosterone, bradykinin, and catecholamines in healthy subjects Yamamoto T., Hiramoto J., Kikuchi Y., Oka M., Takatori O.	197
Two new serum immunoassay (RIA and EIA) for the determination of M ₂ antimitochondrial antibodies in primary biliary cirrhosis Plebani M., Chiaramonte M., Floreani A., Bertorelle R., Naccarato R., Burlina A.	201
An improved spectrophotometric bromide assay for the estimation of extracellular water volume Trapp S.A., Bell E.F.	207
Simultaneous assay of zinc protoporphyrin and protoporphyrin IX in dried blood Orfanos A.P., Guthrie R., Jinks D.C., Vollmer D.W.	213
A micromethod for the determination of fructose in human erythrocytes and plasma Liu S.-Q., Das B., Srivastava S.K.	219
Nonspecific reaction in the sandwich immunoassay for human tumor necrosis factor- α (hTNF- α) Abe Y., Miyake M., Miyazaki T., Horiuchi A., Sagawa T., Ono H., Kimura S.	223
Analysis of acylcarnitines in normal human urine with the radioisotope exchange-high performance liquid chromatography (HPLC) method Schmidt-Sommerfeld E., Penn D., Kerner J., Bieber L.L.	231

Vol. 181/3, 1989

Proteinuria in sickle cell trait and disease: An electrophoretic analysis Lonsdorfer A., Comoe L., Yapoe A.E., Lonsdorfer J.	239
Tetrahydrobiopterin metabolism in the streptozotocin induced diabetic state in rats Hamon C.G., Cutler P., Blair J.A.	249
Pharmacokinetic study of butyric acid administered in vivo as sodium and arginine butyrate salts Daniel P., Brazier M., Cerutti I., Pieri F., Tardivel I., Desmet G., Baillet J., Chany C.	255

An improved formula for the judgement of intrathecally produced IgG in the presence of blood brain barrier damage	
Ohman S., Forsberg P., Nelson N., Vrethem M.	265
Hydroxycholesterols in serum from hypercholesterolaemic patients with and without bile acid sequestrant therapy	
Van Doormaal J.J., Smit N., Koopman B.J., Van der Molen J.C., Wolthers B.G., Doorenbos H.	273
Serum human chorionic gonadotrophin levels in early pregnancy	
Ooi D.S., Perkins S.L., Claman P., Muggah H.F.	281
One step sandwich enzyme immunoassay for human type IV collagen using monoclonal antibodies	
Obata -i. K., Iwata K., Ichida T., Inoue K., Matsumoto E., Muragaki Y., Ooshima A.	293
Detection of elevated amounts of urinary pseudouridine in cancer patients by use of a monoclonal antibody	
Itoh K., Mizugaki M., Ishida N.	305
Assay of serum hyaluronic acid in clinical application	
Chichibu K., Matsuura T., Schichijo S., Yokoyama M.M.	317
The quantitation of human ventricular myosin light chain 1 in serum after myocardial necrosis and infarction	
Wang J., Shi Q., Wu T.-W., Jackowski G., Mickle D.A.G.	325
A new approach for measuring the erythrocyte life span with a nonradioisotope	
Dever M., Smith J.E., Hausler D.W.	337

Vol. 182/1, 1989

Radioisotopic assay for erythrocyte adenosine 5'-monophosphate deaminase	
Dale G.L.	1
Carbohydrate removal fails to eliminate the heterogeneity of human prostatic acid phosphatase	
Morris M.F., Waheed A., Risley J.M., Van Etten R.L.	9
Immunoglobulins from the sera of immunologically activated persons with pairs of electrophoretic restricted bands show a greater tendency to aggregate than normal	
Levinson S.S., Keren D.F.	21
Inhibition of neutrophil and natural killer cell function by human seminal fluid acid phosphatase	
Mukhopadhyay N.K., Saha A.K., Smith W., Dowling J.N., Hiserodt J., Glew R.H.	31
Effects of N-hydroxyethyl-1-deoxynojirimycin (BAY m 1099) on the activity of neutral- and acid α -glucosidases in human fibroblasts and HepG2 cells	
Wisselaar H.A., Van Dongen J.M., Reuser A.J.J.	41
Simultaneous determination of hemoglobin and coproporphyrin by second derivative differential spectrophotometry: Application to the diagnosis of meconium aspiration	
Ficheux H., Levillain P., Francoual J., Leluc R., Lemonnier A.	53
Adenylate degradation products release from the human myocardium during open heart surgery	
Smolenski R.T., Skladanowski A.C., Perko M., Zydowo M.M.	63
Binding of fibronectin to human lipoproteins	
Cseh K., Karadi I., Riachak K., Szollar L., Janoki G., Jakab L., Romics L.	75
A spectrophotometric method for the determination of free and esterified carnitine	
Schafer J., Reichmann H.	87
Simple enzymatic screening assay for ethylene glycol (ethane-1,2-diol) in serum	
Hansson P., Masson P.	95
Quantitative estimation of an α_2 pregnancy-associated protein in normal pregnancy by electro-immunoassay	
Nakai M., Atsumi M., Sone Y., Arai M.	103
Elimination of free glycerol interference in a colorimetric enzymic triglyceride assay	
Artiss J.D., Strandbergh D.R., Zak B.	109
New enzymatic endpoint assay of serum sialic acid	
Horiuchi T., Kurokawa T.	117

Vol. 182/2, 1989

The effect of hypoalbuminaemia, hyperbilirubinaemia and renal failure on serum fructosamine concentration in non-diabetic individuals Dominiczak M.H., Orrell J.M., Finlay W.E.I.	123
Measurement of glomerular filtration rate in children with insulin-dependent diabetes mellitus Gibb D.M., Dalton N.R., Barratt M.T.	131
Measurement of propionate turnover in vivo using sodium [$^2\text{H}_5$]propionate and sodium [^{13}C]propionate Walter J.H., Thompson G.N., Leonard J.V., Heatherington C.S., Bartlett K.	141
Detection of point mutations in amplified single copy genes by biotin-labelled oligonucleotides: Diagnosis of variants of alpha-1-antitrypsin Gegersen N., Winter V., Petersen K.B., Koch J., Kolvraa S., Rudiger N., Heinsvig E.-M., Bolund L.	151
Red cell membrane phosphatidylinositol kinase activity in hemolytic anemias and myeloproliferative diseases Boivin P., Galand C.	165
Evidence that free gamma carboxyglutamic acid circulates in serum Fournier B., Gineyts E., Delmas P.D.	173
Serum CK-BB as a tumor marker in patients with carcinoma confirmed histologically Arenas J., Diaz A.E., Alcaide M.J., Santos I., Martinez A., Culebras J.M.	183
Plasma protein S in disseminated intravascular coagulation, liver disease, collagen disease, diabetes mellitus, and under oral anticoagulant therapy Takahashi H., Takewaki W., Wada K., Shibata A.	195
Human serum immuno-reactive copper, zinc-superoxide dismutase assayed with an enzyme monoclonal immunosorbent in patients with digestive cancer Oka S., Ogino K., Matsuura S., Yoshimura S., Yamamoto K., Okazaki Y., Takemoto T., Kato N., Uda T.	209
Serum selenium in liver cirrhosis: Correlation with markers of fibrosis Casaril M., Stanzial A.M., Gabrielli G.B., Capra F., Zenari L., Galassini S., Moschini G., Liu N.Q., Corrocher R.	221
Measurement of free kappa immunoglobulin light chains in the cerebrospinal fluid by a competitive avidin-biotin ELISA Lolli F., Amaducci L.	229

Vol. 182/3, 1989

The effects of Oradexone and Ambroxol pretreatment on the oxidative sensitivity of the red blood cells in preterm infants Novak Z., Varga S.I., Kovacs L., Pal A., Pataki L., Matkovics B.	241
A simple, rapid assay for plasma oxalate in uraemic patients using oxalate oxidase, which is free from vitamin C interference Rolton H.A., McConnell K.N., Modi K.S., Macdougall A.I.	247
Binding of calcium by organic anions, determined by perturbation of the equilibrium solubility of [^{14}C]calcium oxalate Geller D.A., Ostrow J.D., Moore E.W., Celic L., Nancollas G.H.	255
Reference values for amino acids in cerebrospinal fluid of children determined using ion-exchange chromatography with fluorimetric detection Gerrits G.P., Trijbels F.J., Monnens L.A., Gabreels F.J., De Abreu R.A., Theeuwes A.G., Raay-Selten B.	271
N-acetyl- β -glucosaminidase and albuminuria in normal and diabetic pregnancies Skrha J., Perusicova J., Sperl M., Bendl J., Stolba P.	281
Identification of octadeca-9,11-dienoic acid and its significance in the diagnosis of cervical intra-epithelial neoplasia Parham D.M., Ross P.E., Gunstone F.D., Robertson A.J., Duncan I.D.	289

Application of two monoclonal antibodies to either an immunosorbent enzyme assay or a competitive binding enzyme immunoassay for human serum pancreatic lipase Ohkaru Y., Kurooka S., Kitamura T.	295
Sickle cell screening using a rapid automated HPLC system Shapira E., Miller V.L., Miller J.B., Qu Y.	301

Vol. 183/1, 1989

Carnitine: Metabolism and clinical chemistry Siliprandi N., Sartorelli L., Ciman M., Di Lisa F.	3
Clinical biochemistry of neuron specific enolase Kaiser E., Kuzmits R., Pregant P., Burghuber O., Worofka W.	13
The importance of objective data in the diagnosis of pancreatitis Lott J.A., Ellison E.C., Applegate D.	33
Pancreatic α -amylase assay employing the synergism of two monoclonal antibodies Rauscher E., Gerber M.	41
Monoclonal antibodies to enzymes Rosalki S.B.	45
Serum phospholipase A in hospitalized patients Hoffmann G.E., Hiefinger R., Steinbrueckner B.	59
Increase of creatine kinase and lactate dehydrogenase in the serum of rats submitted to experimental intestinal infarction Roth M., Jaquet P.-Y., Rohner A.	65
Serum arylesterase (paraoxonase) activity following myocardial infarction Secchiero S., Mussap M., Zaninotto M., Bertorelle R., Burlina A.	71
Assay of HMG-CoA reductase activity in the evaluation of cholesterol synthesis in man Carulli N., Tripodi A., Carubbi F.	77
Glutathione-related enzyme activities in testis of patients with malignant diseases Aceto A., Di Ilio C., Angelucci S., Tenaglia R., Zezza A., Caccuri A.M., Federici G.	83
Methods for serum cholinesterase assay and classification of genetic variants Panteghini M.	87
Caffeine as indicator of metabolic functions of microsomal liver enzymes Varagnolo M., Plebani M., Mussap M., Nemetz L., Paleari C.D., Burlina A.	91
Evaluation of pancreatic amylase immunoassay in acute pancreatitis Sacchetti L., Cavalcanti E., Cerasuolo D., Oriani G., Uomo G., Rabitti P.G., Lacetti M., Visconti M., Salvatore F.	95
Assays and structure of reverse transcriptase of the AIDS virus Bertazzoni U., Lori F., Achilli G., Cattaneo E.	101

Vol. 183/2, 1989

Calcium absorption in children estimated from single and double stable calcium isotope techniques Miller J.Z., Smith D.L., Flora L., Peacock M., Johnston Jr. C.C.	107
Time resolved immunofluorometric assay for placental protein 14 Riitinen L., Stenman U.-H., Alfthan H., Suikkari A.-M., Bohn H., Seppala M.	115
Detection of kinetically abnormal argininosuccinate synthase in neonatal citrullinemia by conversion of citrulline to arginine in intact fibroblasts Ben-Yoseph Y., Mitchell D.A.	125
Human cerebrospinal fluid acetylcholinesterase and butyrylcholinesterase. Evidence for identity between the serum and cerebrospinal fluid butyrylcholinesterase Rao R.V., Gnanamuthu C., Balasubramanian A.S.	135
Fluorometric assay of protein in native human bile Harvey P.R.C., Upadhy G.A., Toth J.L., Strasberg S.M.	147

Immunoassay of β -hexosaminidase isoenzymes in serum in patients with raised total activities Isaksson A., Hultberg B.	155
Automated enzymatic assays for the determination of intestinal permeability probes in urine. 2. Mannitol Lunn P.G., Northrop C.A., Northrop A.J.	163
A convenient enzymatic method for the determination of 4-methyl-2-oxopentanoate in plasma: comparison with high performance liquid chromatographic analysis Schadewaldt P., Hummel W., Trautvetter U., Wendel U.	171
A rapid, qualitative ELISA test for the specific detection of morphine in serum or urine Laurie D., Manson A.J., Rowell F.J., Seviour J.	183
A competitive microtitre plate enzyme immunoassay for plasma testosterone using polyclonal anti- testosterone immunoglobulins Rajkowski K.M., Hanquez Ch., Bouzoumou A., Cittanova N.	197
An improved rapid thin-layer chromatographic-gas-liquid chromatographic procedure for the determination of free fatty acids in plasma Tichelaar H.Y., Spinnler Benade A.J., Daubitzer A.K., Kotze T.J.V.W.	207
Salivary corticosteroids in the study of adrenal function Bolufer P., Gandia A., Rodriguez A., Antonio P.	217
ELISA for measuring porphobilinogen deaminase in human erythrocytes Lannfelt L., Wetterberg L., Lilius L., Thunell S., Gellerfors P.	227

Vol. 183/3, 1989

Markers of collagen metabolism in sera of patients with various rheumatic diseases Myllyla R., Becvar R., Adam M., Kivirikko K.I.	243
Serum lipids and cholesterol metabolism during guar gum, plantago ovata and high fibre treatments Miettinen T.A., Tarpila S.	253
Increased furosemide-sensitive sodium influx into erythrocytes in non-insulin-dependent diabetes mellitus associated with hypertension Chimori K., Miyazaki S., Goi R., Dodo S., Hayashi H., Izai M., Goto S., Yasuda K., Miura K.	263
Partial purification and characterization of human hepatoma-derived growth factor Nakamura H., Kambe H., Egawa T., Kimura Y., Ito H., Hayashi E., Yamamoto H., Sato J., Kishimoto S.	273
IgA isotyping of antigliadin antibodies. A possible clue for a less invasive diagnosis of coeliac disease Elewaut A., Dacremon G., Robberecht E., Leroy J., De Baets M.H.	285
CSF-IgM measurement in neurovenereological disease Van Kamp G.J., Wolters Ch. E.	295
Glycosyltransferase activities in normal and leukaemic monocytic cells Francina A., Gateau-Roesch O., Couprie N., Leculier C., Col J.-F., Archimbaud E., Campos L., Louisot P., Richard M.	301
Selective determination of hydroxyproline in urine by high-performance liquid chromatography using precolumn derivatization Teerlink T., Tavenier P., Netelenbos J.C.	309
Small bowel biopsy for disaccharidase levels: Evidence that endoscopic forceps biopsy can replace the Crosby capsule Smith J.A., Mayberry J.F., Ansell I.D., Long R.G.	317
Plasminogen activator activity and plasminogen independent amidolytic activity in tear fluid from healthy persons and patients with anterior segment inflammation Tozser J., Berta A., Punyiczki M.	323
Alterations of purine metabolism in mononuclear cells of individuals at risk of developing type I (insulin-dependent) diabetes mellitus Schafer J.H., Jira T.B., Michaelis D., Rjasanowski I.M.	333

Insulin and glucagon degradation in liver are not affected by hepatic cirrhosis Antoniello S., La Rocca S., Calvacanti E., Auletta M., Salvatore F., Cacciatore L.	343
Increased acylphosphatase levels in erythrocytes from hyperthyroid patients Nassi P., Liguri G., Nediani C., Taddei N., Piccinni P., Degl'Innocenti D., Gheri R.G., Ramponi G.	351
Immunodepletion of albumin from human serum samples Burgess-Cassler A., Johansen J.J., Kendrick N.	359

Vol. 184/1, 1989

A high performance liquid chromatography method for the analysis of ³⁵ S-cystine: Application to the diagnosis of cystinosis Hall N.A., Young E.P.	1
Human fibrinogen heterogeneity. A study of limited fibrinogen degradation Reganon E., Vila V., Aznar J., Laiz B.	7
Purification and characterization of endometrial protein PP14 from mid-trimester amniotic fluid Riittinen L., Julkunen M., Seppala M., Koistinen R., Huhtala M.-L.	19
Immunoreactive anionic and cationic trypsin in human serum Kimland M., Russick C., Marks W.H., Borgstrom A.	31
Analysis of maple syrup urine disease in cell culture: Use of substrates Schadewaldt P., Beck K., Wendel U.	47
3-Methylglutaconyl-CoA hydratase, 3-methylcrotonyl-CoA carboxylase and 3-hydroxy-3-methylglutaryl-CoA lyase deficiencies: A coupled enzyme assay useful for their detection Narisawa K., Gibson K.M., Sweetman L., Nyhan W.L.	57
Radioimmunoassay of human platelet-derived growth factor using monoclonal antibody toward a synthetic 73-97 fragments of its B-chain Shiraishi T., Morimoto S., Itoh K., Sato H., Sugihara K., Onishi T., Ogihara T.	65
The sugar-chain heterogeneity of human γ -glutamyl transferases from the reproductive system and kidney Arai K., Yoshida K.-I., Komoda T., Kobayashi N., Saitoh H., Sakagishi Y.	75
Serum F protein: A new sensitive and specific test of hepatocellular damage Foster G.R., Goldin R.D., Oliveira D.B.G.	85
3-Methylisoxazol-5-one, an artefact from acetoacetic acid formed during urinary organic acid analysis Mills G.A., Walker V., Mellor J.M.	93
Removal of bilirubin from jaundiced serum by immunoaffinity chromatography Izumi Y., Nakajima H.	99
Serum lipolytic activity in Reye's syndrome Kang E.S., Matsuo N., Nagai T., Greenhaw J., Williams P.L.	107
Serum deoxyribonuclease I determined by a radioimmunoassay and an enzymatic assay in malignant diseases Miyachi K., Ogawa M., Murata A., Nakano I., Funakoshi A., Mori T.	115

Vol. 184/2, 1989

Con A affinity of rat α 1-acid glycoprotein (rAGP): Changes during inflammation, dexamethasone or phenobarbital treatment as detected by crossed affino immunoelectrophoresis (CAIE) are not only a reflection of biantennary glycan content Pos O., Drechou A., Durand G., Bierhuizen M.F.A., Van der Stelt M.E., Van Dijk W.	121
Fragmented fibronectin and other synovial fluid proteins in chronic arthritis: Their relation to immune complexes Griffiths A.M., Herbert K.E., Perrett D., Scott D.L.	133
The effect of clomipramine treatment on plasma lipoproteins and high density lipoprotein subfractions in healthy subjects Skinner E.R., Watt C., Reid I.C., Besson J.A.O., Ashcroft G.W.	147

Oxalate, phosphate and sulphate determination in serum and urine by ion chromatography	
Politi L., Chiaraluce R., Consalvi V., Cerulli N., Scandurra R.	155
Glucocorticoid receptors in monocytes in Type 1 diabetes mellitus	
Damm P., Binder C.	167
Pyrimidine 5'-nucleotidase deficiency: Improved detection of carriers	
De Korte D., Sijstermans J.M., Seip M., Van Doorn C.C.H., Van Gennip A.H., Roos D.	175
Plasma lipoproteins in infantile visceral Leishmaniasis: Deficiency of apolipoproteins A-I and A-II	
Bekaert E.D., Kallei R., Bouma M.-E., Lontie J.-F., Mebazaa A., Malmendier C.L., Ayrault-Jarrier M.	181
Effect of renal failure on plasma dihydroxyphenylglycol, 3-methoxy-4-hydroxyphenylglycol, and vanillylmandelic acid	
Hoeldtke R.D., Israel B.C., Cavanaugh S.T., Krishna G.G.	195

Vol. 184/3, 1989

Development of an immunoassay for all human isoferritins, and its application to serum ferritin evaluation	
Cozzi A., Levi S., Bazzigaluppi E., Ruggeri G., Arosio P.	197
Amplification of the proto-oncogenes int-2, c-erb B-2 and c-myc in human breast cancer	
Machotka S.V., Garrett C.T., Schwartz A.M., Callahan R.	207
The estimation of total serum lipids by a completely enzymatic 'summation' method	
Akins J.R., Waldrep K., Bernert Jr. J.T.	219
Rapid homogeneous enzyme immunoassay of plasma protein C	
Suzuki K., Hayashi T., Koyama M., Yoshimura T., Shimamoto M., Kimura N., Kita M.	227
Use of 4-heptylumbelliferyl- β -D-glucoside to identify Gaucher's disease heterozygotes	
Butcher B.A., Gopalan V., Lee R.E., Richards T.C., Waggoner A.S., Glew R.H.	235
Determination of succinylacetone and succinylacetoacetate in physiological samples as the common product 5(3)-methyl-3(5)-isoxazole propionic acid using an isotope dilution method and mass spectrometry	
Schierbeek H., Berger R.	243
Abnormally-fucosylated serum haptoglobins in patients with inflammatory joint disease	
Thompson S., Kelly C.A., Griffiths I.D., Turner G.A.	251
Effect of ischaemia on the activities of human hepatic acyl-CoA: cholesterol acyltransferase and other microsomal enzymes	
Smith J.L., Pillay S.P., De Jersey J., Hardie I.R.	259
Glutathione S-transferase in human bile	
Howie A.F., Hayes P.C., Bouchier I.A.D., Hayes J.D., Beckett G.J.	269
Determination of bone Gla protein (osteocalcin) by enzyme-linked immunosorbent assay	
Egemo C., Dagaard H., Lund B.	279
Enhanced net mass transfer of HDL cholesteryl esters to Apo B-containing lipoproteins in patients with peripheral vascular disease	
Ruhling K., Zabel-Langhennig R., Till U., Thielmann K.	289
Time course of serum amyloid A response in myocardial infarction	
Bausserman L.L., Sadaniantz A., Saritelli A.L., Martin V.L., Nugent A.M., Sady S.P., Herbert P.N.	297
Assay of methylmalonyl CoA mutase with high-performance liquid chromatography	
Kikuchi M., Hanamizu H., Narisawa K., Tada K.	307
Enzyme-labelled immunoassay for urinary gonadotropins using penicillinase	
Desai M., Khatkhatay I., Sankolli G., Meherji P., Joshi U.	315
Effect of low dose acetylsalicylic acid upon plasma thromboxane B_2 levels and platelet aggregation in ischemic stroke patients	
Lee T.-K., Chen Y.-C., Kuo T.-L., Lien I.-N., Liu M.-C., Huang Z.-S.	323

Vol. 185/1, 1989

Carbon dioxide tensions in physiological salt solutions: direct measurements Abrahamsen J., Norrie B., Andersen P.K., Stokke D.B., Nedergaard O.A.	1
A sensitive, direct colorimetric assay of serum copper using 5-Br-PSAA Makino T.	7
Bioluminescent assay for serum adenosine deaminase with immobilized bacterial luciferase Oda K., Yoshida S., Hirose S., Takeda T.	17
The relationship of plasma acetate with glucose and other blood intermediary metabolites in non-diabetic and diabetic subjects Akanji A.O., Humphreys S., Thursfield V., Hockaday T.D.R.	25
Serum alkaline DNase activity in normal or nonhospitalised individuals Avall-Lundqvist E., Economidou-Karaoglou A., Sjøvall K., Lans M., Taper H.S., Roberfroid M., Eneroth P.	35
A preliminary study of serum class II levels in healthy individuals and bone-marrow transplant patients Thompson S., Wareham M., Pearson A.D.J., Sviland L., Turner G.A.	45
Intraerythrocytic water relaxation rates in amyotrophic lateral sclerosis. An NMR investigation Monti J.P., Braguer D., Guelton C., Crevat A., Serratrice G.	53
Radioimmunoassay of neuropeptide Y (NPY) in biological fluids. Circulating NRY levels depend on renal function Stridsberg M., Lundqvist G.	61
Direct extraction and assay of collagenase from human osteoarthritic articular cartilage Blancaert A., Mazieres B., Eeckhout Y., Vaes G.	73
Elastase binding capacity of α_2 -macroglobulin in plasma of patients with asthma or chronic obstructive pulmonary disease, without α_1 -protease inhibitor deficiency Kilroe-Smith T.A., Dowdeswell R.J., Gaillard M.C.	81
Biochemical and neurophysiological parameters in hemodialyzed patients with chronic renal failure Schoots A.C., De Vries P.M.J.M., Thiemann R., Hazejager W.A., Visser S.L., Oe P.L.	91
The effect of vitamin E therapy on plasma and erythrocyte lipid peroxidation in chronic hemodialysis patients Yalcin A.S., Yurtkuran M., Dilek K., Kilinc A., Taga Y., Emerk K.	109
Reactive oxygen species production in whole blood from AIDS patients Guillard O., Sonnet J., Lauwerys R.	113
Oxidative hemolysis in erythrocytes with unstable hemoglobins Falcioni G., Grelloni F., De Sanctis G., Bonfigli A.R., Pierani P., Felici L., Coppa G.V.	115

Vol. 185/2, 1989

Bound amino acids in uremic sera: Search for peptidic middle molecules by preparative polyacrylamide gel electrophoresis and high performance liquid chromatography Kaplan B., Gotfried M., Cojocaru M., Pras M., Ravid M.	119
The quantitation of oxalate in amniotic fluid by ion-chromatography Wandzilak T.R., Hanson F.W., Williams H.E.	131
Effect of 8-anilino-1-naphthalene sulfonic acid (ANS) on the interaction between thyroid hormone and anti-thyroid hormone antibodies Matsuda M., Sakata S., Komaki T., Nakamura S., Kojima N., Takuno H., Miura K.	139
A rapid electrophoretic method for the detection of serum Lp(a) lipoprotein Kawakami K., Tsukada A., Okubo M., Tsukada T., Kobayashi T., Yamada N., Murase T.	147
Evaluation and performance characteristics of a novel ELISA using monoclonal antibody to glycosylated albumin Hud E., Cohen M.P.	157

Gas-liquid chromatographic method for routine detection of plasma sulfo-, gluco-, and free 3-methoxy-4-hydroxyphenylglycol	
Piletz J.E., Halaris A.	165
Radioimmunoassay for erythropoietin using anti-recombinant erythropoietin antibody with high affinity	
Matsubara K., Yoshimura T., Kamachi S., Fukushima M., Hino M., Morii H.	177
Lectin binding characteristics of a cholesterol nucleation promoting protein	
Harvey P.R.C., Upadhy A., Toth J.L., Strasberg S.M.	185
Automated analysis of urinary hydroxyproline	
Ho K.-C., Pang C.-P.	191
Microassay for amino acids in human liquid blood and dried blood	
Ho J.W.	197
The effect of non-enzymatically glycosylated collagen on normal human platelets	
Yarat A., Emerkli N.	203
A simplified procedure for the study of LDL kinetics	
Samman S., Khosla P., Carroll K.K.	207
A combined lectin precipitation method for differentiating modified bone alkaline phosphatase in serum	
Kuwana T., Sugita O., Okada M., Yakata M.	211
Binding of nitroblue tetrazolium to human serum albumin in a fructosamine test	
Sakurai T., Tsuchiya S.	213
Unusual 1 β -hydroxylated bile acids in children with a paucity in interlobular bile ducts	
Kimura A., Mahara R., Tohma M., Ushijima K., Yuge K., Ono E., Yamashita F.	215
German Society for clinical chemistry: Workshop on the standardization of enzyme determinations in medicine	
Schmidt E.	S1
1989 Alkaline phosphatase symposium	
Moss D.	S2
11th National Congress of the Italian Society of clinical enzymology	
Moss D.	S3

Vol. 185/3, 1989

Preface	
Okuda K.	221
Development and applications of ultrasensitive enzyme immunoassays for antigens and antibodies	
Ishikawa E., Hashida S., Tanaka K., Kohno T.	223
A new homogenous enzyme immunoassay using recombinant enzyme fragments	
Khanna P.L., Dworschack R.T., Manning W.B., Harris J.D.	231
Microbial enzymes for creatinine assay: A review	
Shimizu S., Kim J.-M., Yamada H.	241
Diagnostic application of mitochondrial enzymes and isoenzymes	
Schmidt F.W., Schmidt E.	253
Different patterns of leakage of cytosolic and mitochondrial enzymes	
Kamiike W., Fujikawa M., Koseki M., Sumimura J., Miyata M., Kawashima Y., Wada H., Tagawa K.	265
Immunological and ultrastructural analysis of loss of tubular membrane-bound enzymes in patients with renal damage	
Scherberich J.E.	271
Structure of the ornithine transcarbamylase (OTC) gene and DNA diagnosis of OTC deficiency	
Matsuda I., Hata A., Matsuura T., Tsuzuki T., Shimada K.	283
Genetic analysis of familial amyloidotic polyneuropathy, an autosomal dominant disease	
Sakaki Y., Sasaki H., Yoshioka K., Furuya H.	291
Laboratory and clinical features of lactate dehydrogenase subunit deficiencies	
Maekawa M., Kanno T.	299

Phosphoinositide metabolism and oncogenes	
Takenawa T., Fukami K.	309
Separation of hepatoma-associated γ -glutamyltransferase isoenzyme on cellulose acetate media with Triton X-100 and concanavalin A	
Yoshikawa C., Shimojo N., Naka K., Okuda I., Ohkawa J.	317
N-Acetylglucosaminyltransferase III in human serum, and liver and hepatoma tissues: Increased activity in liver cirrhosis and hepatoma patients	
Ishibashi K., Nishikawa A., Hayashi N., Kasahara A., Sato N., Fujii S., Kamada T., Taniguchi N.	325
Regulation of protein kinase C by sphingosine and lysosphingolipids	
Hannun Y.A., Bell R.M.	333
Small molecular weight GTP-binding proteins and signal transduction	
Yamamoto K., Tanimoto T., Kim S., Kikuchi A., Takai Y.	347
Pathogenic potentials of bacterial proteases	
Maeda H., Molla A.	357

Vol. 186/1, 1989

A lack of coordination in the release of urinary lysosomal and brush border enzymes following renovascular surgery	
Holmes R.P., Craddock G., Espeland M.A., Assimos D.G., Dean R.H.	1
Hyperhistaminemia in patients with histidinemia due to increased decarboxylation of histidine	
Tanabe M., Sakura N.	11
Immunochemical characterization of corticosteroid-binding globulin in human bronchoalveolar fluid	
Loric S., Egloff M., Domingo M., Lacroque J., Degrelle H.	19
Ornithine carbamoyltransferase deficiency. A new variant with subnormal enzyme activity	
Rabier D., Benoit A., Petit F., Chekoury A., Bonnefont J.P., Saudubray J.M., Kamoun P.	25
Measurement of the unbound fraction of long chain nonesterified fatty acids (NEFA) in serum: Methodological considerations	
Lim C.-F., Wynne K.N., Stockigt J.R.	31
Origin of serum catalase activity in acute pancreatitis	
Goth L.	39
A rapid and sensitive method for estimating low concentrations of albumin in human urine	
Horton J.K., Davies M., Woodhead J.S., Weeks I.	45
Octadeca-9,11-dienoic acid in cervical intraepithelial neoplasia: A colposcopic study	
Fairbank J., Hollingworth A., Griffin J., Ridgway E., Wickens D., Singer A., Dormandy T.	53
Microheterogeneity of the carbohydrate moiety of human α 1-acid glycoprotein in two benign liver diseases: Alcoholic cirrhosis and acute hepatitis	
Biou D., Chanton P., Konan D., Seta N., N'Guyen H., Feger J., Durand G.	59
Progress and problems in immunoassays for serum pituitary gonadotrophins: Evidence from the UK external quality assessment schemes (EQAS) 1980-1988	
Seth J., Hanning I., Bacon R.R.A., Hunter W.M.	67
The evaluation of a chemiluminescent assay for free thyroxine by comparison with equilibrium dialysis in clinical samples	
Beaman J., Woodhead J.S., Liewendahl K., Mahonen H.	83
On optimising the diagnostic yield of secretin pancreozymin tests	
Hunt L.P., Braganza J.M.	91
A simple method for the measurement of total and glycosylated apolipoprotein B and its relevance to apolipoprotein-B metabolism in diabetes mellitus	
Kortlandt W., Benschop C., Erkelens D.W., Thijssen J.H.H.	109
Characterization of trypsin-activatable latent collagenases and their inhibitors in rheumatoid arthritis synovial fluid	
Tagawa M., Kobayashi J., Akama T.	119

Vol. 186/2, 1990

On the importance of being (stereo) specific Fishman W.H.	129
The human alkaline phosphatases: What we know and what we don't know Harris H.	133
Purification and some properties of the fast migrating alkaline phosphatase in FL-amnion cells (the Kasahara isoenzyme) and its cDNA cloning Higashino K., Muratani K., Hada T., Imanishi H., Amuro Y., Yamamoto Y., Furuyama J., Hirano K., Hong Y.-M., Shimokura M., Hirano T., Kishimoto T.	151
Comparison of the three PLAP-related genes on human chromosome 2 Martin D., Tucker D., Campbell I., Trowsdale J.	165
Cloning and characterization of a cDNA encoding alkaline phosphatase in mouse embryonal carcinoma Hahnel A.C., Schultz G.A.	171
Structure of alkaline phosphatases Kim E.E., Wyckoff H.W.	175
An active site mutant of human placental alkaline phosphatase with deficient enzymatic activity and preserved immunoreactivity Narisawa S., Sowadski J.M., Millan J.L.	189
Purified tetrameric alkaline phosphatase: The effect of treatments with phosphatidylinositol phospholipase C and sodium dodecyl sulfate Hawrylak K., Kihn L., Rutkowski D., Stinson R.A.	197
Allelic and ectopic polymorphisms in human placental alkaline phosphatases: Sugar chain heterogeneities Komoda T., Sato M., Furiya K., Sakagishi Y., Sekine T.	203
Secretion of hepatic and intestinal alkaline phosphatases: Similarities and differences Alpers D.H., Eliakim R., DeSchryver-Kecakemeti K.	211
Human fetal intestinal alkaline phosphatase: Molecular heterogeneity and immunological detection in amniotic fluids Verpooten G.F., Hoylaerts M.F., Nouwen E.J., De Broe M.E.	225
Alkaline phosphatase of rat intestinal lymph: Its characterization and the effect of fat administration Miura S., Yoshioka M., Hamada Y., Morita A., Asakura H., Tsuchiya M.	239
Alkaline phosphatase releasing activity in human tissues Hamilton B.A., McPhee J.L., Hawrylak K., Stinson R.A.	249
Enzymatic heterogeneity of seminomas Koshida K., Wahren B.E.	255
Purification and partial characterization of the placental-like alkaline phosphatase in human lung tissue Hirano K., Koyama I., Stigbrand T.	265
Serum placental alkaline phosphatase (PLAP) in gynecologic malignancies — With special reference to the combination of PLAP and CA54/61 assay Nozawa S., Udagawa Y., Ohkura H., Negishi Y., Akiya K., Inaba N., Takamizawa H., Kimura E., Terashima Y.	275
Placental-type alkaline phosphatase in peritoneal fluid of women with endometriosis Kang J.O., Hudak W.A., Crowley W.J., Criswell B.S.	285
Placental-like alkaline phosphatase from ovary and ascites of a patient with a malignant ovarian tumour Koyama I., Stendahl U., Stigbrand T.	295
Histochemical, ultracytochemical and biochemical study of alkaline phosphatase activity during gastric carcinogenesis Hirota N., Sakai T., Komoda T.	301
An alkaline phosphatase reacting with both monoclonal antibodies to intestinal and placental isoenzymes Imanishi H., Hada T., Muratani K., Hirano K., Higashino K.	309

- The preparation of monoclonal antibodies which react preferentially with human bone alkaline phosphatase and not liver alkaline phosphatase
Hill C.S., Wolfert R.L. 315

Vol. 186/3, 1990

- Two ELISA's for measurement of protein S, and their use in the laboratory diagnosis of protein S deficiency
Deutz-Terlouw P.P., Ballering L., Van Wijngaarden A., Bertina R.M. 321
- Transmembrane cation fluxes and fatty acid composition of erythrocytes in psoriatic patients
Corrocher R., Bassi A., Gandini A., Guarini P., Trevisan M.T., Schena D., Olivieri O., Ferrari S. 335
- Experimental studies on unconjugated bilirubin binding by human erythrocytes
Hayer M., Piva M.-T., Sieso V., Magnan de Bornier B. 345
- Serial measurements of Ca antigen in sera from patients with advanced malignant breast disease
Searle F., Chantler S.M., Goodall A.B., Adam T., Bagshawe K.D., Mitchell H.D.C., Boxer G.M., Southall P.M. 351
- Human liver nicotinamide N-methyltransferase: Ion-pairing radiochemical assay, biochemical properties and individual variation
Rini J., Szumlanski C., Guercioli R., Weinshilboum R.M. 359
- Quantitation of hydrogen peroxide formed during UV-visible irradiation of protoporphyrin, coproporphyrin and uroporphyrin
Menon I.A., Becker M.A.C., Persad S.D., Haberman H.F. 375
- Quantitative assay of free and total carnitine using tandem mass spectrometry
Kodo N., Millington D.S., Norwood D.L., Roe C.R. 383
- Enzyme immunoassay for urinary albumin at low concentration in diabetes mellitus
Niwa T., Katsuzaki T., Tatemichi N., Kato K., Yazawa T., Maeda K. 391
- Low biotinidase activities in the sera of patients with impaired liver function: evidence that the liver is the source of serum biotinidase
Grier R.E., Heard G.S., Watkins P., Wolf B. 397

Vol. 187/1, 1990

- Characterization of prolidase I and II from erythrocytes of a control, a patient with prolidase deficiency and her mother
Ohhashi T., Ohno T., Arata J., Sugahara K., Kodama H. 1
- Sensitive enzyme immunoassay for human Cu/Zn superoxide dismutase
Kurobe N., Suzuki F., Okajima K., Kato K. 11
- Immunochemical assays of serum proteins: A European external quality assessment survey and the effects of calibration procedures on interlaboratory agreement
Bullock D.G., Dumont G., Vassault A., Aguzzi F., Chambers R.E., Milford Ward A., Whicher J.T., Bienvenu J. 21
- Validation of a method to measure the uptake of [³H]dopamine by human platelets
Dean B., Kulkarni J., Copolov D.L. 37
- Influence of angiotensin converting enzyme inhibitor (captopril) on kidney epithelial cells in vitro: Studies on potassium (⁸⁶Rb) influx and cellular proliferation
Bauer H., Lametschwandtner A., Steiner M., Bauer H.C. 47
- Accumulation of mannosyl- β (1-4)-N-acetylglucosamine in fibroblasts and leukocytes of patients with a deficiency of β -mannosidase
Van Pelt J., Hokke C.H., Dorland L., Duran M., Kamerling J.P., Vliegenthart J.F.G. 55

Vol. 187/2, 1990

Detection of neuron specific enolase concentrations in cerebrospinal fluid from patients with neurological disorders by means of a sensitive enzyme immunoassay Vermuyten K., Lowenthal A., Karcher D.	69
Automated enzymatic assays for the determination of intestinal permeability probes in urine. I. Lactulose and lactose Northrop C.A., Lunn P.G., Behrens R.H.	79
A new sensitivity microassay for the measurement of erythrocyte glycogen Farquharson J., Jamieson E.C., MacPhee G.B., Logan R.W.	89
Development and validation of an automated enzyme assay for paracetamol (acetaminophen) Morris H.C., Overton P.D., Ramsay J.R., Campbell R.S., Hammond P.M., Atkinson T., Price C.P.	95
Characterization of aminopeptidases in human kidney soluble fraction Mantle D., Lauffat B., McDermott J., Gibson A.	105
Post-transcriptional modification of serum creatine kinase in infected intensive care patients Delanghe J.R., De Buyzere M.L., De Scheerder I.K., Van Rostenberghe H.L., Faust U., Rodenbach J.M., Kruse-Jarres J.D., Wieme R.J.	115
Comparison of serum and urine neopterin concentrations in patients with HIV-1 infection Fuchs D., Jaeger H., Popescu M., Reibnegger G., Werner E.R., Kaboth W., Wachter H.	125
A fluorimetric enzyme assay for the diagnosis of Morquio disease type A (MPS IV A) Van Diggelen O.P., Zhao H., Kleijer W.J., Janse H.C., Poorthuis B.J.H.M., Van Pelt J., Kamerling J.P., Galjaard H.	131
Elevation of hormone-binding globulins in acute intermittent porphyria Herrick A.L., McColl K.E.L., Wallace A.M., Moore M.R., Goldberg A.	141
Dye-stained gelatin microcarriers as insoluble chromolytic substrates for the determination of proteolytic activity Safarik I.	149
Enzymatic erythrocyte creatine determinations as an index for cell age Buysse A.M., Delanghe J.R., De Buyzere M.L., De Scheerder I.K., De Mol A.M., Noens L.	155
A microalbuminuria assay using bromphenol blue Jung K., Nickel E., Pergande M.	163
A sensitive and specific assay for granulocyte elastase in inflammatory tissue fluid using L-pyroglyutamyl-L-prolyl-L-valine-p-nitroanilide Tanaka H., Shimazu T., Sugimoto H., Yoshioka T., Sugimoto T.	173
Alterations in manganese and iodide contents in human thyroid tumors; a correlation between the contents of essential trace elements and the states of malignancy Nishida M., Sakurai H., Tezuka U., Kawada J., Koyama M., Takada J.	181
Spectrophotometric evaluation of results of the ascorbate and the ascorbate-cyanide screening tests for glucose-6-phosphate dehydrogenase deficiency Rakitzis E.T., Manolakou K., Papandreou P.	189

Vol. 187/3, 1990

Kawasaki disease and oxidative metabolism Lebranchu Y., Malvy D., Richard M.J., Arnaud J., Favier A., Bardos P.	193
Sialylacto-N-fucopentaose II (sialylated Le ^a) coupled to human serum albumin used as standard in immunoassays of tumour associated antigens CA 19-9 and CA-50 in serum Masson P., Lundblad A., Wieslander J.	199
Characterization of a monoclonal antibody specific for prostatic secretory protein of 94 amino acids (PSP94) and development of a two-site binding enzyme immunoassay for PSP94 Von Der Kammer H., Kraus E., Aumüller G., Scheit K.H.	207
Demonstration of a combined deficiency of xanthine oxidase and aldehyde oxidase in xanthinuric patients not forming oxipurinol Reiter S., Simmonds H.A., Zollner N., Braun S.L., Knedel M.	221

Microheterogeneity of α_1 -acid glycoprotein: Variation during the menstrual cycle in healthy women, and profile in women receiving estrogen-progestogen treatment	
Succari M., Foglietti M.-J., Percheron F.	235
Comparative evaluation of the diagnosis of acute pancreatitis based on serum and urine enzyme assays	
Gwozdz G.P., Steinberg W.M., Werner M., Henry J.P., Pauley C.	243
Anti-skeletal muscle antibodies in the sera from myasthenic patients with thymoma: Identification of anti-myosin, actomyosin, actin, and α -actinin antibodies by a solid-phase radioimmunoassay and a Western blotting analysis	
Ohta M., Ohta K., Itoh N., Kurobe M., Hayashi K., Nishitani H.	255
Sensitive enzyme immunoassay for human aldolase A	
Okajima K., Kurobe N., Shimizu K., Kato K.	265
Variations in serum alpha-L-fucosidase activity during childhood and pregnancy	
Vayse J., Pilardeau P., Gattegno L.	273
A routine method for the measurement of the sodium, potassium, magnesium and calcium content of human lymphocytes	
Weissberg N., Brooks B.A., Schwartz G., Eylath U., Abraham A.S.	281
How should urinary cotinine concentrations be adjusted for urinary creatinine concentration?	
Thompson S.G., Barlow R.D., Wald N.J., Van Vunakis H.	289
Isoelectric focusing versus quantitative measurements in the detection of intrathecal local synthesis of IgG	
Luxton R.W., McLean B.N., Thompson E.J.	297
Non-M CK — A practical measure of creatine kinase isoenzymes in cancer patients	
McGing P.G., Teeling M., McCann A., Kyne F., Carney D.N.	309
Measurement of L-carnitine and acylcarnitines in body fluids and tissues in children and in adults	
De Sousa C., English N.R., Stacey T.E., Chambers R.A.	317
Early amniocentesis and amniotic fluid organic acid levels in the prenatal diagnosis of organic acidemias	
Coude M., Chadeaux B., Rabier D., Kamoun P.	329
A carry-over study in three different instruments for the determination of serum glucose: The importance of student's <i>t</i> test	
Yucel D., Kantaroglu N., Saydam G.S., Balk M., Top S.	333

Vol. 188/1, 1990

Metal inhibition of ferrochelatase activity in human lymphocytes	
Rossi E., Attwood P.V., Garcia-Webb P., Costin K.A.	1
Isolation, identification, and analysis of 4-acetylamino-phenol-glucuronide in body fluids of dialyzed renal patients; a molecular mass marker for peritoneal diffusive transport	
Schoots A.C., Koomen G.C.M., Struijk D.G., Krediet R.T., Arisz L.	15
The effect of synthetic very low calorie diets on the GH-IGF-I axis in obese subjects	
Gama R., Teale J.D., Marks V.	31
A new hemoglobin variant found during investigations of diabetes mellitus: Hb Pavie (α 135 (H18) Val—Glu)	
Wajcman H., Blouquit Y., Riou J., Kister J., Poyart C., Soria J., Galacteros F.	39
A simple serum galactosyltransferase assay method suitable for routine use	
Sichel F., Malas J.-P., Gauduchon P., Aubert M., Bar E., Le Tallec J.-Y.	49
A comparison of serum mucoprotein with serum α_1 acid glycoprotein, haptoglobin, and α_1 antitrypsin assays in monitoring inflammatory bowel disease	
Lubega J., Davies T.J.	59
Does electrophoresis reliably screen for high serum lipoprotein Lp(a)?	
Bruckert E., Truffert J., De Gennes J.L., Lagarde J.P., Federspiel C.B.M.C., Giral P., Dairou F., Bernard C.	71
Urinary paraquat in acute intoxication	
Kuo T.-L.	79

Urinary organic acids in elderly Japanese patients with ketosis and encephalopathy who have taken panto-yl-gamma-aminobutyrate, calcium salt (calcium hopantenate, HOPA)	
Nakanishi T., Funahashi S., Shimizu A., Hayashi A.	85
Newletter of the Societa Italiana di Biochimica Clinica	
Deledda R.	92

Vol. 188/2, 1990

Urinary excretion of brush-border antigen and plasma proteins in early stages of diabetic nephropathy	
Mutti A., Alinovi R., Ghiggeri G.M., Bergamaschi E., Candiano G., Rasi A., Gusmano R., Franchini I., Borghetti A.	93
Early alterations of plasma free amino acids in chronic renal failure	
Ceballos I., Chauveau P., Guerin V., Bardet J., Parvy P., Kamoun P., Jungers P.	101
Identification of an intermediate form of β -N-acetylhexosaminidase in chorionic villi	
Pieretti M., Beccari T., Cosmi E.V., Stirling J.L., Orlacchio A.	109
The increased plasma Lp(a): B lipoprotein particle concentration in angina pectoris is not associated with hypofibrinolysis	
Alessi M.C., Parra H.J., Joly P., Vu-Dac N., Bard J.M., Fruchart J.C., Juhan-Vague I.	119
Usefulness of a prognostic inflammatory and nutritional index in pediatric clinical practice	
Pressac M., Vignoli L., Aymard P., Ingenbleek Y.	129
Decreased xanthine oxidase activities and increased urinary oxypurines in heterozygotes for hereditary xanthinuria	
Kawachi M., Kono N., Mineo I., Yamada Y., Tarui S.	137
Quantitative determination of IgM-rheumatoid factor by enzyme immunoassay — Standardization using a serum from a rheumatoid arthritis patient	
Oka H., Hirohata S., Inoue T., Iwamoto S., Miyamoto T.	147
Increased urinary excretion of β -hydroxyisovaleric acid in ketotic and nonketotic type II diabetes mellitus	
Yu W., Kuhara T., Inoue Y., Matsumoto I., Iwasaki R., Morimoto S.	161
Nuclear magnetic resonance studies of lithium transport in erythrocyte suspensions of hypertensives	
Ramasamy R., Mota de Freitas D., Bansal V.K., Dorus E., Labotka R.J.	169
Improved specific assay for serum arylesterase using a water-soluble substrate	
Kawai H., Sakamoto F., Inoue Y.	177
Elevated spermidine level in pleural fluid of patients with lung cancer	
Abe K., Saito T., Kitamura S., Konaka R.	183
Stability of gonadotropins in urine	
Khatkhatay I., Desai M., Sankolli G., Joshi U.	185

Vol. 188/3, 1990

Immunochemical determination of lipoprotein Lp(a): Comparison of Laurell electrophoresis and ELISA	
Kostner G.M., Gries A., Pometta M., Molinari E., Pichler F., Aicher H.	187
Additional fucosyl residues on membrane glycoproteins but not a secreted glycoprotein from cystic fibrosis fibroblasts	
Wang Y.-M., Hare T.R., Won B., Stowell C.P., Scanlin T.F., Glick M.C., Hard K., Van Kuik J.A., Vliegthart J.F.G.	193
The phospholipid composition of erythrocyte ghosts and plasma lipoproteins in diabetes type I in children	
Watala C., Jozwiak Z.	211
O-Phosphohydroxylysineuria: A new inborn error of metabolism?	
Dorland L., Duran M., De Bree P.K., Smith G.R., Horvath A., Tibosch A.S., Wadman S.K.	221

Diurnal variation of erythrocyte sodium-lithium countertransport rate and intracellular cation concentrations	
Siebers R.W.L., Maling T.J.B.	227
Carboxypeptidase activity in human urine from healthy subjects and renal disease patients	
Hamai K., Ikeda R., Sumi H., Mihara H.	233
Inosine triphosphate pyrophosphohydrolase deficiency in a kindred with adenosine deaminase deficiency	
Duley J.A., Simmonds H.A., Hopkinson D.A., Levinsky R.J.	243
Evaluation of a radioimmunoassay for somatomedin-C/insulin-like growth factor I (Sm-C/IGF-I) in human plasma	
Giannella-Neto D., Cavaleiro A.M., Sadoyama R., Wajchenberg B.L., Spencer E.M.	253
The effect of incubation with <i>Staphylococcus aureus</i> Cowan I on subsequent determinations of IgG, IgM, and IgM rheumatoid factor by ELISA	
Nesher G., Moore T.L.	261
Spectrophotometric estimation of erythrocyte ATP and deoxyadenosine triphosphate using hexokinase and glucokinase	
Nakashima K., Nakashima H., Shimoyama M.	267
Succinic acidemia is not a new syndrome of organic acidemia	
Elpeleg O.N., Hurvitz H.	271
Monoclonal gammopathies. Their identification and biological significance (I)	
Radl J.	273

Vol. 189/1, 1990

Influence of human transferrin and group-specific protein on endotoxicity in vitro	
Berger D., Winter M., Beger H.G.	1
Diagnosis of alcaptonuria: Rapid analysis of homogentisic acid by HPLC	
Bory C., Boulieu R., Chantin C., Mathieu M.	7
65 Zinc absorption in patients with insulin-dependent diabetes mellitus assessed by whole-body counting technique	
Kiilerich S., Hvid-Jacobsen K., Vaag A., Sorensen S.S.	13
Effect of diet on urinary excretion of unconjugated catecholamines and some of their metabolites in healthy control subjects and depressed patients	
Moleman P.	19
The effect of n-3 polyunsaturated fatty acids on lipids, platelet function, coagulation, fibrinolysis and monocyte chemotaxis in patients with hypertension	
Schmidt E.B., Nielsen L.K., Pederson J.O., Kornerup H.J., Dyerberg J.	25
Direct reading glucose electrodes detect the molality of glucose in plasma and whole blood	
Fogh-Anderson N., Wimberley P.D., Thode J., Siggaard-Andersen O.	33
Use of a new simplified assay for phospholipase A2 to measure bacterial enzyme levels	
Lumb M.R., Roseblade C.K., Helmig R., Uldbjerg N., Sullivan M.H.F., Elder M.G.	39
The acute phase protein response in mice does not show tolerance to recurrent sterile inflammation	
Todd N.J., Whicher J.T., Westacott C., Gilbert A.	47
International study on the comparability of Apo A-I and Apo B methods	
Grafnetter D., Molinari E., Lonsky L.	55
High performance liquid chromatography methods for studying protein binding of aluminium in human serum in the absence and in the presence of desferrioxamine	
Garcia Alonso J.I., Lopez Garcia A., Parajon J., Blanco Gonzalez E., Sanz Medel A., Cannata J.B.	69
S-Adenosylhomocysteine hydrolase and adenosine deaminase activities in human red cell ageing	
Bozzi A., Furciniti-La Chiusa B., Strom R., Crifo C.	81
Changes in phospholipid composition of tracheal aspirates from newborns with hyaline membrane disease or transient tachypnoea	
Bourbon J.R., Francoual J., Magny J.F., Lindenbaum A., Leluc R., Dehan M.	87

Vol. 189/2, 1990

Development of a double antibody radioimmunoassay for quantitation of 1 α ,25-dihydroxyvitamin D	
Armbruster F.-P., Reichel H., Vogel G., Georgousis H., Schmidt-Gayk H.	97
Distribution of HDL ₂ and HDL ₃ in a random population of healthy French males and females — evaluation by a two-step precipitation procedure	
Atger V., Wirbel E., Roche D., Apfelbaum M., Burstein M., Girard-Globa A.	111
Detection and estimation of the barley prolamins content of beer and malt to assess their suitability for patients with coeliac disease	
Ellis H.J., Freedman A.R., Ciclitira P.J.	123
The levels of adenine nucleotides and pyridine coenzymes in red blood cells from the newborn, determined simultaneously by HPLC	
Formato M., Masala B., De Luca G.	131
Human liver L-alanine-glyoxylate aminotransferase: Characteristics and activity in controls and hyperoxaluria type I patients using a simple spectrophotometric method	
Wanders R.J.A., Ruiter J., Van Roermund C.W.T., Schutgens R.B.H., Ofman R., Jurriaans S., Tager J.M.	139
Hyperphosphatasemia related to three intestinal alkaline phosphatase isoforms: Biochemical study	
Bentouati L., Samadi Baboli M., Hachem H., Hamza M., Canal P., Soula G.	145
Detection of apolipoprotein E polymorphisms using PCR/ASO probes and Southern transfer: Application for routine use	
Houlston R.S., Wenham P.R., Humphries S.E.	153
Is basic red cell ferritin a more specific indicator than serum ferritin in the assessment of iron stores in the elderly?	
Galan P., Sangare N., Preziosi P., Roudier M., Hercberg S.	159
Energy dependence of cytosolic enzyme efflux from rat skeletal muscle	
West-Jordan J.A., Martin P.A., Abraham R.J., Edwards R.H.T., Jackson M.J.	163
Determination of α_1 -antitrypsin in fecal extracts by enzyme immunoassay	
Brouwer J., Smekens F.	173
Cutaneous interstitial fluid protein concentrations in the inflammatory syndrome: Pharmacological consequences	
Clerc C., Pibouin M., Ruelland A., Legras B., Chevrant-Breton J., Cloarec L.	181
Combined diagnostic value of biochemical markers in acute pancreatitis	
Viel J.F., Foucault P., Bureau F., Albert A., Drosowsky M.A.	191
Rapid spectrophotometric assay of serum iron	
Johnson D.J., Williams H.L.	199
Protease activities in gastric cancer tissues	
Chung S.M., Kawai K.	205
Glomerular and urinary glycosaminoglycans in diabetic rats	
Reddi A.S.	211
Human platelet thermostable phenol sulfotransferase: Assay of frozen samples and correlation between frozen and fresh activities	
Anderson R.J., Liebentritt D.K.	221
Inhibition of lactate dehydrogenase by an EDTA-Ca ²⁺ complex	
Shina R., Condrea E.	231
Plasma polymorphonuclear leukocyte elastase- α_1 -proteinase inhibitor complex in patients with gastric ulcer	
Okajima K., Inoue M., Koga S., Okabe H., Takatsuki K.	237
Kinetic enzymatic assay for ethylene glycol	
Hansson P.	243
Studies on the clinical significance of serum α_2 HS-glycoprotein measurement in metabolic bone diseases	
Scarnecchia L., Minisola S., Pacitti M.T., Carnevale V., Bigi F., Romagnoli E., Mazzuoli G.F.	245

- Proper handling of blood samples is crucial for the accurate ion chromatographic determination of oxalate
Petraculo M., Marangella M. 249
- Inhibition by a free-radical scavenger of ascorbate-induced hemoglobin denaturation in glucose-6-phosphate dehydrogenase deficient erythrocytes
Papandreou P.T., Rakitzis E.T. 253

Vol. 189/3, 1990

- A time-resolved fluoroimmunoassay for the determination of prostaglandin F(2 α)
Luke F.J., Schlegel W. 257
- Immunochemical detection of human blood in feces
Uchida K., Matsuse R., Miyachi N., Okuda S., Tomita S., Miyoshi H., Hirata I., Tsumoto S., Ohshiba S. 267
- Development of a radioimmunoassay for neurone specific enolase (NSE) and its application in the study of patients receiving intra hepatic arterial streptozotocin and floxuridine
Cunningham R.T., Johnston C.F., Irvine G.B., McIlrath E.M., McNeill A., Buchanan K.D. 275
- Apolipoprotein B-48 analysis by high-performance liquid chromatography in VLDL: A sensitive and rapid method
Hidaka H., Kojima H., Nakajima Y., Aoki T., Nakamura T., Kawabata T., Nakano T., Harano Y., Shigeta Y. 287
- Enzymatic diagnosis of 3-hydroxy-3-methylglutaryl-CoA lyase deficiency with high-performance liquid chromatography
Kikuchi M., Narisawa K., Tada K., Sweetman L. 297
- Decreased proapolipoprotein A-I processing in liver disease: Evidence for hepatic participation in proapolipoprotein A-I conversion
Isobe H., Sakai H., Satoh M., Sakamoto S., Koga S., Nawata H. 303
- Lipoamidase activity in human serum is due to biotinidase
Garganta C.L., Wolf B. 313
- Rapid diagnosis of 3-hydroxy-3-methylglutaryl-coenzyme A lyase deficiency via enzyme activity measurements in leukocytes or platelets using a simple spectrophotometric method
Wanders R.J.A., Zoeters P.H.M., Schutgens R.B.H., De Klerk J.B.C., Duran M., Wadman S.K., Van Sprang F.J., Hemmes A.M., Voorbrood B.S. 327
- Evidence of acid hydrolysis as responsible for 2-(2-furoyl)-4(5)-(2-furanyl)-1H-imidazole (FFI) production
Gerhardinger C., Lapolla A., Crepaldi G., Fedele D., Ghezzi E., Seraglia R., Traidi P. 335

Vol. 190/1-2, 1990

- The biology of transferrin
De Jong G., Van Dijk J.P., Van Eijk H.G. 1
- Incidence and properties of aspartate aminotransferase-immunoglobulin complexes in patients with a high serum aspartate to alanine aminotransferase ratio
Moriyama T., Nobuoka M., Makino M. 47
- A compensatory mechanism improving red cell membrane fluidity in hemodialysed patients
Peuchant E., Motta C., Sales C., Clerc M. 57
- Liquid chromatographic studies of vitamin B6 metabolism in man
Edwards P., Liu P.K.S., Rose G.A. 67
- A method for the rapid detection of urinary glycopeptides in α -N-acetylgalactosaminidase deficiency and other lysosomal storage diseases
Schindler D., Kanzaki T., Desnick R.J. 81
- Heterogeneity of urinary albumin from diabetic patients
Hayashi Y., Morikawa A., Makino M. 93

Lack of correlation between human serum angiotensin converting enzyme activity and the renin-angiotensin-aldosterone system	
Pedraza-Chaverri J., Herrero B., Ibarra-Rubio M.E., Cruz C., Tapia E., Pena J.C.	105
Seed oils containing fatty acids with conjugated double bonds as substrates for lipase assay	
Anzai T., Kotake I., Komine S.	107
Serum immunoreactive prolyl 4-hydroxylase as a specific marker of hepatocellular necrosis	
Nagai Y., Toda G.	109
Urinary adenosine deaminase binding protein, a marker of contrast media induced acute renal damage	
Parvez Z., Patel N.B., Nelson J.E.	111
Glucose interferes with inulin determination by the resorcinol method: Overestimation of glomerular filtration rate in diabetic patients	
Erman A., Van Dijk D.J., Pansky R., Wittenberg C., Boner G., Rosenfeld J.B.	115
Plasma fibronectin in patients affected by psoriasis: Relation with the extension of skin involvement and complications	
Cecchi R., Brunetti L., Giomi A., Gironi A., De Giorgio L.A., Seghieri G.	119
Improved HPLC method for the determination of cephalosporins in human plasma and a new solid-phase extraction procedure for cefazolin and ceftizoxime	
Moore C.M., Sato K., Hattori H., Katsumata Y.	121
Scientific Division. Committee on enzymes. IFCC methods for the measurement of catalytic concentration of enzymes. Part 7. IFCC method for creatine kinase (ATP: creatine N-phosphotransferase, EC 2.7.3.2). IFCC recommendation	
Horder M., Elser R.C., Gerhardt W., Mathieu M., Sampson E.J.	S4
Scientific Division. Committee on pH, blood gases and electrolytes. Guidelines for transcutaneous p(O ₂) and p(CO ₂) measurement	
Wimberley P.D., Burnett R.W., Covington A.K., Maas A.H.J., Mueller-Plathe O., Siggaard-Andersen O., Weisberg H.F., Zijlstra W.G.	S41
Education division expert panel on quantities and units. A protocol for the conversion of clinical laboratory data	
Lehmann H.P., Worth H.G.J., Zinder O.	S51

Vol. 190/3, 1990

Redox electrode for monitoring dehydrogenase-catalyzed reactions	
Chen S.S.-H.	129
Enzyme immunoassay for flecainide	
Freche J.P., Coudert G., Wellman M., Siest G.	139
Myoadenylate deaminase deficiency studies on normal and deaminase-deficient skeletal muscle	
Kaletha K., Nowak G.	147
Elevation of blood lactate and pyruvate levels in acute intermittent porphyria — A reflection of haem deficiency?	
Herrick A.L., Fisher B.M., Moore M.R., Cathcart S., McColl K.E.L., Goldberg A.	157
Comparison of the effects of purified human α_1 -antichymotrypsin and α_1 -proteinase inhibitor on NK cytotoxicity: Only α_1 -proteinase inhibitor inhibits natural killing	
Laine A., Leroy A., Hachulla E., Davril M., Dessaint J.-P.	163
Enzyme immunosorbant assay of oestradiol in unextracted plasma using penicillinase as label	
Pandey P.K., Shrivastav T.G., Kumari G.L., Rao P.N., Grover P.K., Murthy H.G.K.	175
Immunoreactive vasopressin in end stage renal failure	
Argent N.B., Baylis P.H., Wilkinson R.	185
Theoretical isoelectric points and electric charges of mutated human hemoglobin subunits	
Meireles Ribeiro J., Ruiz A., Sillero M.A.G., Sillero A.	189
A sensitive radioimmunoassay using a monoclonal antibody that is equipotent for ercalcitriol and calcitriol (1,25-dihydroxy vitamin D ₂ and D ₃)	
Mawer E.B., Berry J.L., Cundall J.P., Still P.E., White A.	199

The effect of oral contraceptive use and pregnancy on the daily rhythm of cortisol and cortisone Meulenberg M.N. P.M., Hofman J.A.	211
Does Tamm-Horsfall mucoprotein inhibit or promote calcium oxalate crystallization in human urine? Grover P.K., Ryall R.L., Marshall V.R.	223
Two-site assay of intact parathyroid hormone in primary hyperparathyroidism: Studies in basal conditions, following adenoma removal and during calcium and EDTA infusion Mazzuoli G., Minisola S., Scarnecchia L., Pacitti M.T., Carnevale V., Romagnoli E., Bigi F., Bianchi G.	239
Urinary proteins and red blood cell membrane negative charges in diabetes mellitus Bernard A., Amor A.O., Goemare-Vanneste J., Antoine J.-L., Lauwerys R., Colin I., Vandeleene B., Lambert A.	249
Immunological study on hypophosphatasia Goseki M., Oida S., Takagi Y., Okuyama T., Watanabe J., Sasaki S.	263
Dopamine-mediated effects induced by metoclopramide simultaneously on adrenocorticotrophic hormone and prolactin in patients with pituitary and/or hypothalamic disorders Nishida S., Matsuki M., Horino M., Kawai Y., Tsushima K., Yoneda M., Oyama H., Ishii R.	269
Different molecular forms of acid-stable trypsin inhibitor in normal and uremic plasma Yoshida E., Sumi H., Mihara H.	277
The effect of non-insulin-dependent diabetes on serum concentrations of tumor-associated carbohydrate antigens of Ca19-9, CA-50, and sialyl SSEA-I in association with the Lewis blood phenotype Shimojo N., Naka K., Nakajima C., Ishizaki T., Okuda K., Murai J., Yamamoto M.	283
Serum alkaline phosphatase in sickle cell anaemia Ogunleye I.O., Adekile A.D., Fakunle J.B.	291
Presence of thiosulfate in human plasma Lundquist P.	293

Vol. 191/1-2, 1990

Concentrations of N-terminal ProANP in human plasma: evidence for ProANP (1-98) as the circulating form Buckley M.G., Sagnella G.A., Markandu N.D., Singer D.R.J., MacGregor G.A.	1
Detection of aspartylglycosaminuria using urine specimens recovered from absorbent filter paper Kartinen V., Mononen I.	15
Free testosterone levels in plasma and saliva as determined by a direct solid-phase radioimmunoassay: a critical evaluation Rey F., Chiodoni G., Braillard K., Berthod C., Lemarchand-Beraud T.	21
Determination of aluminium in bone in haemodialyzed patients, using inductively coupled argon plasma emission spectrometry Leflon P., Plaquet R., Moriniere P., Fournier A.	31
Stable isotope dilution analysis of cystine in granulocyte suspensions as cysteine: a powerful method for the diagnosis, the follow up, and treatment of patients with cystinosis Schierbeek H., Bijsterveld K., Chapman T.E., Van Luijk W.H.J., Reijngoud D.J., Berger R.	39
Studies on lipamidase: characterization of the enzyme in human serum and breast milk Backman-Gullers B., Hannestad U., Nilsson L., Sorbo B.	49
Presence of reverse transcriptase in human leukemias and lymphomas Moyret C.F., Bernard D.J., Maurizis J.-C., Chassagne J., Plagne R., Chollet P.	61
Plasma oxalate and creatinine and oxalate/creatinine clearance ratios in normal subjects and in primary hyperoxaluria. Evidence for renal hyperoxaluria Kasidas G.P., Nemat S., Rose G.A.	67
Longitudinal changes in serum fructosamine do not parallel those in glycated haemoglobin in young adults with insulin-dependent diabetes Fisken R.A., Chan A.W., Hanlon A., MacFarlane I.A.	79

Lectin binding to serum 5'-nucleotidase and alkaline phosphatase in liver disease Massoubre C., Verdier P., Mefflah K., Masliah C., Perrin D., Orsonneau J.-L., Harb J.	87
Why serum F protein is a sensitive and specific marker of liver damage in man Grewal G.K., Oliveira D.B.G.	93
Enzymatic determination of serum total and esterified cholesterol with the same reagents Matsushita M., Irino T., Maehata E.	97
Ribonuclease activities in liver tissue of patients with chronic liver diseases Tsuji H., Murai K., Akagi K., Fujishima M.	101
Labile HbA(1c) formation in fractionated erythrocytes in aerobic and anaerobic conditions Yoshida Y., Nakashima K.	105
Storage and thawing influence plasma levels of immunoreactive atrial natriuretic peptide Tan A.C.I.T.L., Kloppenborg P.W.C., Benraad Th. J.	111
Bayes' theorem should be considered when making decision limits Ohman S.	115
On the role of thyroglobulin as an antigen of thyroid hormone autoantibodies Sakata S.	119

Vol. 191/3, 1990

Clinical biochemistry of preeclampsia and related liver diseases of pregnancy: A review Freund G., Arvan D.A.	123
Increased immunoreactivity of apolipoprotein B epitopes during prolonged storage of low density lipoproteins La Belle M., Mok K.S., Karu A.E., Krauss R.M.	153
Histamine release with a solid-phase-coupled allergen Mita H., Tadokoro K., Yasueda H., Shida T.	161
A simple enzyme-linked immunoassay of human carbonic anhydrase I for the study of developing erythroid cells Kondo T., Murakami K., Isobe H., Taniguchi N., Kawakami Y.	169
Angiotensin I-converting enzyme activity in puromycin aminonucleoside-nephrotic syndrome Arevalo A.E., Ibarra-Rubio M., Cruz C., Pena J.C., Pedraza-Chaverri J.	175
Comparative biochemical and immunological studies on gamma-glutamyltransferases from human kidney and renal cell carcinoma applying monoclonal antibodies Fischer P., Scherberich J.E., Schoepp W.	185
Amino acid profiles by HPLC after filter paper sampling: 'Appropriate technology' for monitoring of nutritional status Lundsjo A., Hagelberg S., Palmer K., Lindblad B.S.	201
One-step sandwich enzyme immunoassay for human laminin using monoclonal antibodies Iwata K.	211
Gamma-glutamyltransferase from human hepatoma cell lines: Purification and cell culture of HepG2 on microcarriers Visvikis A., Goergen J.-L., Oster T., Bagrel D., Wellman M., Marc A., Engasser J.-M., Siest G.	221
A reversed-phase high-performance liquid chromatographic method for the determination of pyruvate dehydrogenase complex activity in human platelets and cultured skin fibroblasts Maehara M., Kajita M., Kitoh M., Watanabe K.	233
Interference of abnormal hemoglobins on the measurement of hemoglobin A(1c) by ion-exchange chromatography Engbaek F., Sorensen G.H., MacIntyre B., Clausen I., Lund H.T., Jastrup B.	239

Vol. 192/1, 1990

Ferritin and transferrin levels in human breast cyst fluids: Relationship with intracystic electrolyte concentrations Troccoli R., Stella F., Biagioni S., Battistelli S., Cerroni L., Marcheggiani F., Artico M., Terzano C., Mannello F.	1
---	---

Cytochrome oxidase deficiency affecting the structure of the myofibre and the shape of mitochondrial cristae membrane	
Sumegi B., Melegh B., Adamovich K., Trombitas K.	9
Interaction between endogenous circulating sulfated-glycosaminoglycans and plasma proteins	
Pasquali F., Oldani C., Ruggiero M., Magnelli L., Chiarugi V., Vannucchi S.	19
Gunn rats: A reproducible experimental model to compare the different methods of measurements of bilirubin serum concentration and to evaluate the risk of bilirubin encephalopathy	
Labruere P., Myara A., Trivin F., Odievre M.	29
Radioimmunoassay for sequence 38-54 of human progastrin: Increased diagnostic specificity of gastrin-cell diseases	
Van Solinge W.W., Rehfeld J.F.	35
Radioimmunoassay for plasma neuropeptide-Y in physiological and physiopathological states and response to sympathetic activation	
Miller M.A., Sagnella G.A., Markandu N.D., MacGregor G.A.	47
Evaluation of a simple method for the measurement of cytidine deaminase in serum and comparison with a reference method	
Thompson P.W., James I.T., Rowell L., Jones D.D.	55
Urinary polyamine and metabolite excretion by children with Zellweger's syndrome	
Govaerts L.C.P., Van den Berg G.A., Theeuwes A., Muskiet F.A.J., Monnens L.A.H.	61
The effect of excess daily methionine intake on plasma homocysteine after a methionine loading test in humans	
Andersson A., Brattstrom L., Israelsson B., Isaksson A., Hultberg B.	69
One-step solvolysis of 3-, 7- and 12-sulfated free and conjugated bile acids	
Princen H.M.G., Meijer P., Kuipers F.	77
A simple procedure to exclude false-positive galactose screening test results caused by antibiotics	
Clemens P.C., Pletner C., Vellguth B.	85

Vol. 192/2, 1990

The relationship between erythrocyte aspartate aminotransferase activity and the 'TPP effect' in Wernicke's encephalopathy and other thiamine deficiency states	
Nixon P.F., Price J., Norman-Hicks M., Williams G.M., Kerr R.A.	89
Measurement of β_2 -microglobulin, retinol-binding protein, α_1 -microglobulin and urine protein I in healthy children using enzyme-linked immunosorbent assay	
Tomlinson P.A., Dalton R.N., Turner C., Chantler C.	99
Enzyme immunoassay of urinary β -hexosaminidase isoenzymes in patients with renal transplants	
Hultberg B., Isaksson A., Sterner G., Weibull H.	107
A new, rapid, one-step method for the isolation of platelets from human blood	
Ford T.C., Graham J., Rickwood D.	115
Erythrocyte nucleotide stability and plasma hypoxanthine concentrations: Improved ATP stability with short-term storage at room temperature	
Simmonds H.A., Micheli V., Davies P.M., McBride M.B.	121
Stable isotope dilution analysis of very long chain fatty acids in plasma, urine and amniotic fluid by electron capture negative ion mass fragmentography	
Stellaard F., Ten Brink H.J., Kok R.M., Van Den Heuvel L., Jakobs C.	133
Is dehydroepiandrosterone sulfate a natriuretic digitalis-like factor in human serum?	
Goto A., Yamada K., Ishii M., Yoshioka M., Sugimoto T.	145
The antioxidant activity of human semen	
Kocak T., Unal T., Tokar N.K.	153

Vol. 192/3, 1990

Purines, lactate and phosphate release from child and adult heart during cardioplegic arrest	
Smolenski R.T., Swierczynski J., Narkiewicz M., Zydzio M.M.	155

A new approach to the newborn screening for hyperphenylalaninemia: Use of L-phenylalanine dehydrogenase and microtiter plates	
Wendel U., Koppelkamm M., Hummel W., Sander J., Langenbeck U.	165
Sensitive enzyme immunoassay for human Mn superoxide dismutase	
Kurobe N., Inagaki T., Kato K.	171
Failure to conserve lactose and glucose polymers during frozen storage of fecal specimens: Methods for preservation	
Murray R.D., Ailabouni A.H., Powers P.A., Juhling McClung H., Ulysses B., Li K., Heitlinger L.A., Sloan H.R.	181
One-step sandwich enzyme immunoassay for soluble human thrombomodulin using monoclonal antibodies	
Kodama S., Uchijima E., Nagai M., Mikawatani K., Hayashi T., Suzuki K.	191
Excretion of epidermal growth factor (EGF) in diabetes	
Lev-Ran A., Hwang D.L., Miller J.D., Josefsberg Z.	201
HPLC separation of alkaline phosphatase isoenzymes. Application to the amniotic fluid	
Ragoucy-Sengier C., Rolland H., Pileire B.	207

Vol. 193/1-2, 1990

Evidence for locally synthesized and clonally restricted immunoglobulin in the synovial fluid from rheumatoid arthritis patients	
Carpenter A.B., Huczko E., Eisenbeis Jr. C.H., Kelly R.H.	1
Serum alpha-1-proteinase inhibitor with abnormal properties in ovarian cancer	
Thompson S., Wong E., Cantwell B.M.J., Turner G.A.	13
Collagen of the normal and the varicose human saphenous vein: A biochemical study	
Maurel E., Azema C., Deloly J., Bouissou H.	27
Serum baseline hyaluronate and disease activity in rheumatoid arthritis	
Kontinen Y.T., Saari H., Honkanen V.E.A., Szocsik K., Mussalo-Rauhama H., Tulensalo R., Friman C.	39
Biliary excretion of iron in healthy man and in patients with alcoholic cirrhosis of the liver	
Raedsch R., Stiehl A., Walker S., Theilmann L., Kommerell B., Waldherr R., Senn M.	49
Studies of circulating parathyroid hormone following parathyroidectomy in renal osteodystrophy	
Straffen A.M., Carmichael D.J.S., Fairney A., Hulme B., Snell M.E.	55
The immunological evidence for a phenylalanine hydroxylase like immunoreactive protein in different human cells and tissues	
Petruschka L., Rebrin I., Grimm U., Herrmann F.H.	65
Altered protein kinase C and protein kinase A activities in erythrocyte membrane, platelets and lymphocytes of Duchenne muscular dystrophy (DMD) patients	
Jagadeesh G., Lavanya M., Anandaraj M.P.J.S., Anjaneyulu A.	79
Evaluation of the particle size distribution of plasma lipoprotein in a patient with Tangier disease by high performance liquid chromatography	
Takizawa A., Komoda T., Hokari S., Sakagishi Y., Tanaka A., Hara I.	85
Decreased muscle enoyl-CoA hydratase activities in patients with exercise intolerance	
Carroll J.E., McGuire B.S., Hartlage P.L., Howard J.C.	89
Differential diagnosis of an hyperamylasemia in an acute abdominal syndrome	
Foucault P., Drosowsky M.A.	91
Hyperphosphatasemia related to three intestinal alkaline phosphatase isoforms: Biochemical study	
Bentouati L., Samadi Baboli M., Hachem H., Hamza M., Canal P., Soula G.	93

Vol. 193/3, 1990

Polymorphonuclear elastase in neonatal sepsis	
Tsaka T., Herkner K.R.	103

Reduction of urinary bile alcohol excretion and serum cholestanol in patients with cerebrotendinous xanthomatosis after oral administration of deoxycholic acid Wolthers B.G., Van der Molen J.C., Walrecht H., Hesselmann L.F.G.M.	113
Use of pralidoxime-Pd(II) complex for the spectrophotometric determination of the cholinesterase reactivator (PAM-2C1) in the urine Karlijkovic-Rajic K.D., Stankovic B.S.	119
Mass concentration measurements of creatine kinase BB isoenzyme as an index of brain tissue damage Delanghe J.R., De Winter H.A., De Buyzere M.L., Camaert J.J., Martens F.E., De Praeter C.	125
Proteolysis of human apolipoprotein B: Effect on quantitative immunoturbidimetry De Bruin T.W.A., Vos M.C., Kortlandt W., Bouma B.N., Erkelens D.W.	137
Effect of fish-oil and vitamin E supplementation on lipid peroxidation and whole-blood aggregation in man Brown J.E., Wahle K.W.J.	147
Radial immunodiffusion and immunoelectrophoresis compared for identifying autoantibodies to lactate dehydrogenase in human serum Harff G.A., Backer E.T.	157
Lipoprotein particles in homozygous familial hypercholesterolemic patients treated with portacaval shunt and LDL apheresis Gairin D., Monard F., Cachera C., Bard J.-M., Amouyel P., Duriez P., Tacquet A., Fruchart J.-C.	165
Effect of ethanol on metabolism of the hypouricemic agents allopurinol and benzbromarone Kaneko K., Fujimori S., Ishizuka I., Akaoka I.	181
Redox electrode for monitoring oxidase-catalyzed reactions Chen P.T., Chen S.S.-H.	187
Quantitation of tyrosine-O-sulfate in human urine by ion-pair reverse-phase high-performance liquid chromatography Suiiko M., Prasantha Fernando P.H., Arino Y., Terada M., Nakatsu S., Liu M.-C.	193
Spectrophotometric determination of oxalate in whole blood Hatch M.	199

Vol. 194/1, 1990

The widening horizons of bioanalytical mass spectrometry Hill R.E.	1
Analytical methodology for immunoassays and DNA hybridization assays — Current status and selected systems — Critical review Diamandis E.P.	19
Ultrasensitive enzyme immunoassay Ishikawa E., Hashida S., Kohno T., Hirota K.	51
Chemiluminescent DNA probes: A comparison of the acridinium ester and dioxetane detection systems and their use in clinical diagnostic assays Nelson N.C., Kacian D.L.	73
Fluorescence spectroscopy and its application to a new generation of high sensitivity, multi-microspot, multianalyte, immunoassay Ekins R., Chu F., Biggart E.	91

Vol. 194/2-3, 1990

Simple method for the measurement of antioxidants Novak Z., Varga S.I., Pataki L., Matkovic B.	115
The effect of doxorubicin on the glutathione content and viability of cultured human lung cancer cell lines A549 and GLC ₄ 210 Al-Kabban M., Stewart M.J., Watson I.D., Reglinski J.	121

Serum concentrations of carotenoids, retinol, and α -tocopherol in healthy persons determined by high-performance liquid chromatography Ito Y., Ochiai J., Sasaki R., Suzuki S., Kusuha Y., Morimitsu Y., Otani M., Aoki K.	131
A delayed-addition enzyme immunoassay for the relative cholesteryl ester transfer protein mass in patients with deficient plasma cholesteryl ester transfer activity Yamashita S., Sprecher D.L., Matsuzawa Y., Sakai N., Tarui S., Harmony J.A.K., Wetterau J.R.	145
The facile detection of the nt 1226 mutation of glucocerebrosidase by 'mismatched' PCR Beutler E., Gelbart T., West C.	161
Inhibition of protein kinase C activation by low concentrations of aluminium Cochran M., Elliott D.C., Brennan P., Chawtur V.	167
gamma-Glutamylglutamine identified in plasma and cerebrospinal fluid from hyperammonaemic patients Hammond J.W., Potter M., Truscott R., Wilcken B.	173
Immunological quantification of human hexokinase type I Magnani M., Serafini G., Chiarantini L., Corsi D., Stocchi V.	185
Further characterization of a heat-stable alkaline phosphatase with low sensitivity to L-phenylalanine Onica D., Rosendahl K., Waldenlind L.	193
Creatine metabolism during metabolic perturbations in patients with organic acidurias Davies S.E.C., Iles R.A., Stacey T.E., Chalmers R.A.	203
An HPLC-ECD procedure for measuring total phenolsulfotransferase (PST) activity in human liver, platelets and blood Khoo B.Y., Sit K.H., Wong K.P.	219
The contribution of sex, electrophoretic phenotype, pregnancy and race to the variability of delta-aminolevulinate dehydrase (ALADH) levels in human erythrocytes. A study in Black mixed Brazilians De Freitas Sousa M.G., Passos Bueno M.R., Olympio da Silva M.C.B., De Borja Goncalves Duarte A.F., Azevedo E.S.	229

Vol. 195/1,2, 1990

Activities of mitochondrial aspartate aminotransferase and creatine kinase isoenzyme MB in serum following coronary bypass surgery Huseby N.-E., Grostad M., Almdahl S.M., Vaage J.	1
Short- and long-term effects of LDL-apheresis on lipoprotein (a) serum levels Ritter M.M., Suhler K., Richter W., Schwandt P.	9
A new sandwich-ELISA method for the determination of keratan sulphate peptides in biological fluids employing a monoclonal antibody and labelled avidin biotin technique Kongtawelert P., Ghosh P.	17
Interactions of pancreatic secretory trypsin inhibitor in small intestinal juice: Its hydrolysis and protection by intraluminal factors Freeman T.C., Davies R., Calam J.	27
The action of myocristin on membrane bound sulphhydryl groups McGrae F., Reglinki J., Smith W.E., Brzeski M., Sturrock R.D.	41
Atomic absorption and UV-VIS absorption spectrophotometric determination of oxalate in urine by ligand exchange extraction Leyva J.A.M., Artiga M.P.H., Mendez M.M.A., Perez J.J.Q.	47
Binding of corticotropin-releasing hormone (CRH) in maternal and fetal plasma and in amniotic fluid Salminen-Lappalainen K., Laatikainen T.	57
Characterisation of the main molecular forms of human fecal immunoglobulin A Kapel N., Meillet D., Isacki S., Bouvet J.P., Gobert J.G., Raichvarg D.	67
The effect of polycythemia and hypoxia on urinary N-acetyl- β -D-glucosaminidase activity in newborns Csathy L., Poci I., Kiss L., Balla G., Modi N., Price R.G.	77

- Immunoelectrophoretic pattern of α_2 -macroglobulin and α_2 -macroglobulin-protease complexes
Donnelly P.K., Trotter P., White M., Shenton B.K. 87
- Letter to the Editor*
- Plasma lipids and apolipoproteins in end-stage liver disease
Lontie J.-F., Dubois D.Y., Malmendier C.L., Mathe D., Adam R., Gigou M., Bismuth H. 93

Vol. 195/3, 1991

- Superiority of thyrotropin to thyroxine as a tool in the screening for congenital hypothyroidism by the filter paper spot technique
Bourdoux P.P., Van Thi H.V., Courtois P.A., Ermans A.M. 97
- Total and ultrafiltrable plasma magnesium in hyper- and hypoparathyroidism, and in calcium-related metabolic disorders
Claeysens S., Lavoine A., Daragon A., Josse S., Godin M., Matray F., Kuhn J.M. 107
- A rapid extraction method for serum calcitonin
Bucht E., Granberg B., Sjostedt U., Torring O. 115
- Exopeptidases in human platelets: An indication for proteolytic modulation of biologically active peptides
Scharpe S.L., Vanhoof G.C., De Meester I.A., Hendriks D.F., Van Sande M.E., Muylle L.M., Yaron A. 125
- Stability of β_2 -microglobulin and retinol binding protein at different values of pH and temperature in normal and pathological urine
Blumsohn A., Morris B.W., Griffiths H., Ramsey C.F. 133
- Effects of smoking on HDL subfractions in myocardial infarction patients: Effects on lecithin-cholesterol acyltransferase and hepatic lipase
Moriguchi E.H., Fusegawa Y., Tamachi H., Goto Y. 139

Vol. 196/1, 1991

- A rapid method for determination of human C-peptide in plasma
Saelsen L., Tronier B., Madsbad S., Juel Christensen N. 1
- Effects of selenium supplementation on blood and urine selenium levels and liver function in patients with primary biliary cirrhosis
Valimaki M., Alfthan G., Vuoristo M., Ylikahri R. 7
- High-performance liquid chromatographic determination of plasma glycolic acid in healthy subjects and in cases of hyperoxaluria syndromes
Petrarulo M., Marangella M., Linari F. 17
- Influx of urea during bronchoalveolar lavage depends on the permeability of the respiratory membrane
Van de Graaf E.A., Jansen H.M., Weber J.A., Koolen M.G.J., Out T.A. 27
- Letters to the Editor*
- Erythrocyte sodium, potassium and sodium fluxes with cell and subject ageing
Heseltine D., Thomas T.H., Wilkinson R., James O.F.W., Potter J.F. 41
- Disparity between apolipoprotein E phenotypes and genotypes (as determined by polymerase chain reaction and oligonucleotide probes) in patients with non-insulin-dependent diabetes mellitus
Snowden C., Houlston R.S., Arif M.H., Laker M.F., Humphries S.E., Alberti K.G.M.M. 49
- In vivo effect of simvastatin on lipoprotein cholesterol ester metabolism in normocholesterolemic volunteers
Magot T., Malmendier C.L., Ouguerram K., Lontie J.F., Lutton C. 59

Vol. 196/2-3, 1991

- Modulative effects of cell adhesion peptide (Arg-Gly-Asp-Ser) on the aggregation of stimulated platelets from ophthalmic patients
Babizhayev M.A., Mucha A.I. 77

Aminopeptidase P and dipeptidyl peptidase IV activity in human leukocytes and in stimulated lymphocytes	
Hendriks D., De Meester I., Umel T., Vanhoof G., Van Sande M., Scharpe S., Yaron A.	87
Application of sialidase antibodies for the diagnosis of clostridial infections	
Roggentin P., Hobrecht R., Tirpitz D., Rothe B., Schauer R.	97
Restriction fragment length polymorphisms in the Apo B gene in relation to coronary heart disease in a Southern Asian population	
Mendis S., Shepherd J., Packard C.J., Gaffney D.	107
Lipid peroxidation and protective enzymes during myocardial infarction	
Loeper J., Goy J., Rozenzajn L., Bedu O., Moisson P.	119
Hydrogen sulphide and total acid-volatile sulphide in faeces, determined with a direct spectrophotometric method	
Florin T.H.J.	127
Characterization of dipeptidyl and tripeptidyl aminopeptidases in human kidney soluble fraction	
Mantle D.	135
A simple method for determination of serum catalase activity and revision of reference range	
Goth L.	143
Influence of splenectomy on the properties of erythrocytes with Hemoglobin Volga disease	
Falcioni G., Grelloni F., Bertoli E., Felici L., Valentino Coppa G.	153
Assessment of the automated colorimetric and the high-performance liquid chromatographic methods for nicotine intake by urine samples of smokers' smoking low- and medium-yield cigarettes	
Kolonen S.A., Puhakainen E.V.J.	159
Clinical relevance of the quantification of apolipoprotein E in cerebrospinal fluid	
Carlsson J., Armstrong V.W., Reiber H., Felgenhauer K., Seidel D.	167
The association between plasma β -hexosaminidase and its isoenzyme patterns and retinopathy in Type 1 diabetes mellitus	
Hultberg B., Isaksson A., Agardh E., Agardh C.-D.	177
Serum laminin P1 fragment concentration in renal diseases	
Horikoshi S., Koide H.	185
Purification and properties of human serum carnosinase	
Jackson M.C., Kucera C.M., Lenney J.F.	193
Aminopeptidase A in human placenta and pregnant serum	
Okuyama T., Ishiura S., Nojima M., Tsukahara T., Yanagida M., Sugita H.	207
Erythrocyte sodium in chronic liver disease with ascites	
Ramakrishna B.S., Kumaresan R., Balasubramanian K.A.	217
4-Hydroxybutyric aciduria: Application of a fluorometric assay to the determination of succinic semialdehyde dehydrogenase activity in extracts of cultured human lymphoblasts	
Gibson K.M., Lee C.F., Chambliss K.L., Kamali V., Francois B., Jaeken J., Jakobs C.	219

Vol. 197/1, 1991

The nature of the proteins lost from isolated rat skeletal muscle during experimental damage	
Jackson M.J., Page S., Edwards R.H.T.	1
An immunologic pathway for intravascular catabolism of creatine kinase subform MM3	
Kanemitsu F., Okigaki T.	9
A sensitive assay of transthyretin (prealbumin) in human cerebrospinal fluid in nanogram amounts by ELISA	
Vatassery G.T., Quach H.T., Smith W.E., Benson B.A., Eckfeldt J.H.	19
Carnitine in dried blood spots: A method suitable for neonatal screening	
Barns R.J., Bowling F.G., Brown G., Clague A.E., Thompson A.	27
Purification and characterization of leucyl aminopeptidase and pyroglutamyl aminopeptidase from human skeletal muscle	
Mantle D., Lauffart B., Gibson A.	35

Modified hemoglobin in hypoglycemia caused by hyperinsulinism	
Niketic V., Maric S., Diklic A., Neskovic S., Tomasevic N.	47
Collagen peptidase and type III procollagen peptide serum levels in chronic liver diseases	
Muller A., Krombholz B., Pott G., Machnik G., Vollandt R., Reinhardt M., Jorke D.	59
Low density lipoprotein receptor levels and the response to dietary and pharmacological intervention in polygenic hypercholesterolemia	
Joven J., Vilella E., Masana L., Ansoleaga J.	67
Porphobilinogen deaminase in acute intermittent porphyria: Activity and concentration in erythrocytes and lymphocytes	
Lilius L., Lannfelt L., Wetterberg L., Floderus Y., Henrichson A., Thunell S.	77

Vol. 197/2, 1991

Serum ferritin spectrotypes in patients with heterozygous β -thalassaemia	
Lasne Y., Francina A., Benzerara O.	85
The relationship between plasma free fatty acids and experimentally induced hepatic encephalopathy in the rat	
Smit J.J.H., Bosman D.K., Jorning G.G.A., De Haan J.G., Maas M.A.W., Chamuleau R.A.F.M.	95
Determination of gangliosides and sulfatide in human cerebrospinal fluid with a microimmunoaffinity technique	
Davidsson P., Fredman P., Manasson J.-E., Svennerholm L.	105
Low serum creatine kinase in patients with infective endocarditis	
De Scheerder I.K., Delanghe J.R., De Buyzere M.L., Hollanders G., Clement D.L., Leroux-Roels G.G.	117
Measurement of serum Apo A1 and Apo B: Comparison of immunoturbidimetric and rate nephelometric techniques	
Glueck C.J., McCray C., Speirs J.	123
Multiple forms of γ -glutamyl transpeptidase in human submandibular gland	
Sajjan A.R., Hinchigeri S.B., Datta K.S.	133
Determination of serum iron by aas: Deproteinization versus direct analysis	
De Benzo Z.A., Fraile R., Gomez C., Carrion N.	141
Letter to the Editor	
Reference intervals for retinol-binding protein in serum and urine	
Berg B., Fex G., Tryding N., Nilsson J.-E., Ekman S.	149
Erratum: One-step sandwich enzyme immunoassay for human laminin using monoclonal antibodies (<i>Clinica Chimica Acta</i> (1990) 191 (211-220))	
Fujimoto N., Sejima H., Kyotani M., Iawata K., Ichida T., Ooshima A.	157

Vol. 197/3, 1991

An enzyme immunoassay for determining total thyroxine in human serum using an ultramicro-analytical system	
Gonzalez R.R., Robaina R., Rodriguez M.E., Blanca S.	159
The presence and measurement of secretory component in human bile and blood	
Perez J.H., Van Schaik M., Mullock B.M., Bailyes E.M., Price C.P., Luzio J.P.	171
Stability of human blood serum aminoacids after storage at different pH and temperature conditions	
Kornhuber M.E., Balabanova S., Heiligensetzer G.V., Kornhuber C., Zettlmeissl H., Kornhuber A.W.	189
Modulation of cholesterol 7α -hydroxylase activity by nonspecific lipid transfer protein in human liver — possibly altered regulation of its cytosolic level in patients with gallstones	
Kawata S., Imai Y., Inada M., Inui Y., Kakimoto H., Fukuda K., Maeda Y., Tarui S.	201

A sensitive, direct colorimetric assay of serum zinc using nitro-PAPS and microwell plates Makino T.	209
Hypogalactosylation of immunoglobulin G sugar chains and elevated serum interleukin 6 in Castleman's disease Nakao H., Nishikawa A., Nishiura T., Kanayama Y., Tarui S., Taniguchi N.	221
Intact parathyroid hormone assay is superior to mid region assay in the EDTA-infusion test in hyperparathyroidism Bergenfels A., Norden N.E., Ahren B.	229
Characterization of human plasma sialyltransferase using a novel fluorometric assay Gross H.J., Brossmer R.	237
Quantitative analysis of urinary pyroglutamic acid in patients with hyperammonemia Nakanishi T., Shimizu A., Saiki K., Fujiwara F., Funahashi S., Hayashi A.	249
Isoforms of human serum biotinidase Hart P.S., Hymes J., Wolf B.	257
Delay in cell separation, storage and anticoagulants induced inaccuracies in measuring plasma non-esterified fatty acids McGann A.M., Hodson A.W.	265

Vol. 198/1-2, 1991

Analysis of immunoglobulin and T cell receptor genes. Part I: Basic and technical aspects Van Dongen J.J.M., Wolvers-Tettero I.L.M.	1
Analysis of immunoglobulin and T cell receptor genes. Part II: Possibilities and limitations in the diagnosis and management of lymphoproliferative diseases and related disorders Van Dongen J.J.M., Wolvers-Tettero I.L.M.	93

Vol. 198/3, 1991

Ionselective minielectrode determination of ionic and total calcium concentrations in mixed saliva Lagerlof F., Matsuo S.	175
An enzyme-linked lectin assay for sialidase Lambre C.R., Terzidis H., Greffard A., Webster R.G.	183
Detection of the common Hb F Sardinia [(A) γ (E19)Ile \rightarrow Thr] variant by isoelectric focusing in normal newborns and in adults affected by elevated fetal hemoglobin syndromes Masala B., Manca L.	195
Glyceroluria in healthy adults, mentally ill adults and children selected for metabolic screening Kohlshutter A., Seitz H.J., Feldmann B., Lehnert W., Langenbeck U.	203
Facts and artefacts in mevalonic aciduria: Development of a stable isotope dilution GCMS assay for mevalonic acid and its application to physiological fluids, tissue samples, prenatal diagnosis and carrier detection Hoffmann G.F., Sweetman L., Bremer H.J., Hunneman D.H., Hyanek J., Kozich V., Lehnert W., Nyhan W.L., Speidel I., Trefz F.K., Gibson K.M.	209
Significance of variation in turnover of glycated albumin on indices of diabetic control Johnson R.N., Easdale R.W., Tatnell M., Baker J.R.	229
Decreased oxidant-induced proteolysis in erythrocytes with enhanced antioxidative defence enzymes due to Down's syndrome Bartosz G., Kedziora J., Retelewska W.	239
Accelerated in vitro degradation of CCK-58 in blood and plasma of patients with acute pan- creatitis Springer C.J., Eberlein G.A., Eysselein V.E., Schaeffer M., Goebell H., Calam J.	245
Consequences of the presence of elongated variant of the major transmembrane protein (Band 3 protein) in the human erythrocyte Retelewska W., Gaczynska M., Bartosz G., Judkiewicz L.	255

Serum aluminium and zinc and other variables in patients with and without cognitive impairment in the community	
Haines A., Iliffe S., Morgan P., Dormandy T., Wood B.	261
Changes in proteolytic susceptibility of erythrocyte membrane proteins in hereditary spherocytosis	
Gaczynska M., Bartosz G., Judkiewicz L.	267
The effect of n-3 polyunsaturated fatty acids on Lp(a)	
Berg Schmidt E., Klausen I.C., Kristensen S.D., Lervang H.-H., Faergeman O., Dyerberg J.	271
<i>Letter to the Editor</i>	
Nomenclature standardization on an international basis	
Fuentes-Arderiu X.	279

Vol. 199/1, 1991

Lysosomal enzyme activity in human aqueous humor	
Weinreb R.N., Jeng S., Miller A.L.	1
Leukocyte sonicates as a source for both enzyme assay and DNA amplification for mutational analysis of certain lysosomal disorders	
Louie E., Rafi M.A., Wenger D.A.	7
Screening for disorders of pyruvate metabolism by measuring the ratio of the rates of lactate production and pyruvate decarboxylation in cultured skin fibroblasts	
Kuroda Y., Kawakami I., Kobashi H., Naito E., Ito M., Saijo T., Yokota I., Takeda E.	17
Determination of α -amylase using a new blocked substrate (3-ketobutylidene β -2-chloro-4-nitrophenyl-maltopentaoside)	
Teshima S., Hayashi Y., Emi S., Ishimaru K.	23
High-performance liquid chromatographic determination of urinary cysteine and cystine	
Birwe H., Hesse A.	33
Clinical evaluation of high-performance affinity chromatography for the separation of bone and liver alkaline phosphatase isoenzymes	
Gonchoroff D.G., Branum E.L., Cedel S.L., Riggs B.L., O'Brien J.F.	43
Serum fructosamine/haemoglobin A ₁ ratio predicts the future changes in haemoglobin A ₁ in type 2 (non-insulin dependent) diabetic patients	
MacRury S.M., Kilpatrick E.S., Paterson K.R., Dominiczak M.H.	51
Determination of membrane cholesterol in normal and pathological red blood cells	
Macchia T., Mancinelli R., Barbini D.A., Taggi F., Avico U., Cantafora A.	59
Investigation of glycosylation processes in mitochondria and microsomal membranes from human skeletal muscle	
Gasnier F., Lerne F., Rousson R., Roussouly P., Vaganay E., Louisot P., Gateau-Roesch O.	69
Stable isotope dilution assay for 5 β -cholestane-3 α ,7 α ,12 α ,25-tetrol and 5 β -cholestane-3 α ,7 α ,12 α ,23,25-pentol in human serum using [26,27-D ₈] labeled internal standards; a highly accurate approach to the serological diagnosis of cerebrotendinous xanthomatosis	
Kosaka D., Hiraoka T., Kohoda T., Kajiyama G., Yamauchi T., Kihira K., Kuramoto T., Hoshita T.	83
Simultaneous analysis of plasma phenethylamine, phenylethanolamine, tyramine and octopamine in patients with hepatic encephalopathy	
Yonekura T., Kamata S., Wasa M., Okada A., Kawata S., Tarui S.	91
Quantitative analysis of human erythrocyte spectrin by sodium dodecyl sulfate-polyacrylamide gel electrophoresis using pancreatic trypsin inhibitor (Kunitz) as an internal standard	
Yoshida H., Takamatsu S., Satoh K., Kawamura Y., Imaizumi T., Teshimori R., Kawachi K.	99
<i>Letters to the Editor</i>	
Elevated serum bile acids in hepatic vein occlusion and idiopathic portal hypertension	
Imai Y., Kawata S., Tarui S.	109
The effective inhibition of the Maillard reaction by guanabenz acetate, and its relevance to the prevention of diabetic complications	
Igaki N., Yamada H., Oimomi M., Kasuga M.	113

Effect of temperature on red cell ATP

Beutler E.

117

Vol. 199/2, 1991

- An enzyme-linked immunosorbent assay (ELISA) for the detection and quantitation of the tumor marker 1-methylxanthine in human urine
D'Ambrosio S.M., Gibson-D'Ambrosio, Trewyn R.W. 119
- Lipoprotein-cyclodextrin interaction
Sharma A., Janis L.S. 129
- Platelet activating factor acetylhydrolase activity in the urine of patients with renal disease
Kirschbaum B. 139
- Quantitation of serum apolipoprotein A-I, A-II, B, C-II, C-III and E in healthy Japanese by turbidimetric immunoassay: Reference values, and age- and sex-related differences
Noma A., Hata Y., Goto Y. 147
- Ecdysteroid-like compounds in human urine: They can occur in the absence of any parasitic infection
Gharib B., Baswaid S., Quilici M., De Reggi M. 159
- Changes in ceruloplasmin activity and lactate concentration in patients at high risk of acute organ system failure
Dauberschmidt R., Mrochen H., Forster I., Stumpe C., Dressler C., Grajetzki H., Meyer M. 167
- Determination of medium chain acyl-CoA dehydrogenase activity in cultured skin fibroblasts using mass spectrometry
Niezen-Koning K.E., Chapman T.E., Mulder I.E., Smit G.P.A., Reijngoud D.J., Berger R. 173
- N-Acetyl- β -D-glucosaminidase (NAG) isoenzymes release from human monocyte-derived macrophages in response to zymosan and human recombinant interferon- γ
Bourbouze R., Raffi F., Dameron G., Hali-Mirafab H., Loko F., Vilde J.-L. 185
- Response of urinary angiotensin to challenges of the renin-angiotensin system
Boer P., Vos P.F., Koomans H.A., Rabelink A.J., Beutler J.J., Dorhout Mees E.J. 195
- A gas chromatographic method for the determination of neutral steroid profiles in urine, including studies on the effect of oxytetracycline administration on these profiles in men
Hamalainen E., Fotsis T., Adlercreutz H. 205

Vol. 199/3, 1991

- Immunological studies of plasma cholinesterase during pregnancy and the puerperium
Whittaker M., Jones J.W., Braven J. 223
- Characteristics of hexosaminidase A in homogenates of white blood cells using methylumbelliferyl-N-acetyl- β -D-glucosaminide-6-sulphate as substrate
Den Tandt W.R., Scharpe S. 231
- O-(2,3,4,5,6-Pentafluorobenzyl)oxime-trimethylsilyl ester derivatives for sensitive identification and quantitation of aldehydes, ketones, and oxoacids in biological fluids
Hoffmann G.F., Sweetman L. 237
- Affinity electrophoresis of alkaline phosphatase using polyacrylamide gels
Ramasamy I. 243
- Spectrofluorimetric determination of urinary p-aminobenzoic and p-aminosalicylic acids in the BT-PABA/PAS test of pancreatic function
Pemberton P.W., Gagjee P., Chaloner C., Braganza J.M., Lobley R.W. 253
- Isotope dilution gas chromatographic-mass spectrometric method for the determination of lignans and isoflavonoids in human urine, including identification of genistein
Adlercreutz H., Fotsis T., Bannwart C., Wahala K., Brunow G., Hase T. 263
- Agarose gel electrophoresis of glucose-6-phosphate-dehydrogenase isoenzymes
Forbes J., Steytler J.G., Van Heerden R. 279

Studies on the effects of aminopterin and methotrexate on the release of prostaglandin PGE ₂ , PGD ₂ , and PGI ₂ from small intestine in the rat in vivo	
Weiler H., Moser U., Weiler C., Gerok W.	283
Bile acid and phospholipid fatty acid composition in bile of patients with cholesterol and pigment gallstones	
Van Erpecum K.J., Van Berge Henegouwen G.P., Stolk M.F.J.	295
A photometric assay for hydrogen peroxide production by polymorphonuclear leucocytes	
Negri M., Bellavite P., Lauciello C., Guzzo P., Zatti M.	305
Elevated serum β -glucuronidase activity in acquired immunodeficiency syndrome	
Saha A.K., Glew R.H., Kotler D.P., Omene J.A.	311

Vol. 200/1, 1991

Atherogenic risk factors in cerebrotendinous xanthomatosis	
Fujiyama J., Kuriyama M., Arima S., Shibata Y., Nagata K., Takenaga S., Tanaka H., Osame M.	1
Measurement of plasma free steroids by direct radioimmunoassay of ultrafiltrate in association with the monitoring of free components with [¹⁴ C]glucose	
Adachi K., Yasuda K., Fuwa Y., Goshima E., Yamakita N., Miura K.	13
Serum release of the creatine kinase tissue-specific isoforms MM3 and MB3 is simultaneous during myocardial reperfusion	
Christenson R.H., Magnus Ohman E., Vollmer R.T., Clemmensen P., Grande P., Wagner G.S.	23
A new sensitive method for determining endotoxin in whole blood	
Tamura H., Tanaka S., Obayashi T., Yoshida M., Kawai T.	35
A new mutant erythrocyte pyrimidine 5'-nucleotidase characterized by fast electrophoretic mobility in a Chinese boy with chronic hemolytic anemia	
Li J.-Y., Wan S.-D., Ma Z.-M., Zhao Y.-H., Zhou G.-P.	43
Letters to the Editor	
A universal eluent system for free amino acid analysis in biological fluids	
Pospisil R., Valik D.	49
Macro creatine kinase type 2 in a patient with prostatic carcinoma	
Arenas J., Bornstein B., Mayor J.J., Santos I., Leiva O., Martinez A.	53
Effects of dimethyl sulfoxide (DMSO) on carbon (CCl ₄)-induced hepatotoxicity in mice	
Achudume A.C.	57
A simplified analysis of fatty acids in serum phospholipids using Sep-Pak cartridges	
Ellin A., Van den Berg S., Strandvik B.	59
Measurement of total fasting-state serum bile acids: Comparison of a solid-phase bioluminescence enzymatic assay, a homogeneous fluorescence enzymatic assay and isotope dilution gas chromatography-mass spectrometry	
Rossi S.S., Angellotti M.A., Setchell K.D.R., Hofmann A.F.	63
Suppression/stimulation of chemiluminescence by human seminal plasma	
Haq A., Sheth K., Abdullatif M., Jabbar F.A.-A., Hamilton C., Al-Sedairy S.	67

Vol. 200/2-3, 1991

β -N-Acetylhexosaminidases in human cerebrospinal fluid and serum of patients with multiple sclerosis	
Datti A., Emiliani C., Capocchi G., Orlacchio A.	73
CA 15-3 and carcinoembryonic antigen in the clinical evaluation of breast cancer	
Dnistrian A.M., Schwartz M.K., Greenberg E.J., Smith C.A., Schwartz D.C.	81
A sensitive fluorescence assay for quantitation of fludarabine and metabolites in biological fluids	
Kemena A., Fernandez M., Bauman J., Keating M., Plunkett W.	95
Pyrophosphate inhibition of Proteus mirabilis-induced struvite crystallization in vitro	
McLean R.J.C., Downey J., Claphan L., Wilson J.W.L., Nickel J.C.	107

Mechanized assay of plasma prekallikrein by activation with <i>Pseudomonas aeruginosa</i> elastase and amidolysis of chromogenic substrate Shibuya Y., Yamamoto T., Tanaka H., Semba U., Kambara T., Okabe H.	119
Human aortic elastin from normal individuals and atherosclerotic patients: Lipid and cation contents; susceptibility to elastolysis Saulnier J.M., Hauck M., Fulop Jr. T., Wallach J.M.	129
Development of enzyme-linked immunosorbent assay for free human pro-colipase activation peptide (APGPR) Bowyer R.C., Jehanli A.M.T., Patel G., Hermon-Taylor J.	137
Biochemical properties of blood cells from histiocytosis X patients Pertino E., Marra E., Petragallo V.A., Quagliariello E., De Terlizzi M., Santostasi T., Ceci A.	153
Stability of platelet activating factor (PAF) in human saliva. Quantitation by radioimmunoassay Cooney S.J., Smal M.A., Baldo B.A.	161
Immunological assay of erythrocyte acetylcholinesterase Jones J.W., Whittaker M., Braven J.	175
Absence of increased urinary excretion of adenosine-deaminase-binding protein by patients with chronic renal tubular malfunction Flynn F.V., Lapsley M., Sansom P.A., Norden A.G.W.	183
Serum amyloid P component in chronic renal failure and dialysis Nelson S.R., Tennent G.A., Sethi D., Gower P.E., Ballardie F.W., Amatayakul-Chantler S., Pepys M.B.	191
Membrane properties of the sarcolemma and sarcoplasmic reticulum of pigs susceptible to malignant hyperthermia. Action of halothane Thomas M.A., Rock E., Viret J.	201
Increased elastase- α_1 -antitrypsin complex in fulminant hepatic failure: Relationship to bacterial infection and activation of coagulation Langley P.G., Hughes R.D., Rolando N., Williams R.	211
<i>Letters to the Editor</i>	
Human liver vasopressin receptors Howl J., Strain A.J., Wheatley M.	221
Usefulness of faecal water potassium/sodium ratios in the investigation of diarrhoea Duncan A., Robertson C., Russell R.I.	223
Impossibility to study crystalluria directly with Coulter Counter Hennequin C., Gaubil J., Bader C., Daudon M., Drueke T., Lacour B.	227
Glycohistones in diabetic human liver Jobst K., Lakatos A., Horvath A.	231

Vol. 201/1-2, 1991

Neopterin in clinical practice Muller M.M., Curtius H.-C., Herold M., Huber C.H.	1
Analysis of bismuth in serum and blood by electrothermal atomic absorption spectrometry using platinum as matrix modifier Slikkerveer A., Helmich R.B., Edelbroek P.M., Van der Voet G.B., De Wolff F.A.	17
Bone-alkaline phosphatase as indicator of bone formation Van Straalen J.P., Sanders E., Prummel M.F., Sanders G.T.B.	27
High-performance liquid chromatographic determination of bilirubin distribution on serum proteins Soltes L., Sebille B., Fugedi A.	35
Clinical analysis in intact erythrocytes using ^1H spin echo NMR Reglinski J., Smith W.E., Wilson R., Buchanan L.M., McKillop J.H., Thomson J.A., Brzeski M., Marabani M., Sturrock R.D.	45
Changes in fucose metabolism associated with heavy drinking and smoking: A preliminary report Thompson S., Matta K.L., Turner G.A.	59

Differential low density lipoprotein hydrated density distribution in female and male patients with familial hypercholesterolemia	
Tilly-Kiesi M.K., Tikkanen M.J.	65
Increased plasma cyclic guanosine monophosphate concentration in chronic obstructive pulmonary disease patients with increased circulating atrial natriuretic peptide	
Rota M., Brulot N., Rezgui N., Labrousse J., Allard D.	75
Reference values for free gamma-aminobutyric acid determined by ion-exchange chromatography and fluorescence detection in the cerebrospinal fluid of children	
Carchon H.A., Jaeken J., Jansen E., Eggermont E.	83
Pitfalls in the determination of diamine oxidase activity	
Smith C.H., Maghsoudloo M., Himuna K., Murphy G.M.	89
Enzymatic phenylalanine estimation for the management of patients with phenylketonuria	
Wendel U., Koppelkamm M., Hummel W.	95
HbA(1C) and serum fructosamine in diabetic patients: Relationship to age, clotting and fibrinolysis parameters and urinary microalbumin excretion	
Van Wersch J.W.J., Westerhuis L.W.J.J.M., Venekamp W.J.R.R.	99
Detection of an eosinophil derived neurotoxin in the urine of a patient with idiopathic chronic eosinophilic pneumonia	
Deviller P., Gruart V., Prin L., Tardy F., Bulle C., Capron M., Capron A., Cordier J.-F.	105
The common finding of covert hypothyroidism at initial clinical evaluation for hyperlipoproteinemia	
Glueck C.J., Lang J., Tracy T., Speirs J.	113
Diurnal and postural variation in serum α_1 microglobulin in normal individuals	
Itoh Y., Kawai T.	123
Hyperornithinemia, hyperammonemia and homocitrullinuria — A case report and study of ornithine metabolism using in vivo deuterium labelling	
Tsujino S., Suzuki T., Azuma T., Higa S., Sakoda S., Kishimoto S.	129
<i>Letters to the Editor</i>	
Nomenclature of hormones used as drugs for biochemical function tests	
Fuentes-Arderiu X.	135
Instability of fecal lysozyme	
Brouwer J., Smeekens F.	137
Serum beta-2-microglobulin levels and cognitive function in chronic dialysis patients	
Annunziata P., Biondi B., Aucone A.M., Poggi A., Moschini F.	139
Measurement of intact parathormone in hyperthyroidism	
Roiter I., Da Rin G., Galliazzo S., Legovini P., Di Virgilio R., Foscolo G., Breda F., Conte N.	143
Diagnosis of organic acidemia in developing countries	
Lin Y.Y., Wisniewski K.E., Inoue Y., Matsumoto I., Schmidt B.J., Raiman E., Cornejo V.	145
Amino-peptidases in the soluble fraction of human placenta	
Mizutani S., Nomura S., Kurauchi O., Kasugai M., Ino K., Goto S., Tomoda Y.	151

Vol. 201/3, 1991

Improved synthesis of [1- ¹⁴ C]acyl-sphingosine-galactose-3-sulfate (sulfatide) for diagnosis of metachromatic leukodystrophy: Usefulness of radioscanning	
Masson M., Li W.X., Fluharty A.L., Beaucourt J.P., Turpin J.C., Baumann N.	157
Some aspects of the use of dextran sulfate-cellulose columns for the treatment of familial hypercholesterolemia	
Erich D., Aubert I., Neel D., Scrobohaci M.L., Loreau J., Bussel A., Rouffy J., Dreux C., Goussault Y., Weintraub H.	169
A radiometric assay for aspartoacylase activity in human fibroblasts: Application for the diagnosis of Canavan's disease	
Barash V., Flhor D., Morag B., Boneh A., Elpeleg O.N., Gilon C.	175
Sensitive enzyme immunoassay for human 28 kDa calbinin-D	
Zhu Y., Takashi M., Miyake K., Kato K.	183

Brain natriuretic peptides in human plasma, spinal cord and cerebrospinal fluid Togashi K., Kameya T., Ando K., Marumo F., Kawakami M.	193
Anti-skeletal muscle and anti-acetylcholine receptor antibodies in patients with thymoma without Myasthenia gravis: Relation to the onset of Myasthenia gravis Ohta M., Itoh M., Hara H., Itoh N., Nishitani H., Hayashi K., Ohta K.	201
Identification of human ventricular myosin heavy chain fragments with monoclonal antibody 2F4 in human sera after myocardial necrosis Simeonova P.P., Kehayov I.R., Kyurkchiev S.D.	207
Immunonephelometric assay of myoglobin Massoubre C., Chivot L., Mainard F., Bridji B., Madec Y.	223
Determination by latex immunoassay of protein I in normal and pathological urine Bernard A., Lauwerys R., Noel A., Vandeleene B., Lambert A.	231
Examination of the role of oxygen free radicals in bronchial asthma in childhood Novak Z., Nemeth I., Gyurkovits K., Varga S.I., Matkovic B.	247

Vol. 202/1-2, 1991

Quantitation of urobilinogen in feces, urine, bile and serum by direct spectrophotometry of zinc complex Kotal P., Fevery J.	1
Investigation and validation of the affinity chromatography method for measuring glycated albumin in serum and urine Silver A.C., Lamb E., Cattell W.R., Dawney St. B.J.	11
A competitive radioimmunoassay for peptidoglycan monomer Vranesic B., Ljevakovic D., Tomasic J., Ladesic B.	23
In thiamine deficiency, activation of erythrocyte transketolase by thiamine in vivo exceeds activa- tion by cofactor in vitro Price J., Clague A.E., Kerr R.A., Nixon P.F.	39
Comparison of normal subjects and asthma patients with respect to age-related elastase-binding activity of α_2 -macroglobulin in plasma Kilroe-Smith T.A., Becker P.J., Gaillard M.C.	47
α_2 -Macroglobulin and generation of oxygen radicals by granulocytes: Potential role in prevention and treatment of reperfusion injury Donnelly P.K., Booth H., White M., Shenton B.K.	55
Microheterogeneity of α_1 -acid glycoprotein in healthy elderly subjects: Patterns obtained by crossed affino-immunoelectrophoresis Kawerk N., Succari-Aderschlag M., Foglietti M.-J.	65
Serum F-protein concentration following halothane or isoflurane anaesthesia Cooper C.M., Foster G.R., Cooper A.E., Kenny M., Olivera D.B.	73
Comparison of sampling systems for the collection of $^{14}\text{CO}_2$ in ^{14}C breath tests Peled Y., Gilat T.	83
Skeletal muscle cytochrome c in X-linked muscle dystrophies Nigro G., Comi L.I., Politano L., Limongelli F.M.	89
Platelet-activating factor acetylhydrolase in plasma lipoproteins of healthy men and women Satoh K., Imaizumi T.-A., Yoshida H., Kawamura Y., Takamatsu S., Takamatsu M., Mizuno S.	95
Carbon monoxide production from L-3,4-dihydroxyphenylalanine; a method for assessing the oxidant/antioxidant properties of drugs Cooper M.J., Engel R.R.	105
Plasma lactoferrin levels after bone marrow transplantation monitored by a two-site enzyme immunoassay Suzuki T., Takizawa-Mizuno M., Yazaki M., Wada Y., Asai K., Kato T.	111
<i>Letter to the Editor</i>	
Lactate dehydrogenase isoenzymes in serum of patients with preeclampsia/eclampsia complicated by the HELLP syndrome Beyer C.	119

IFCC Recommendation — Theory of reference values. Part 4. Control of analytical variation in the production, transfer and application of reference values Solberg H.E., Stamm D.	55
IFCC Recommendation — Recommendation on sampling, transport and storage for the determination of the concentration of ionized calcium in whole blood, plasma and serum Boink A.B.T.J., Buckley B.M., Christiansen T.F., Covington A.K., Maas A.H.J., Muller-Plathe O., Sachs Ch., Siggaard-Andersen O.	S13
Guidelines (1990) for selection of safe laboratory centrifuges and for their safe use. With general purpose appendices concerning centrifuge nomenclature, quantities and units, and calculation of centrifugal acceleration Uldall A., Damgaard P.T., Drachmann O., Jorgensen F., Kennedy D., Lauritzen M., Magnussen E., Rigg J.C., Voss P.	523

Vol. 202/3, 1991

Accumulation and impaired in vivo metabolism of di- and trihydroxycholestanic acid in two patients Wanders R.J.A., Casteels M., Mannaerts G.P., Van Roermund C.W.T., Schutgens R.B.H., Kozich V., Zeman J., Hyaneck J.	123
Body fat distribution and serum lipoproteins in relation to age and body weight Sonnichsen A.C., Richter W.O., Schwandt P.	133
Raised urinary excretion of inorganic pyrophosphate in asymptomatic members of a hypophosphatasia kindred Macfarlane J.D., Poorthuis B.J.H.M., Mulivor R.A., Caswell A.M.	141
Arylsulfatases A and B in EBV-transformed lymphoid cell lines: Studies on their molecular forms in cells from patients with inborn sulfatase deficiencies. Comparative diagnostic value of enzymatic assays Tempesta M.-C., Levade T., Salvayre R.	149
A rapid fluorometric enzyme immunoassay for the determination of neonatal TSH from blood spots Tuuminen T., Rakkolainen A.E.M., Welin M.-G.I., Weber T.H., Nylander P.L., Kapyaho K.I.	167
Sandwich enzyme immunoassay for serum integrins using monoclonal antibodies Katayama M., Kurome T., Yamamoto K., Uchida H., Hino F., Kato I.	179
A new immunochemical assay for biotin Thuy L.P., Sweetman L., Nyhan W.L.	191
Isolation of amniotic fluid proteins of non-maternal serum origin by negative immuno-affinity chromatography Sorensen S.	199
Effect of alcohol on δ -aminolevulinic acid dehydratase and porphyrin metabolism in man Sieg I., Doss M.O., Kandels H., Schneider J.	211
ELISA using monoclonal antibody to human serum arylesterase Kawai H., Yomoda S., Inoue Y.	219
A new enzymatic assay of urinary guanidinoacetic acid Shirokane Y., Nakajima M., Mizusawa K.	227
Role of iron in the photosensitization by uroporphyrin Menon I.A., Becker M.A.C., Haberman H.F.	237
Radiimmunoassay of growth hormone-releasing hormone (GHRH) with a polyclonal antibody against synthetic GHRH(1-29)-Gly ₄ -Cys-NH ₂ : Method and clinical studies Zhang C., Yamasaki R., Mitsuhashi S., Takahashi H., Bando H., Saito S.	243

Vol. 203/1, 1991

Isotransferrins and pregnancy: A study in the guinea pig Van Dijk J.P., Van Noort W.L., Kroos M.J., Starreveld J.S., Van Eijk H.G.	1
---	---

Marked variation in blood beta-hexosaminidase in Gaucher disease Natowicz M.R., Prencz E.M., Cajolet A.	17
Specific diagnosis of medium-chain acyl-CoA dehydrogenase (MCAD) deficiency in dried blood spots by a polymerase chain reaction (PCR) assay detecting a point-mutation (G985) in the MCAD gene Gregersen N., Blakemore A.I.F., Winter V., Andresen B., Kolvraa S., Bolund L., Curtis D., Engel P.C.	23
Interest of zinc determination in leucocyte fractions for the assessment of marginal zinc status Peretz A., Neve J., Jeghers O., Leclercq N., Praet J.-P., Vertongen F., Famaey J.-P.	35
Aminoterminal propeptide of type III procollagen in cerebrospinal fluid. Variation with age and in childhood leukemia Vainionpää L., Risteli L., Lanning M., Myllylä V., Risteli J.	47
Increased concentrations of endogenous 13-cis- and all-trans-retinoic acids in diffuse idiopathic skeletal hyperostosis, as demonstrated by HPLC Periquet B., Lambert W., Garcia J., Lecomte G., De Leenheer A.P., Mazieres B., Thouvenot J.P., Arlet J.	57
Flow injection system based on the sandwich technique for saving expensive reagents Araujo A.N., Lima J.L.F.C., Alonso-Chamarro J., Bartroli J., Poch M.	67
Synthesis of ¹³ C-labeled chenodeoxycholic, hyodeoxycholic, and ursodeoxycholic acids for the study of bile acid metabolism in liver disease Matern S., Marzschall H.-U., Schill A., Schumacher B., Lehnert W., Sjøvall J., Matern H.	77
<i>Letters to the Editor</i>	
Refinement of a specific assay for hydroxyproline in urine Sela B.-A., Doolman R.	91
Rapid quantitation of serum myoglobin by latex agglutination turbidimetry Abe J., Yamaguchi T., Isshiki T., Naka H., Kawada Y., Katayama Y.	95
A discrepancy between in vivo and in vitro lipoprotein lipase activity in a patient with Tangier disease Takizawa A., Komoda T., Hokari S., Sakagishi Y., Tanaka A.	97
Wheat germ agglutinin binding to serum 5'-nucleotidase and alkaline phosphatase Novo F.J., Tutor J.C.	101
Further data on an RIA kit for the direct measurement of free testosterone in plasma Rivera-Coll A., Navarro M.A.	103
Micromethod for the assay of glutamate: Glyoxylate aminotransferase and modifications of a micromethod for the assay of alanine: Glyoxylate aminotransferase. Implications for the prenatal diagnosis of Type I hyperoxaluria by fetal liver biopsy Toone J.R., Applegarth D.A.	105

Vol. 203/2-3, 1991

Effect of simvastatin on the apparent size of LDL particles in patients with type IIB hyperlipoproteinemia Zhao S.P., Hollaar L., Van 't Hooft F.M., Smelt A.H.M., Leuven J.A.G., Van Der Laarse A.	109
Energy metabolism during damaging contractile activity in isolated skeletal muscle: A ³¹ P-NMR study West-Jordan J.A., Martin P.A., Abraham R.J., Edwards R.H.T., Jackson M.J.	119
Abnormal serum immunoglobulin concentrations in patients with diabetes mellitus Rodriguez-Segade S., Camina M.F., Paz J.M., Del Rio R.	135
Phosphoribosylpyrophosphate synthetase in human erythrocytes: Assay and kinetic studies using high-performance liquid chromatography Sakuma R., Nishina T., Yamanaka H., Kamatani N., Nishioka K., Maeda M., Tsuji A.	143
Antibodies to dietary antigens in rheumatoid arthritis — Possible molecular mimicry mechanism Perez-Maceda B., Lopez-Bote J.P., Langa C., Bernabeu C.	153

Changes in α_1 -acid glycoprotein serum concentrations and glycoforms in the developing human fetus	
Seta N., Tissot B., Forestier F., Feger J., Daffos F., Durand G.	167
Determination of apolipoprotein E phenotypes from stored or postmortem serum samples	
Lehtimäki T.	177
Detection of the most common mutation of adenine phosphoribosyltransferase deficiency among Japanese by a non-radioactive method	
Kawaguchi R., Higashimoto H., Hikiji K., Hakoda M., Kamatani N.	183
Rapid automated quantitation of isoleucine, leucine, tyrosine and phenylalanine from dried blood filter paper specimens	
Qu Y., Miller J.B., Slocum R.H., Shapira E.	191
Urinary oligosaccharide excretion and severity of galactosialidosis and sialidosis	
Takahashi Y., Nakamura Y., Yamaguchi S., Orii T.	199
High-performance liquid chromatography of urinary oligosaccharides in the diagnosis of glycoprotein degradation disorders	
Hommes F.A., Varghese M.	211
Assay for type III collagenolytic activity in lung cancer tissue	
Kawamura M., Kato R., Kikuchi K., Kobayashi K., Ishihara T., Shibata T., Inayama S.	225
In vitro stability of human atrial natriuretic peptide (h-ANP)	
Artner-Dworzak E., Lindner H., Puschendorf B.	235
The glycoconjugate content of human faeces and ileostomy effluents	
Clamp J.R., Gough M.	243
Phospholipase A ₂ is associated with albumin in patients with acute pancreatitis	
Langton S.R., Dench J.	249
A re-evaluation of the lipid-bound sialic acid determination	
Vegh Z., Kremmer T., Boldizsar M., Gesztesi K.A., Szajani B.	259
Deoxycholic acid is not reconverted to cholic acid in humans — A study by isotope ratio mass spectrometry	
Fischer S., Beuers U., Berr F., Paumgartner G.	269
Assay of plasma glycine by HPLC with electrochemical detection in patients undergoing glycine irrigation during gynaecological surgery	
Sherwood R.A., Bayliss E.M., Chappatte O.	275
Urinary hydroxyproline and serum alkaline phosphatase in sickle cell disease	
Mohammed S.M., Suleiman S.A., Addae S.K., Annobil S.H., Adzaku F.K., Kadoum O.F., Richards J.T.	285
Rapid and sensitive discriminating determination of acetylcholinesterase activity in amniotic fluid with a choline sensor	
Morelis R.M., Coulet P.R., Simplot A., Boisson C., Guibaud G.	295
Serum levels of human alkaline phosphatase isozymes in relation to blood groups	
Domar U., Hirano K., Stigbrand T.	305
Glycosylation of α_1 -acid glycoprotein in relation to duration of disease in acute and chronic infection and inflammation	
Faasbender K., Zimmerli W., Kissling R., Sobieska M., Aeschlimann A., Kellner M., Müller W.	315
Effect of acute hyperinsulinemia on fatty acid composition of serum lipids in non-insulin-dependent diabetics and healthy men	
Pelikanova T., Kohout M., Base J., Stefka Z., Kovar J., Kazdova L., Valek J.	329
Azopigments from serum bilioprotein and accuracy of total bilirubin measurement	
Franzini C., Cattozzo G.	339
Lysosomal enzymes and severity of saphen varicosis	
Hayer M., Caporiccio A., Pages A., Caporiccio G., Toueri M.F., Magnan de Bornier B.	343
The identification of type 1 Gaucher disease patients, asymptomatic cases and carriers in The Netherlands using urine samples: An evaluation	
Aerts J.M.F.G., Sa Miranda M.C., Wanzeller De Lacerda L., Van Weely S., Donker-Koopman W., Brouwer-Kelder B., Jansen D.C., Van Leeuwen M., Schram A.W., Tsipara A., Tager J.M.	349

Elevated plasma histamine in sickle cell anaemia Enwonwu C.O., Lu M.	363
Prenatal diagnosis of isovaleric acidemia by fast atom bombardment and tandem mass spectrometry Shigematsu Y., Kikawa Y., Sudo M., Kanaoka H., Fujioka M., Dan M.	369
Serum gamma glutamyl transpeptidase in head and neck cancer Singh J., Sharma A., Yadav S.P.S., Lal H.	375
Variations in relative concentrations of variants of human α_1 -acid glycoprotein after acute-phase conditions Eap C.B., Fischer J.-F., Baumann P.	379
Identification of renal calculi by their Raman spectra Sudlow K., Woolf A.	387
Measurement of ornithine carbamyl transferase (OCT) in plasma by means of enzymatic determination of ammonia Peltenburg H.G., Janssen M.A., Soeters P.B., Flendrig J.G., Hermens W.T.	395
<i>Letters to the Editor</i>	
Isoelectric focusing of neutrophil alkaline phosphatase in trisomy 21 pregnancies Brisson-Lougarre A., Vergnes H., Grozdea J., Bourrouillou G., Colombies P.	403
A practical new assay for determining N-acetyl- β -D-glucosaminidase activity in urine Jung K., Priem F.	405
Biochemical properties of γ -glutamyl transpeptidase in human submandibular gland Sajjan A.R., Hinchigeri S.B., Datta K.S.	409
Improved method for the determination of the magnesium concentration of mononuclear blood cells Nozue T., Kodama T., Kobayashi A.	411
Level is not a synonym of concentration Fuentes-Arderiu X.	417
The effects of common matrices for assay standards on performance of 'ultra sensitive' immunometric assays for TSH. Report of a joint WHO/IFCC collaborative study Gaines-Das R.E., Brettschneider H., Bristow A.F.	S5

Vol. 204/1-3, 1991

Alterations of the glycan moiety of human α_1 -acid glycoprotein in late-term pregnancy Biou D., Bauvy C., N'Guyen H., Codogno P., Durand G., Aubery M.	1
Strenuous physical activity, aspirin and heat stress increase urinary dolichols: Evidence for lysosomal origin of urinary dolichols Roine R., Heinonen T., Salmela K., Heikkonen E., Suokas A., Luurila O.J., Koskinen P., Palo J., Salaspuro M.	13
Effectiveness of palladium matrix modification for the determination of thallium by graphite furnace atomic absorption spectrometry Yang Q., Smeyers-Verbeke J.	23
In vitro glycation of human apolipoprotein A(I) reduces its efficiency in lecithin:cholesterol acyl-transferase activation Gugliucci A., Stahl A.J.C.	37
Determination of protein-bound urinary gamma-carboxyglutamic acid in calcium nephrolithiasis Colette C., Benmbarek A., Boniface H., Astre C., Pares-Herbute N., Monnier L., Guitter J.	43
The adrenergic control of lipolysis: Development of tolerance to beta-2 receptor stimulation with terbutaline Kendall M.J., Hafner C.A., Hughes B., Florkowski C.	51
Predisposition to LDL oxidation during copper-catalyzed oxidative modification and its relation to α -cotopherol content in humans Cominacini L., Garbin U., Cenci B., Davoli A., Pasini C., Ratti E., Gaviraghi G., Lo Cascio V., Pastorino A.M.	57

Arylsulfatase A in serum from patients with cancer of various organs Laidler P., Kowalski D., Silberring J.	69
Gas chromatography method for the separation of amino acids enantiomers in plasma and urine. Application in a case of short bowel syndrome Ketling D., Wadman S.K., Spaapen L.J.M., Van Der Meer S.B., Duran M.	79
Urine thiocyanate:creatinine ratio as a reliable indicator of cigarette smoking Pre J., Vassy R.	87
Cellular and humoral responses in coeliac disease. 1. Wheat protein fractions Penttilä I.A., Devery J.M., Gibson C.E., LaBrooy J.T., Skerrett J.H.	95
Cellular and humoral responses in coeliac disease. 2. Protein extracts from different cereals Skerrett J.H., Devery J.M., Penttilä I.A., Labrooy J.T.	109
Influence of serum proteins on fructosamine concentration in multiple myeloma Montagna M.P., Laghi F., Cremona G., Zuppi C., Barbaresi G., Castellana M.L.	123
3-Hydroxy-3-methylglutaric aciduria: A possible pitfall in diagnosis Mills G.A., Hill M.A.W., Buchanan R., Corina D.L., Walker V.	131
Optimisation of non-denaturing polyacrylamide gel electrophoretic analysis of glyoxalase I phenotypes in clinical blood samples McLellan A.C., Thornalley P.J.	137
The nature of small bowel luminal fluid lactase Quak S.H., Brown G.A., Booth I.W., McNeish A.S.	145
Composition of very low density lipoproteins and in vitro effect of lipoprotein lipase Saheki S., Takahashi I., Murase M., Takeuchi N., Uchida K.	155
Fatty acid composition of hepatic triglycerides in Reye's syndrome: Implications for hepatic desaturase abnormalities Kang E.S., Johnson S.B., Holman R.T.	167
Purification and structural study of two albumin variants in an Irish population Sakamoto Y., Davis E., Madison J., Watkins S., McLaughlin H., Leahy D.T., Putnam F.W.	179
Capillary plasma lipid profiles Mace R., May B., McCray C., Tracy T., Speirs J., Glueck C.J.	189
Comparison of nonparametric recursive partitioning to parametric discriminant analyses in labo- ratory differentiation of hypercalcemia Lacher D.A.	199
Ethinyl estradiol and norethindrone radioimmunoassay following Sephadex LH-20 column chro- matography Zacur H.A., Linkins S., Chang V., Smith B., Kimball A.W., Burkman R.	209
Assay of inulin in biological fluids by high-performance liquid chromatography with pulsed amperometric detection Ruo T.I., Wang Z., Dordal M.S., Atkinson Jr. A.J.	217
Creatinine deiminase (EC 3.5.4.21) from bacterium BN11: Purification, properties and applicabili- ty in a serum/urine creatinine assay Gottschalk E.M., Hippe H., Patzke F.	223
Betaine:homocysteine methyltransferase — A new assay for the liver enzyme and its absence from human skin fibroblasts and peripheral blood lymphocytes Wang J., Dudman N.P.B., Lynch J., Wilcken D.E.L.	239
Platelet aggregation and fatty acid composition of platelets in type 1 diabetes mellitus Myrup B., Bregengaard C., Petersen L.R., Winther K.	251
Carnitine therapy and metabolism in the disorders of propionyl-CoA metabolism studies using ¹ H-NMR spectroscopy Davies S.E.C., Iles R.A., Stacey T.E., De Sousa C., Chalmers R.A.	263
Increased susceptibility to peroxide-induced haemolysis with normal vitamin E concentrations in cystic fibrosis James D.R., Alfaham M., Goodchild M.C.	279
Serum Lp(a) concentrations are unaffected by treatment with the HMG-CoA reductase inhibitor Pravastatin: Results of a 2-year investigation Fieseler H.-G., Armstrong V.W., Wieland E., Thierry J., Schutz E., Walli A.K., Seidel D.	291

Plasma bromine concentrations and lipid profiles

Corrigan F.M., Skinner E.R., MacIntyre F., Ijomah G., Holliday J., Horrobin D.F., Ward N.I. 301

Letter to the Editor

Isoelectric focusing of neutrophil alkaline phosphatase in trisomy 21 pregnancies

Brisson-Lougarre A., Vergnes H., Grozdea J., Bourrouillou G., Colombies P. 305

Vol. 205/1-2, 1992

Generation of analytic plasma lipoprotein profiles using two prepacked superose 6B columns

Gerdes L.U., Gerdes C., Klausen I.C., Faergeman O. 1

High-performance liquid chromatographic determination of free polyamines in human seminal plasma

Oefner P.J., Wongyai S., Bonn G. 11

Discriminant analysis of urinary calculi by near-infrared reflectance spectroscopy

Peuchant E., Heches X., Sess D., Clerc M. 19

Tubular secretion of Tamm-Horsfall protein in type 1 (insulin-dependent) diabetes mellitus using a simplified enzyme linked immunoassay

Torffvit O., Agardh C.-D., Kjellson B., Wieslander J. 31

Influences of diet and surgical trauma on serum alkaline DNase activity levels

Avall Lundqvist E.H., Sjovalk K., Eneroth P.H.E. 43

Limitations of the triolein breast test

Duncan A., Cameron A., Stewart M.J., Russell R.I. 51

An enzyme-linked immunosorbent assay for heparin cofactor II (HCII). Application to the measurement of HCII in clinical materials

Toulon P., Costa J.M., Amiral J. 65

Selenium and glutathione peroxidase variations induced by polyunsaturated fatty acids oral supplementation in humans

Bellisola G., Galassini S., Moschini G., Poli G., Perona G., Guidi G. 75

Characterization of 6-hexadecanoylamino-4-methylumbelliferyl- β -D-galactopyranoside as fluorogenic substrate of galactocerebrosidase for the diagnosis of Krabbe disease

Wiederschain G., Raghavan S., Kolodny E. 87

Effects of unsaturated and saturated dietary plant sterols on their serum contents

Vanhanen H.T., Miettinen T.A. 97

Homocarnosinosis patients and great apes have a serum protein that cross-reacts with human serum carnosinase

Jackson M.C., Lenney J.F. 109

Mechanized assay of serum cholinesterase by specific colorimetric detection of released acid

Takeuchi T., Kabasawa Y., Horikawa R., Tanimura T. 117

A coupled assay detecting defects in fibroblast isoleucine degradation distal to enoyl-CoA hydratase: Application to 3-oxothiolase deficiency

Gibson K.M., Lee C.F., Kamali V., Sovik O. 127

Separation of plasma lipoproteins with a tabletop ultracentrifuge

Sykes E., Meany M., Schulz V., Kessel D. 137

Letters to the Editor

More on monoamine oxidase activity in whole blood

Konings C.H., Kuiper M.A., Wolters E.C. 145

Solving one of the problems of in vitro cytokine production and endotoxin measurement in blood

Redl H., Bahrami S., Leichtfried G., Schlag G. 149

Intestinal permeability: Update on the enzymatic assay of mannitol

Lunn P.G., Northrop-Clewes C.A. 151

The precipitation of histone-proteins with trichloroacetic acid

Lakatos A., Jobst K. 153

Erratum: Purification and structural study of two albumin variants in an Irish population (*Clinica Chimica Acta*, 204 (1991) 179-188)

Sakamoto Y., Davis E., Madison J., Watkins S., McLaughlin H., Leahy D.T., Putnam F.W. 155

Vol. 205/3, 1992

Cholinesterases exhibiting aryl acylamidase activity in human amniotic fluid Jayanthi L.D., Balasubramanian N., Balasubramanian A.S.	157
Activity of cholesterol in human gallbladder bile in relation to nucleation of cholesterol monohydrate crystals Noshiro H., Chijiwa K., Hirota I.	167
Increased serum concentration of type IV collagen peptide and type III collagen peptide in hyperthyroidism Inui T., Ochi Y., Chen W., Nakajima Y., Kajita Y.	181
The determination of asialoglycoforms of serum glycoproteins by lectin blotting with Ricinus communis agglutinin Lundy F.T., Wisdom G.B.	187
Clearance of purified human liver γ -glutamyltransferase after intravenous injection in the rat Huseby N.-E., Kindberg G.M., Grostad M., Berg T.	197
Increase in plasma total and lipoprotein cholesterol during incubation of whole blood samples at 37°C — Influence of LCAT inhibitors Ruhling K., Lang A., Holtz H., Winkler L., Schlag B., Till U.	205
A competitive ELISA for lipoprotein(a) Yeo K.H.J., Walmsley T.A., Owen M.C., George P.M.	213
Serum phospholipase A ₂ and lysolecithin changes following myocardial infarction Langton S.R., Jarnicki A.	223
Direct gas chromatographic assay of urinary medium-chain fatty acylcarnitines by their thermal decomposition Farquharson J., Jamieson E.C., Muir J., Cockburn F., Logan R.W.	233
<i>Letters to the Editor</i>	
The influence of antioxidants on the oxidative stress of red blood cells Varga S.I., Novak Z., Pataki L., Patocsai M., Matkovics B.	241
Competitive enzyme inhibition immunoassay of apolipoprotein B based on monoclonal antibody Rojas A., Zayas M., Falero G., Rodriguez I.	245
Use of homocysteine to stabilise ascorbic acid, or to reduced dehydroascorbic acid, during HPLC separation of large volumes of tissue extracts Bates C.J.	249
Quantities and units for electrophoresis in the clinical laboratory Ferard G.	S17
Quantities and units for centrifugation in the clinical laboratory Lauritzen M.	S5

Vol. 206/1-2, 1992

International Week of Laboratory Science — Rome, Italy, November 12-17, 1990: Introduction Arvan D.A., Burlina A.	1
Tumor cell heterogeneity: An overview Arvan D.A.	3
Phenotypic heterogeneity in acute leukemia Ryan D.H.	9
Drug-transporter proteins in clinical multidrug resistance Scheper R.J., Broxterman H.J., Scheffer G.L., Meijer C.J.L.M., Pinedo H.M.	25
Evaluation of tumor cell heterogeneity based on response to chemotherapy Asselin B.L.	33
Enzymes as agents for the treatment of disease Goldberg D.M.	45
Enzymes as prognostic markers and therapeutic indicators in patients with cancer Schwartz M.K.	77

Application of clinical laboratory measurements to issues of environmental health	
Rej R., Silkworth J.B., DeCaprio A.P.	83
Glutathione, protein sulphydryls and cysteine proteases in gastric mucosal injury and protection	
Szabo S., Nagy L., Plebani M.	95
Measurement of trypsin in endoscopic gastroduodenal biopsies: Distribution and relationship with ulcer disease	
Plebani M., Di Mario F., Battistel M., Basso D., Mantovani G., Giacomini A., Burlina A.	107
Magnetic resonance spectroscopy of the brain	
Prichard J.W.	115
Carbon-13 magnetic resonance in biology and medicine	
Seelig J., Burlina A.P.	125
Quantitative analysis of rat synaptosomes and cerebrum using high-resolution ¹ H magnetic resonance spectroscopy	
Petroff O.A.C., Burlina A.P., Black J., Prichard J.W.	137
ATP-stimulated glutamate-dependent calcium uptake by rat synaptosomes	
Galzigna L., Bianchi M., Battistin T., Scarpa M., Rigo A.	147

Vol. 206/3, 1992

A sequential and simple determination of zinc, copper and aluminium in blood samples by inductively coupled plasma atomic emission spectrometry	
Chappuis P., Poupon J., Rousselet F.	155
Characteristics of the interactions between acid glycosaminoglycans and protein in normal human plasma as revealed by the behaviour of the protein-polysaccharide complexes in ultrafiltration and chromatographic procedures	
Calatroni A., Vinci R., Ferlazzo A.M.	167
Relationship of urinary pseudouridine and 1-methyladenosine to activity of leukemia and lymphoma	
Itoh K., Konno T., Sasaki T., Ishiwata S., Ishida N., Misugaki M.	181
Creatine kinase MB isoforms for early diagnosis and monitoring of acute myocardial infarction	
Kanemitsu F., Okigaki T.	191
Organic acids in post-mortem cerebrospinal fluid	
Coude M., Kamoun P.	201
Effects of storage, iron and time of day on indices of lipid peroxidation in plasma from healthy volunteers	
Duthie G.G., Morrice P.C., Ventresca P.G., McLay J.S.	207
Determination of fetal hemoglobin by cation exchange liquid chromatography	
Koskinen L., Sillanauke P.	215
Serum α_1 -antitrypsin in patients with hepatocellular carcinoma	
Lee H.B., Yoo O.J., Ham J.S., Lee M.H.	225
β -N-Acetylhexosaminidase in the urine, kidney and serum of bromobenzene-intoxicated mice	
Tassi C., Beccari T., Casini A., Orlicchio A.	231
Letters to the Editor	
Use of tox-lab thin layer chromatography in plasma very long chain fatty acid analysis	
Smith N.B., Brooks K.E.	241
Reference intervals for α_1 -microglobulin in urine	
Jung K., Pergande M., Schreiber G., Thierfelder W.	245
Improved isolation of mononuclear cells from stored blood	
Ford T.C., Rickwood D.	249

Vol. 207/1-2, 1992

Assay of γ -glutamyltransferase with amino acid dehydrogenases from <i>Bacillus stearothermophilus</i> as auxiliary enzymes	
Kondo H., Hashimoto M., Nagata K., Tomita K., Tsubota H.	1

Enzyme-linked immunosorbent assay for serum pseudocholinesterase	
Itoh Y., Fukumachi I., Hashimoto H., Hachisu T., Kawai T.	11
A specific method for determination of peroxisomal β -oxidation activity in cultured human skin fibroblasts using a specific substrate, C ₆ : A possible application for screening of peroxisomal disorders	
Suzuki H., Yamada J., Watanabe T., Suga T.	19
Serum analysis for potassium ions using a fibre optic sensor	
Pereiro Garcia R., Alava Moreno F., Diaz Garcia M.E., Sanz-Medel A., Narayanaswamy R.	31
Aluminum-binding serum proteins: Desferrioxamine alters serum aluminum speciation	
Favarato M., Mizzen C.A., Sutherland M.K., Krishnan B., Kruck T.P.A., Crapper McLachlan D.R.	41
Insulin binding to erythrocytes of nonpregnant women: A reevaluation, underlining the importance of body weights even in nonobese subjects	
Desoye G., Schmon B., Gmoser G., Friedl H., Urdl W., Weiss P.A.M.	57
An indirect sandwich ELISA for LP(a) in serum and dried blood spots	
Wang X.L., Wilcken D.E.L., Dudman N.P.B.	73
Distribution of SP ₁ immunoreactivity among different plasma proteins: Real molecular heterogeneity or adsorption of SP ₁ -beta to other plasma proteins?	
Pala A., Di Ruzza A., Rossetto G., Moro M., Strom R., Padula F., Cosmi E.V.	87
Influence of triglycerides and urate on methods for determination of fructosamine	
Kallner A.	99
Comparison of soluble aminopeptidases in human cerebral cortex, skeletal muscle and kidney tissues	
Mantle D.	107
The effect of blood sample aging and food consumption on plasma total homocysteine levels	
Ubbink J.B., Hayward Vermaak W.J., Van der Merwe A., Becker P.J.	119
A case of Hashimoto's thyroiditis with antibodies to thyroid hormone analogues	
Sakata S., Maekawa H., Ogawa T., Sarui H., Matsuda M., Endo J., Wada S., Nagai K.	129
Selenium, glutathione peroxidase (GSH-Px) and lipid peroxidation products before and after selenium supplementation	
Wilke B.C., Vidailhet M., Favier A., Guillemin C., Ducros V., Arnaud J., Richard M.J.	137
Critical evaluation of the serum kappa/lambda light-chain ratio in the detection of M proteins	
Laine S.T., Soppi E.T., Morsky P.J.	143
Letter to the Editor	
3-Methylglutaconic aciduria in two adults	
Kuhara T., Matsumoto I., Saiki K., Takabayashi H., Kuwabara S.	151

Vol. 207/3, 1992

Increased prostaglandin PGE ₂ and 6-keto-PGF(1 α) immunoactivity in incubates of gastric mucosa and forestomach of nephrectomized rats	
Weiler H., Esser D., Weiler C., Gerok W.	161
Routine use of hair root or buccal swab specimens for PCR analysis: Advantages over using blood	
Thomson D.M., Brown N.N., Clague A.E.	169
Determination of enterokinase activity by measuring the disappearance of trypsinogen	
Carchon H.A., Jansen E., Eggermont E.	175
The measurement of carnitine and acyl-carnitines: Application to the investigation of patients with suspected inherited disorders of mitochondrial fatty acid oxidation	
Bhuiyan A.K.M.J., Jackson S., Turnbull D.M., Aynsley-Green A., Leonard J.V., Bartlett K.	185
A flow-through system for the determination of salicylate in blood	
Bertocchi P., D'Ottavio D., Evangelisti M.E., Mascini M., Palleschi G.	205
A simple, sensitive technique for classification of apolipoprotein(a) isoforms by sodium dodecyl sulphate-polyacrylamide gel electrophoresis	
Farrer M., Game F.L., Adams P.C., Laker M.F., Alberti K.G.M.M.	215
Purification and characterisation of antigenic gliadins in coeliac disease	
Sjostrom H., Friis S.U., Noren O., Anthonsen D.	227

Human urinary protein I: Evidence for identity with the Clara cell protein and occurrence in respiratory tract and urogenital secretions Bernard A., Roels H., Lauwerys R., Witters R., Gielens C., Soumillion A., Van Damme J., De Ley M.	239
Need for both angiotensin converting enzyme and angiotensinase specific inhibition for a reliable assessment of angiotensin II levels in human plasma Boddi M., Coppo M., Rostagno C., Gensini G.F., Neri Serneri G.G.	251
<i>Letters to the Editor</i>	
Within-subject biological variation of pituitary-ovarian axis hormones and desirable imprecision Fuentes-Arderiu X., Albert P.	257
Expression of CuZn-superoxide dismutase and glutathione peroxidase in erythrocytes from diabetic and non-diabetic subjects Strange R.C., Jones P., Bicknell J., Scarpello J.	261
Structure and genetic engineering of antigens and antibodies: Applications in immunoassays Vihko P., Wagener C.	S5
Testing urine for drugs Catlin D., Cowan D., Donike M., Fraisse D., Oftebro H., Rendic S.	S13

Vol. 208/1-2, 1992

Possible role for chondroitin sulfate in urolithiasis: In vivo studies in an experimental model Michelacci Y.M., Boim M.A., Bergamaschi C.T., Rovigatti R.M., Schor N.	1
Serum immunoreactive β -glucuronidase determined by an enzyme-linked immunosorbent assay in patients with hepatic diseases Ohta H., Ono M., Sekiya C., Namiki M.	9
Procedure for development of an enzyme-linked immunosorbent assay. Development of an assay for human apolipoprotein A-I Gaver A., Weedy N.K., Maldonado B.R., Huang S., Wong L.	23
^{13}C and ^{31}P NMR studies of glucose and 2-deoxyglucose metabolism in normal and enzyme-deficient human erythrocytes Ferretti A., Bozzi A., Di Vito M., Podo F., Strom R.	39
Optimized determination of malondialdehyde in plasma lipid extracts using 1,3-diethyl-2-thiobarbituric acid: Influence of detection method and relations with lipids and fatty acids in plasma from healthy adults Hoving E.B., Laing C., Rutgers H.M., Teggeler M., Van Doormaal J.J., Muskiet F.A.J.	63
The glycosaminoglycan composition of the lung with acute and chronic pathology and in senescence Grundboeck-Jusco J., Binette J.P., Kimura A., Talarico L., Kaelin R., Schmid K.	77
Determination of (S)- and (R)-2-oxo-3-methylvaleric acid in plasma of patients with maple syrup urine disease Wendel U., Even G., Langenbeck U., Schadowaldt P., Hummel W.	85
An immunoreactive xanthine oxidase protein-possessing xanthinuria and her family Yamamoto T., Moriwaki Y., Suda M., Takahashi S., Hada T., Nanahoshi M., Agbedana E.O., Higashino K.	93
A receptor assay for serum low density lipoprotein-apolipoprotein B-100: Preliminary report Pham Q., Soldin S.J., Rifai N.	99
Amniotic fluid granulocyte colony-stimulating factor in preterm and term labor Saito S., Kato Y., Ishihara Y., Ichijo M.	105
Abnormal lysosomal cathepsin activities in leukocytes and cultured skin fibroblasts in late infantile, but not in juvenile neuronal ceroid-lipofuscinosis (Batten disease) Bennett M.J., Chern L., Carpenter K.H., Sladky J.T.	111
Liquid chromatography determination of α -tocopherol in erythrocytes Sierra C., Pastor M.C., De Ramon M.	119

Letters to the Editor

Effect of anaesthesia and acute intestinal ischemia on serum β -N-acetylhexosaminidase activity in rabbit as biological model	
Risueno J.C., Gomez-Alonso A., Garcia-Criado F.J., Garcia-Sastre A., Corral J., Cabezas J.A., Villar E.	127
Fetal and amniotic α_1 -acid glycoprotein	
Johnson A.M.	133
A method for quantitative collection of stool and urine for nitrogen balance studies in infants	
Kandil H.M., Switzer B.R., Heizer W.D.	135
Rapid and sensitive colloidal silver staining on cellulose acetate membranes	
Kovarik A., Ondrkalova M., Prachar J., Stofko J.	137
The effect of reducing and non-reducing sugars on the bichinchoninic acid reaction for protein determination	
Milton J.D., Mullen P.J.	141
Erratum: The influence of antioxidants on the oxidative stress of red blood cells (<i>Clinica Chimica Acta</i> (1992) 205 (241-244))	
Varga S.I., Novak Z., Pataki L., Patocskaï M., Matkovic B.	147

Vol. 208/3, 1992

N-glycosylation of serum proteins in disease and its investigation using lectins	
Turner G.A.	149
Prenatal diagnosis of Niemann-Pick disease type C	
De Winter J.M., Janse H.C., Van Diggelen O.P., Loss F.J., Beemer F.A., Kleijer W.J.	173
High-performance liquid chromatographic microassay for L-alanine:glyoxylate aminotransferase activity in human liver	
Petrarulo M., Pellegrino S., Marangella M., Cosseddu D., Linari F.	183
Serum triglyceride and insulin levels are associated with erythrocyte sodium-lithium counter-transport activity in normoglycaemic individuals	
Winocour P.H., Thomas T.H., Brown L., Laker M.F., Wilkinson R., Alberti K.G.M.M.	193
Liver transplantation modifies serum cholestanol, cholesterol precursor and plant sterol levels	
Nikkila K., Hockerstedt K., Miettinen T.A.	205
A simple electrophoretic method for separating elevated liver and bone alkaline phosphatase isoenzymes in plasma after neuraminidase treatment	
Chamberlain B.R., Buttery J.E., Pannall P.R.	219
2,3-Butanediol: A potential interfering substance in the assay of ethylene glycol by an enzymatic method	
Nilsson L., Jones A.W.	225
<i>Letters to the Editor</i>	
Oligosaccharides in (galacto)sialidosis urine	
Van Pelt J.	231
Enzyme immunoassay for serum cortisol	
Candito M., Krebs B., Chambon P.	233
Effect of long-term storage on human plasma osteocalcin level	
Tarallo P., Balland M., Henny J.	237

Vol. 209/1-2, 1992

Decrease of pyruvate dehydrogenase phosphatase activity in patients with congenital lactic acidemia	
Ito M., Kobashi H., Naito E., Saijo T., Takeda E., Mahbubul Huq A.H.M., Kuroda Y.	1
Clinical test of renal guanidinoacetic acid metabolism by oral citrulline and creatine loading	
Kadono K., Yamaguchi T., Tetsutani T., Yasunaga K.	9

Plasmodium falciparum induced perturbations of the erythrocyte antioxidant system Dubey M.K., Dubey M.L., Ganguly N.K., Mahajan R.C.	19
Characterization of the carbohydrate moiety of human γ -glutamyltransferase using lectin-blotting and glycosidase treatment Evjen G., Huseby N.-E.	27
Use of an azo derivative immobilized on an anionic resin for the determination of aluminium in water and dialysis fluids Hernandez Torres O., Jimenez Moreno F., Jimenez Abizanda A., Arias Leon J.J.	35
Investigation of urea cycle enzyme disorders by ^1H -NMR spectroscopy Burns S.P., Woolf D.A., Leonard J.V., Iles R.A.	47
Diagnosis of medium chain acyl CoA dehydrogenase deficiency by measurement of cis-4-decenoic acid in dried blood spots Heales S.J.R., Leonard J.V.	61
Increased urinary excretion of coproporphyrin isomers after transcatheter arterial embolization of hepatocellular carcinoma Horie Y., Yamada S., Kawasaki H.	67
Diagnostic value of adenosine deaminase and lysozyme in tuberculous pleurisy San Jose E., Valdes L., Sarandeses A., Alvarez D., Chomon B.	73
Evaluation of solubilizing agents for 25-hydroxyvitamin D ₃ immunoassays Kobayashi N., Ueda K., Shimada K.	83
Raised monophosphatase activity in schizophrenic patients Zilberman-Kaufman M., Agam G., Moscovits L., Livne A.A., Belmaker R.H.	89
Increased superoxide dismutase activity in erythrocytes of children with pulmonary hypertension Soave M.C., Moulsmas M., Chevalier P., Pillot R., Guidollet J.	95
<i>Letters to the Editor</i>	
Screening of glycerol kinase deficiency in patients affected by Duchenne and Becker muscular dystrophy Schreiber R., Passos-Bueno M.R., Rapaport D., De Cassia Pavanetto R., Vainzof M., Fernandes M.E.S., Zatz M.	103
A colorimetric determination of 5-hydroxyindole acetic acid in urine Mukerjee H., Pincus M.R.	105
<i>9th European Congress of Clinical Chemistry</i>	
Enzymes as reporters of pathophysiological events: Introduction Moss D.W., Naskalski J.	107
Enzyme induction by drugs and toxins Batt A.-M., Siest G., Magdalou J., Galteau M.-M.	109
Alkaline phosphatase as a reporter of cancerous transformation Millan J.L.	123
Changes in enzyme expression related to differentiation and regulatory factors: The acid phosphatase of osteoclasts and other macrophages Moss D.W.	131

Vol. 209/3, 1992

Alterations of purine metabolism in mononuclear cell populations of first degree relatives of insulin-dependent diabetic individuals with disturbed glucose tolerance Schafer J.H., Jira T.B., Kohler E., Michaelis D., Rjasanowski I.M.	141
Melatonin and 6-sulfatoxymelatonin circadian rhythms in serum and urine of primary prostate cancer patients: Evidence for reduced pineal activity and relevance of urinary determinations Bartsch C., Bartsch H., Schmidt A., Ilg S., Bichler K.-H., Fluchter S.-H.	153
Lipid and apolipoprotein changes after orthotopic liver transplantation for end-stage liver diseases Malmendier C.L., Lontie J.-F., Mathe D., Adam R., Bismuth H.	169
Relationships between androgen and estrogen sulfates in breast cyst fluid Nahoul K., Kottler M.-L.	179
Increased membrane activity of glyceraldehyde 3-phosphate dehydrogenase in erythrocytes of patients with homozygous sickle cell anaemia Mohamed A.O., Ronquist G., Al Bayoumi R.	189

An ELISA method to measure human myoglobin in urine Hodson A.W., Skillen A.W., Argent N.B.	197
Comparison of three techniques for the determination of protein content in human tears Meijer F., Van Haeringen N.J.	209
<i>Letter to the Editor</i>	
Consequences of applying parametric methods to non-gaussian distributions in the validation of analytical results Machuca I., Cortes J.D., Fuentes-Arderiu X.	215
Diagnostic applications of repetitive DNA sequences Epplen J.T.	S5
Genes in the diagnosis of malignant disease Wagner C.	S15

Vol. 210/1-2, 1992

Finally! Automated microanalysis of ammonia in arterial blood Conn H.O.	1
Arginase, a new marker of mammary carcinoma Straus B., Cepelak I., Festa G.	5
Cellulose acetate electrophoretic determination of bone alkaline phosphatase activity in healthy subjects and diabetic patients with and without osteopenia Nishikawa Y., Kanda T., Yoshihara H., Fukumoto K., Uematsu I.	13
Characterization of dipeptidyl peptidase IV (CD26) from human lymphocytes De Meester I., Vanhoof G., Hendriks D., Demuth H.-U., Yaron A., Scharpe S.	23
Differences in the enzymatic nature and the sugar-chain structure of γ -glutamyl transferase between normal and carcinomatous human kidney and prostate Arai K., Yoshida K.-I., Komoda T., Kobayashi N., Sakagishi Y.	35
Comparison of urinary excretion of four lysosomal hydrolases in healthy elderly and young adults VanderJagt D.J., Steinberg B.R., Glew R.H.	47
Identification of urinary metabolites of (\pm)-2-(p-isobutylphenyl)propionic acid (Ibuprofen) by routine organic acid screening Bennett M.J., Sherwood W.G., Bhala A., Hale D.E.	55
Intestinal type alkaline phosphatase hyperphosphatasemia associated with liver cirrhosis Saheki S., Ochi M., Horiuchi T., Sakagishi Y., Fujimori-Arai Y., Koyama I., Komoda T., Takeuchi N.	63
Purification of electron transfer flavoprotein from pig liver mitochondria and its application to the diagnosis of deficiencies of acyl-CoA dehydrogenases in human fibroblasts Bertrand C., Dumoulin R., Divry P., Mathieu M., Vianey-Saban C.	75
Interaction of oxidized low density lipoproteins with both apo B,E and scavenger receptors. A model for its production in vitro Vilella E., Joven J., Bargallo T., Turner P.R., Masana L.	93
Concentration of serum laminin and type IV collagen in liver diseases assayed by a sandwich enzyme-immunoassay using monoclonal antibodies Yokoya Y., Iwata K., Muragaki Y., Shiota C., Morimoto Y., Inoue M., Itoh H., Nishioka S., Ooshima A.	109
A hyaluronan-loading test applied to patients with liver and joint diseases Lindqvist U., Groth T., Loof L., Hellsing K.	119
Pregnancy-associated α_2 -glycoprotein in children with acute lymphocytic leukemia, Hodgkin's disease and non-Hodgkin's lymphomas Zalazar F.E., Chiabrando G.A., De Aldao N.A., Ojeda F., Vides M.A., Aldao M.A.J.	133
PAF acetylhydrolase activities in human systemic lupus erythematosus and lupus-prone mice Matsuzaki M., Ishiguro Y., Tetsumoto T., Tsukada Y., Kayahara H., Tadasa K.	139
CA 19-9 in non-neoplastic liver diseases. A clinical and laboratory study Collazos J., Genolla J., Ruibal A.	145

Letter to the Editor

- A rapid method for blood ammonia determination using the new blood ammonia checker (BAC) II
Huizenga J.R., Tangerman A., Gips C.H. 153

Vol. 210/3, 1992

- The role of citrate complexes in preventing urolithiasis
Ashby R.A., Sleet R.J. 157
- Immunochemiluminometric assays (ICMA) specific for growth hormone releasing hormone 1-44
NH₂ and 1-40 OH
Page M.D., Edwards C.A., Brown R., Weeks I., Woodhead J.S., Scanlon M.F. 167
- The inter-relationships between albuminuria, plasma albumin concentration and indices of
glycaemic control in non-insulin-dependent diabetes mellitus
Chan J.C.N., Yeung V.T.F., Cheung C.K., Swaminathan R., Cockram C.S. 179
- IL6 and acute phase plasma proteins in peritoneal fluid of women with endometriosis
Boutten A., Dehoux M., Edelman P., Seta N., Menard A., Madelenat P., Durand G. 187
- Development and validation of a robust specific enzyme mediated assay for phenylalanine in
serum
Campbell R.S., Brearley G.M., Varsani H., Morris H.C., Milligan T.P., Hall S.K., Hammond
P.M., Price C.P. 197
- Evidence that the presence of arginine can lead to overestimation of glucagon levels measured by
radioimmunoassay
Lacey R.J., Scarpello J.H.B., Morgan N.G. 211
- Evaluation of rat intestinal lactase in vivo with 4-galactosylxylose
Aragon J.J., Fernandez-Mayoralas A., Jimenez-Barbero J., Martin-Lomas M., Rivera-Sagredo
A., Villanueva D. 221
- Letters to the Editor*
- Impaired synthesis of taurine in a patient with tyrosinemia type I during the oral L-methionine
loading test
Valik D., Klusacek D., Pintera J. 227
- Lipoprotein(a) and the acute phase response
Craig W.Y., Ledue T.B. 231
- Activity of cathepsin H,B and metalloproteinase in the serum of patients with acute myocardial
infarction
Laszlo A., Sohar I., Sagi I., Kovacs J., Kovacs A. 233
- Disease-coupled specificity in the glycan structures of orosomucoid
Treuhait M.J., Halsall H.B. 237

Vol. 211/1-2, 1992

- Diagnosis of HIV infection
Schochetman G. 1
- Differential isoform profiles of α_2 -macroglobulin from plasma of patients with chronic-
progressive or relapsing-remitting multiple sclerosis
Back S.A., Alhadeff J.A. 27
- High levels of nervous system-specific proteins in cerebrospinal fluid in patients with early stage
Creutzfeldt-Jakob disease
Jimi T., Wakayama Y., Shibuya S., Nakata H., Tomaru T., Takahashi Y., Kosaka K., Asano
T., Kato K. 37
- Renin assay using a fluorogenic substrate and high performance liquid chromatography
Nakamura-Imajo N., Satomura S., Matsuura S., Murakami K. 47
- A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 3 (stromelysin-1)
using monoclonal antibodies
Obata K., Iwata K., Okada Y., Kohrin Y., Ohuchi E., Yoshida S., Shinmei M., Hayakawa T. 59

Isolation of acylcarnitines from urine: A comparison of methods and application to long-chain acyl-CoA dehydrogenase deficiency	
Morrow R.J., Rose M.E.	73
Glycation of human tissue and serum creatine kinase	
Langlois M.R., Delanghe J.R., De Buyzere M.L., Leroux-Roels G.G.	83
Serum apolipoproteins in heterozygous familial hypercholesterolemia	
Kajinami K., Mabuchi H., Koizumi J., Takeda R.	93
Lipid peroxides in blood from patients with atherosclerosis of coronary and peripheral arteries	
Plachta H., Bartnikowska E., Obara A.	101
Photographic chemiluminescent ELISA for detection of anti-human T-cell leukemia virus type-I antibodies by using synthetic peptides as antigens	
Kuroda N., Nakashima K., Akiyama S., Shiraki H., Maeda Y.	113
<i>Letters to the Editor</i>	
Transferrin, cholesterol and aluminium in Alzheimer's disease	
Corrigan F.M.	121
Correlations between the serum pancreatic enzyme activities and lipid peroxidation in acute pancreatitis	
Matkovic B.	125
Effect of long-term storage on human plasma cholinesterase activity	
Henny J.	129

Vol. 211/3, 1992

Bone metal content in patients with chronic renal failure	
Navarro J.A., Granadillo V.A., Salgado O., Rodriguez-Iturbe B., Garcia R., Delling G., Romero R.A.	133
High-performance liquid chromatographic assay for L-glyceric acid in body fluids. Application in primary hyperoxaluria type 2	
Petrarulo M., Marangella M., Cosseddu D., Linari F.	143
Red blood cells and platelet membrane fatty acids in non-dialyzed and dialyzed uremics	
Girelli D., Azzini M., Olivieri O., Guarini P., Trevisan M.T., Lupo A., Bernich P., Panzetta G., Corrocher R.	155
Distribution of lipoprotein species (LpA-I, LpA-I:A-II) in serum and HDL subfractions of untrained and trained normolipemic men	
Frey I., Kamps H., Baumstark M.W., Berg A., Keul J.	167
A screening method for cystine and homocystine in urine	
Thuy L.P., Nyham W.L.	175
Increase in alkaline phosphodiesterase I activity of tumor-derived cultured fibroblasts from neurofibromatosis patients	
Maruyama E., Takashima S., Arima M.	181
<i>Letters to the Editor</i>	
Selenium in Graves' disease	
Reglinski J., Smith W.E., Wilson R., Halls D.J., McKillop J.H., Thomson J.A.	189
Standardization of Lp(a) assays	
Kostner G.M.	191
Quantities and units for metabolic processes as a function of time. Recommendations 1992	
Ferard G.	S5
Flow cytometry in the clinical laboratory. Principles, applications and problems	
Ruiz-Arguelles A.	S13

Vol. 212/1-2, 1992

Measurement of glycated albumin by the nitroblue tetrazolium colorimetric method	
Mashiba S., Uchida K., Okuda S., Tomita S.	3

A new quantitative assay for glycosaminoglycans	
Thuy L.P., Nyhan W.L.	17
Adaptation of transferrin protein and glycan synthesis	
De Jong G., Van Noort W.L., Feelders R.A., De Jeu-Jaspars C.M.H., Van Eijk H.G.	27
Effects of methylprednisolone and uremia on renal and intestinal calbindin-D in the rat	
Egfford M., Staun M., Olgaard K.	47
Determination of free carnitine and 'total' carnitine in human urine: Derivatization with 4'-bromophenacyl trifluoromethanesulfonate and high performance liquid chromatography	
Minkler P.E., Hoppel C.L.	55
Biliary lipid composition and metastability of biliary cholesterol in patients with acalculous cholesterosis	
Chijiwa K., Nakano K., Makino I., Noshiro H., Yamasaki T.	65
Heterogeneity of antigens related to 7S domain of type IV collagen in serum: Application to patients with chronic liver disease	
Yamada S., Kishimoto Y., Suou T., Kawasaki H., Horie Y.	73
An acquired acute hepatic porphyria: A novel type of δ -aminolevulinate dehydratase inhibition	
Akagi R., Prchal J.T., Eberhart C.E., Sassa S.	79
<i>Letter to the Editor</i>	
Seasonal variation in serum total concentrations of cholesterol and protein in elderly subjects	
Fujimara A., Ohashi K., Ebihara A., Hirose T.	85

Vol. 212/3, 1992

Quantitative analysis of extracellular-superoxide dismutase in serum and urine by ELISA with monoclonal antibody	
Adachi T., Ohta H., Yamada H., Futenma A., Kato K., Hirano K.	89
The BT-PABA/PAS test in tropical diabetes	
Gagee P., Pemberton P., Lobleby R., Chaloner C., Snehalatha C., Mohan V., Braganza J.M.	103
Detection of erythrocyte membrane components in hemoglobin-based blood substitutes	
Biessels P.T., Berbers G.A., Broeders G.C., Landsvater R., Huisman H.G., Bleeker W.K., Bakker J.C.	113
Serum neurone-specific enolase levels in patients with neuroendocrine and carcinoid tumours	
Cunningham R.T., Johnston C.F., Irvine G.B., Buchanan K.D.	123
A colorimetric microassay for glycated collagen based on the thiobarbituric acid method	
Ahmed N., Malik N.S., Furth A.J.	133
Development of a method for the purification of human trypsin by single step affinity chromatography suitable for human isotope incorporation studies	
Ogden J.M., O'Keefe S.J.D., Marks I.N.	141
<i>Letters to the Editor</i>	
The variation of serum lipoprotein (a) during surgical operations	
Magnani B., Brega Massone P.P., Meriggi F., Di Jeso F.	149
Antibodies to thyroid hormone analogues	
John R.	153
Kinetic fluorometric micromethod for screening of newborns for galactosemia — A way to improve assay specificity?	
Tuuminen T., Akerlind G.	155

Vol. 213/1-3, 1992

Platelet hyperreactivity in hyperlipidaemia with special reference to platelet lipids and fatty acid composition	
Chetty N., Naran N.H.	1
Measurement of ascorbic acid in platelets and its relationship to polymorphonuclear leukocyte levels	
McCulloch R.K., Vandongen R.	15

Polyclonal immunoglobulin M: Location of glycation sites	
Menini T., Gugliucci A., Stahl A.J.C.	23
Characterization of the gliadin-derived peptides which are biologically active in coeliac disease	
Cornell H., Wieser H., Belitz H.-D.	37
Copper metabolism — A factor in gestational diabetes?	
Loven A., Romem Y., Pelly I.Z., Holcberg G., Agam G.	51
Differences in erythrocyte sodium transport between human plasma and artificial medium: The role and character of sodium efflux and influx stimulating plasma factors	
Tepper T., Jilderda J.F., Huismans R.M., Van der Hem G.K., De Zeeuw D.	61
Protection of vascular basement membrane and microcirculation from elastase-induced damage with a fluorinated β -lactam derivative	
Maillard J.-L., Favreau C., Vergely I., Reboud-Ravaux M., Joyeau R., Kohaier R., Wakselman M.	75
A pulmonary origin of amniotic fluid thermolabile alkaline phosphatase near term	
Ragoucy-Sengler C., Brodent-Bangou J., Pileire B.	87
Neonatal screening for familial hypercholesterolaemia by immunoturbidimetric assay of apolipoprotein B in dried blood spots	
Bangert S.K., Eldridge P.H., Peters T.J.	95
A two-phase colorimetric method for the measurement of pepsin activity	
Heinova D., Blahovec J., Kasafirek E., Bartik M., Rosival I.	103
<i>Letter to the Editor</i>	
Improved immunoassay for perphenazine	
Uebel R., Van Jaarsveld P., Wium C., Hawtrey A.O.	111

Vol. 214/1, 1993

Establishment of assay kits for the determination of microheterogeneities of alpha-fetoprotein using lectin-affinity electrophoresis	
Shimizu K., Taniichi T., Satomura S., Matsuura S., Taga H., Taketa K.	3
Selective and sensitive determination of urinary total proline and hydroxyproline by gas chromatography with flame photometric detection	
Kataoka H., Nabeshima N., Nagao K., Makita M.	13
A rapid GLC method for the direct analysis of plasma medium chain fatty acids	
Mingrone G., De Gaetano A., Greco A.V., Capristo E., Raguso C., Tataranni P.A., Castagneto M.	21
A clinical evaluation of a direct radioimmunoassay of testosterone	
Holownia P., Conway G.S., Shrivastava U., Round J., Honour J.W.	31
Lipoprotein(a) in diabetes mellitus	
Ritter M.M., Loscar M., Richter W.O., Schwandt P.	45
Use of seronegative samples for the calibration and reporting of serological enzyme immunoassay results	
Sorensen K., Viljanen M.K.	55
Enzymes of lysosomal origin in plasma of twin neonates	
Goi G., Burlina A.B., Bairati C., Bordugo A., Zanardo V., Zaccello F., Tettamanti G., Lombardo A.	61
Measurement of insulin-like growth factor binding protein-I in cervical/vaginal secretions: Comparison with the ROM-check Membrane Immunoassay in the diagnosis of ruptured fetal membranes	
Rutanen E.-M., Pekonen F., Karkkainen T.	73
Characterization of γ -glutamyltranspeptidase from human hepatocellular carcinoma, compared with enzymes from normal liver and cirrhotic liver	
Ohta H., Sawabu N., Kawakami H., Watanabe H., Ozaki K., Toya D., Hattori N.	83
Preparation of microsome fractions of human myometrium. A comparison of three methods	
Lamminen S., Rantala I., Tuimala R.	93
Red blood cell adenosine deaminase activity in multiple sclerosis	
Kopff M., Zakrzewska I., Czernicki J., Klem J., Strzelczyk M.	97

Letters to the Editor

A methodological approach to superoxide dismutase (SOD) activity assay based on inhibition of nitroblue tetrazolium (NBT) reduction	
Durak I., Yurtarslan Z., Canbolat O., Akyol O.	103
Activity of phospholipase A in serum from patients with cystic fibrosis	
Laszlo A., Nemeth M., Gyurkovits K.	105
Basal concentration of the isoenzyme BB of the glycogen phosphorylase b in human blood	
Rabitzsch G., Noll F., Hofmann U., Krause E.-G., Armbruster F.P.	109
Impact of circadian rhythms on the interpretation of stat measurements	
Fuentes-Arderiu X., Rivera-Coll A.	113
The role of lipids in the increased mortality following bereavement	
Coulter F., Armstrong K.M.D., Armstrong E.M., Davies A., Merry E., Skinner E.R., Corrigan F.M.	119

Vol. 214/2, 1993

Circulating proalbumin associated with a second case of antitrypsin Pittsburgh	
Brennan S.O., Sheat J.M., Aiach M.	123
Gravimetric determination of the water concentration in whole blood, plasma and erythrocytes and correlations with hematological and clinicochemical parameters	
Lijnema T.H., Huizenga J.R., Jager J., Mackor A.J., Gips C.H.	129
Circulating tumour-associated mucin concentrations, determined by the CASA assay, in healthy women	
McGuckin M.A., Ramm L.E., Joy G.J., Devine P.L., Ward B.G.	139
Determination of lipoprotein(a): Enzyme immunoassay and immunoradiometric assay compared	
Marz W., Siekmeier R., Gross E., Gross W.	153
Variability of kidney tubular interstitial distortions in glomerulonephritis as measured by ¹ H-NMR urinalysis	
Lundina T.A., Knubovets T.L., Sedov K.R., Markova S.A., Sibeldin L.A.	165
Enzymatic determinations in acute rejection after liver transplantation: Preliminary report on necrosis index	
Gozzo M.L., Avolio A., Forni F., Agnes S., Colacicco L., Barbaresi G., Castagneto M.	175
Antioxidant defenses and its modulation by iron in carrageenan-induced inflammation in rats	
Muntane J., Puig-Parellada P., Fernandez Y., Mitjavila S., Mitjavila M.T.	185
High performance liquid chromatographic determination of free and conjugated bile acids in serum, liver biopsies, bile, gastric juice and feces by fluorescence labeling	
Guldutuna S., You T., Kurts W., Leuschner U.	195
Trace element concentrations (Cu, Zn, Fe) in sera from patients with dilated cardiomyopathy	
Oster O.	209
Cholinesterases in cerebrospinal fluid in patients with meningitis and hydrocephaly	
Tornel P.L., Campoy F.J., Vidal C.J.	219
Plasma malondialdehyde in type 1 and type 2 diabetic patients	
Gallou G., Ruelland A., Legras B., Maugeudre D., Allannic H., Cloarec L.	227
Correction factors for single radial immunodiffusion and electroimmunodiffusion assays of IgA in secretory fluids	
Meillet D., Kapel N., Favennec L., Raichvarg D., Savel J., Gobert J.G.	235
<i>Letters to the Editor</i>	
Contamination of biological samples in metal determination	
Villalba A.	243
Biological variability of twenty-five serum analytes during normal pregnancy	
Alberty R.	245

Vol. 215/1, 1993

Relationship between alcohol consumption and the activity of GTP-binding regulatory proteins in human erythrocyte membranes Nakamura J., Harada S., Tanaka N.	1
Further applications of blood gas analysis to gastric acidity determination Niv Y., Abu-Avid S., Neumann G.	9
Magnesium, zinc and copper in plasma and blood cellular components in children with IDDM Rohn R.D., Pleban P., Jenkins L.L.	21
Serum hyaluronic acid in healthy infants and children and its value as a marker of progressive hepatobiliary disease starting in infancy Trivedi P., Cheeseman P., Mowat A.P.	29
Changes of lipid peroxide levels in blood and liver tissue of patients with obstructive jaundice Tsai L.Y., Lee K.T., Tsai S.M., Lee S.C., Yu H.S.	41
Epidermal growth factor in human breast cancer, endometrial carcinoma and lung cancer. Its relationship to epidermal growth factor receptor, estradiol receptor and tumor TNM Bolufer P., Lluch A., Molina R., Alberola V., Vazquez C., Padilla J., Garcia-Conde J., Llopis F., Guillem V.	51
Biotin-specific 110-kDa biotinidase from human serum Oizumi J., Hayakawa K.	63
Increased platelet activating factor in the tracheal aspirates from neonates with patent ductus arteriosus Koyama N., Ogawa Y., Kamiya K., Eguchi H., Tanaka T., Takasaki J., Ohama Y., Itakura Y., Kaneko K.	73
Detection of heterozygotes for fructose-1,6-diphosphatase deficiency by measuring fructose-1,6-diphosphatase activity in monocytes cultured with calcitriol Kikawa Y., Takano T., Nakai A., Shigematsu Y., Sudo M.	81
Frequency of the XbaI, EcoRI, PvuII and MspI polymorphisms of the apolipoprotein B gene in relation to hypercholesterolaemia in the general population Series J.J., Gaffney D., Packard C.J., Shepherd J.	89
A new highly sensitive radioimmunoassay for human calcitonin useful for physiological studies Schifter S.	99
Acceleration of hemoglobin glycation with aging Nakashima K., Nishizaki O., Andoh Y.	111
<i>Letters to the Editor</i>	
Ethanediol stabilised quality control serum for measurement of zinc Swaminathan S., Seshadri M.S., Srikrishna G., Selvakumar R., Kanagasabapathy A.S.	119
Serum trehalase activity in patients with rheumatoid arthritis Yoshida K., Mizukawa H., Haruki E.	123
Erratum: Distribution of lipoprotein species (LpA-I, LpA-I:AI) in serum and HDL subfractions of untrained and trained normolipemic men (Clinica Chimica Acta, 211 (1992) 167-173) Frey I., Kamps H., Baumstark M.W., Berg A., Keul J.	125

Vol. 215/2, 1993

Molecular weight and substrate characteristics of human serum arylesterase following purification by immuno-affinity chromatography Tanaka-Kawai H., Yomoda S.	127
A method for the assay of phosphatidylinositol-specific phospholipase D activity in serum Raymond F.D., Fortunato G., Moss D.W.	139
Survey of methods for measuring human anti-mouse antibodies Kricka L.J., Schmerfeld-Pruss D., Kaladas P., Colcher D., Scassellati G.A., Seccamani E., Courtenay-Luck N.S., Kosmas C., Beatty B.G., Beatty J.D., Chatenoud L., Zimmer A.M., Auerbach B.	153

Enzymatic diagnosis of steroid sulfatase deficiency by high performance liquid chromatography Ohsaki M., Matsumoto T., Sakura N., Ueda K.	165
Glycan microheterogeneity for α_1 -antitrypsin in serum and meconium from normal and cystic fibrosis patients by crossed immunoaffinoelectrophoresis with different lectins (Con A, LCA, WGA) Duthel S., Revol A.	173
Phosphorylation of insulin-like growth factor-binding protein-1 increases in human amniotic fluid and decidua from early to late pregnancy Koistinen R., Angervo M., Leinonen P., Hakala T., Seppala M.	189
Vitamin E attenuates induction of elastase-like activity by tumor necrosis factor- α , cholesterol-3 β ,5 α ,6 β -triol and linoleic acid in cultured endothelial cells Toborek M., Hennig B.	201
Changes in immunoreactive manganese-superoxide dismutase concentration in human serum after 93 h strenuous physical exercise Ohno H., Kayashima S., Nagata N., Yamashita H., Ookawara T., Taniguchi N.	213
Cigarette-smoking, birthweight, thiocyanate and fluorescent lipid-peroxidation products in maternal and cord plasma Pre J., Colau J.C., Vassy R.	221
Measurement of histidase activity in human fingernail by reversed-phase HPLC Fujitaka M., Eguchi T., Sakura N., Ueda K.	227
<i>Letter to the Editor</i>	
Effects of ethanol on the assay of acyl-CoA:cholesterol acyltransferase (ACAT) Smith J.L., Madden L.J., De Jersey J.	233

Vol. 216/1-2, 1993

Deregulated production of interleukin-4 (IL4) in autoimmune thyroid disease assayed with a new radioimmunoassay Hirooka Y., Kayama M., Ohga S., Kimura M., Hasegawa M., Shin K., Nogimori T., Ishizuki Y., Mitsuna T.	1
Variable expression of leukocyte cytosolic broad-specificity β -glucosidase activity Forsyth G.W., Romero K.M., Alverson J., VanderJagt D.J., Glew R.H.	11
A novel lectin-based enzyme-linked immunosorbent assay for the measurement of IgA1 in serum and secretory IgA1 in secretions Cullina M.J., Grealley J.F.	23
Intrathecal synthesis of IgM and IgA in neurological diseases: Comparison of two formulae with isoelectric focusing Kaiser R., Lucking C.H.	39
Determination of acylcarnitines in urine of patients with inborn errors of metabolism using high-performance liquid chromatography after derivatization with 4'-bromophenacylbromide Poorthuis B.J.H.M., Jille-Vlckova T., Onkenhout W.	53
Hyperphenylalaninemia and pterin metabolism in serum and erythrocytes Ponzzone A., Guardamagna O., Spada M., Ponzzone R., Sartore M., Kierat L., Heizmann C.W., Blau N.	63
Release of β -hexosaminidase isoenzymes in cultured human fibroblasts Hultberg B., Isaksson A., Nordstrom M., Kjellstrom T.	73
A 125 I-radioimmunoassay for diethylstilbestrol in serum of patients with prostatic cancer treated with stilphostrol Economou E.V., Livanou E., Evangelatos G.P., Ithakissios D.S.	81
Fetal intestinal and renal origins of trehalase activity in human amniotic fluid El-liger M.-A., Dallaire L., Potier M.	91
Determination of proteins by a reverse biuret method combined with the copper-bathocuproine chelate reaction Matsushita M., Irino T., Komoda T., Sakagishi Y.	103

Lipoprotein lipase mass and activity in severe hypertriglyceridemia	
Kobayashi J., Hashimoto H., Fukamachi I., Tashiro J., Shirai K., Saito Y., Yoshida S.	113
Decreased renal excretion of β -hexosaminidase in adults with insulin-dependent diabetes mellitus and normal renal function	
Romero K.M., Butcher B.A., Boyle P.J., Fry D.E., Glew R.H.	125
Enzyme reactor for urinary acylcarnitines assay by reversed-phase high-performance liquid chromatography	
Matsumoto K., Takahashi M., Takiyama N., Misaki H., Matsuo N., Murano S., Yuki H.	135
Erythrocyte galactokinase assay with high performance liquid chromatography	
Mizoguchi N., Eguchi T., Sakura N., Ueda K.	145
Cerebrospinal fluid ferritin in brain necrosis: Evidence for local synthesis	
Keir G., Tasdemir N., Thompson E.J.	153
Mono-, di- and triacylglycerol concentrations in human plasma: Effects of heparin injection and of a high-fat meal	
Fielding B.A., Humphreys S.M., Allmann R.F.C., Frayn K.N.	167
Characteristics of membrane-bound 5'-nucleotidase on human prostasomes	
Fabiani R., Ronquist G.	175
Hormone-sensitive lipase: Quantitation of enzyme activity and mRNA level in small biopsies of human adipose tissue	
Frayn K.N., Langin D., Holm C., Belfrage P.	183
Apolipoprotein E phenotypes and genotypes as determined by polymerase chain reaction using allele-specific oligonucleotide probes and the amplification refractory mutation system in children with insulin-dependent diabetes mellitus	
Stavijeno-Rukavina A., Sertic J., Salzer B., Dumic M., Radica A., Fumic K., Krajina A.	191
<i>Letters to the Editor</i>	
Prostatic specific antigen and prostatic acid phosphatase measured by radioimmunoassay in vaginal washings from cases of suspected sexual assault	
Roach B.A., Vladutiu A.O.	199
The application of molecular biology techniques to the diagnosis of hyperlipidaemia and other risk factors for cardiovascular disease	
Humphries S.E.	S5

Vol. 217/1, 1993

Clinical impact of genomic analysis in the diagnosis and management of genetic disease and cancer: Introduction	
Werner M., Garrett C.T.	1
The diagnosis of inborn errors of metabolism by examination of the genotype	
Hill R.E.	3
Genotype-phenotype correlations in phenylketonuria	
Trefz F.K., Burgard P., Konig T., Goebel-Schreiner B., Lichter-Konecki U., Konecki D., Schmidt E., Schmidt H., Bickel H.	15
Cystic fibrosis: The impact of analytical technology for genotype-phenotype studies	
Tummler B., Dork T., Kubesch P., Fislage R., Kalin N., Neumann T., Wulbrand U., Wulf B., Steinkamp G., Von der Hardt H.	23
Variable phenotypic expression of genotypic abnormalities in the porphyrias	
Hindmarsh J.T.	29
Comparative clinical evaluation of biochemical and genomic tumor markers	
Werner M., Ballo M.S., Gallagher J.V.	39
myc Amplification: Regulation of Myc function	
Koskinen P.J., Makela T.P., Vastrik I., Alitalo K.	57
Genetic and molecular heterogeneity of breast cancer cells	
Callahan R., Cropp C., Merlo G.R., Diella F., Venesio T., Lidereau R., Cappa A.P.M., Liscia D.S.	63

Detection of minimal residual leukemia by the polymerase chain reaction: Potential implications for therapy	
Bartram C.R.	75
Molecular diagnostics: Issues of utilization, regulation and organization	
Garrett C.T., Ferreira-Centeno A., Nasim S.	83

Vol. 217/2, 1993

In vitro degradation of very low density lipoprotein from diabetic patients by lipoprotein lipase	
Saheki S., Hitsumoto Y., Murase M., Takeuchi N., Uchida K.	105
Increased fragmentation of urinary fibronectin in cancer patients detected by immunoenzymometric assay using domain-specific monoclonal antibodies	
Katayama M., Kamihagi K., Nakagawa K., Akiyama T., Sano Y., Ouchi R., Nagata S., Hino F., Kato I.	115
Demonstration of immunoglobulin G with affinity for dopamine in cerebrospinal fluid from psychotic patients	
Bergquist J., Bergquist S., Axelsson R., Ekman R.	129
Comparison of a mechanized version of the 'Konig' reaction and a fluorescence polarization immunoassay for the determination of nicotine metabolites in urine	
Pickert A., Lingenfelser T., Pickert C., Birbaumer N., Overkamp D., Eggstein M.	143
Value of faecal alpha-1-antitrypsin, haemoglobin and a chemical occult blood test in the detection of gastrointestinal disease	
Moran A., Lawson N., Morrow R., Jones A., Asquith P.	153
Ascorbate concentration in human cerebrospinal fluid (CSF) and serum. Intrathecal accumulation and CSF flow rate	
Reiber H., Ruff M., Uhr M.	163
Semi-automatic method for determination of different isoforms of carbohydrate-deficient transferrin	
Lof K., Koivula T., Seppa K., Fukunaga T., Sillanauke P.	175
Inactivation of glutathione peroxidase following entrapment of purified α or β hemoglobin chains in human erythrocyte	
Grelloni F., Gabbianelli R., Santroni A.M., Falcioni G.	187
The preferential site of non-enzymatic glycation of human apolipoprotein A-I in vivo	
Calvo C., Ulloa N., Campos M., Verdugo C., Ayrault-Jarrier M.	193
A study of growth factors in human renal cysts with or without renal cell carcinoma	
Taide M., Kanda S., Eguchi J., Igawa T., Kanetake H., Saito Y.	199
Cyclooxygenase pathway in dermal fibroblasts from patients with metabolic disorders of peroxisomal origin	
Malle E., Leis H.J., Steinmetz A., Paschke E., Hoefler G.	205
Studies on zinc and angiotensin-converting enzyme	
Buttery J.E., Stuart S.	213
<i>Letters to the Editor</i>	
Bile acid metabolism in three patients with mevalonic aciduria due to mevalonate kinase deficiency	
Gibson K.M., Stellaard F., Hoffmann G.F., Rating D., Hrebicek M., Jakobs C.	217
Urinary excretion patterns of modified nucleosides, pseudouridine and 1-methyladenosine, in healthy individuals	
Itoh K., Aida S., Ishiwata S., Sasaki S., Ishida N., Mizugaki M.	221
Fab treatment in acute digitalis intoxication: Reliability of serum digoxin determination with the Stratus system	
Bizzaro N., Finco B., Milani L.	225
A rare case of IgD gammopathy associated with free light-chain cryoglobulinemia and a gelling Bence Jones protein	
Myara I., Llibat B., Rossert J., Cosson C., Tenenhaus D., Moatti N.	229
Abnormal serum lipid levels in subjects with gastric xanthoma	
Chang F.-Y., Shih C.-Y., Lee S.-D.	233

Vol. 218/1, 1993

High concentrations of immunoreactive gliostatin/platelet-derived endothelial cell growth factor in synovial fluid and serum of rheumatoid arthritis	
Asai K., Hirano T., Matsukawa K., Kusada J., Takeuchi M., Otsuka T., Matsui N., Kato T.	1
Human seminal deoxyribonuclease I (DNase I): Purification, enzymological and immunological characterization and origin	
Yasuda T., Sawazaki K., Nadano D., Takeshita H., Nakanaga M., Kishi K.	5
A radioimmunoassay for pancreatic elastase I: Use of α_1 -antitrypsin-elastase I conjugate as both standard and tracer to eliminate the influence of anti-elastase I autoantibodies	
Asada H., Shibata K., Uchida K., Takeyama Y., Saitoh Y., Ohyanagi H.	17
Repertoires of natural autoantibodies against muscle tissue are independent of age or gender in normal human adults. A Western blot study	
Zimmermann C.W.	29
Malondialdehyde adducts to, and fragmentation of, apolipoprotein B from human plasma	
Lecomte E., Artur Y., Chancerelle Y., Herbeth B., Galteau M.-M., Jeandel C., Siest G.	39
Serum tissue inhibitor of metalloproteinases in patients with chronic liver disease and with hepatocellular carcinoma	
Murawaki Y., Yamamoto H., Kawasaki H., Shima H.	47
A rate turbidimetric immunoassay for theophylline using biotin-avidin system	
Oh C., Kim J., Kearns B., Cheng A., Dobashi T.	59
5,6-Dihydrouridine: A marker ribonucleoside for determining whole body degradation rates of transfer RNA in man and rats	
Topp H., Duden R., Schoch G.	73
Comparison of lipoprotein(a) assay methods in serum and in a plasminogen-free fraction	
Baldo-Enzi G., Baiocchi M., Crepaldi G.	83
Quantitation of plasma oxidatively modified low density lipoprotein by sandwich enzyme linked immunosorbent assay	
Wang H., Chen S., Kong X., Wang X., Chang G., Xu S., Luo Z., Xie Y.	97
Biochemical and immunological characteristics of neutrophil alkaline phosphatase in Down's syndrome	
Vergnes H.A., Denier C.J., Brissan-Lougarre A., Grozdea J.G., Bourrouillou G., Colombies P., Klæbe A.G., Perie J.J.	105
Vigabatrin — Interference with urinary amino acid analysis	
Preece M.A., Sewell I.J., Taylor J.A., Green A.	113
Letter to the Editor	
Oxidative metabolism and severe asthma in children	
Malvy J.-M.D., Lebranchu Y., Richard M.-J., Arnaud J., Favier A.	117

Vol. 218/2, 1993

Rapid enzyme immunoassay of anti-streptokinase antibodies in human plasma	
Bom V.J.J., Brugemann J., Van Der Schaaf W., Van Wijk R.T., Van Der Meer J.	121
The ins/del polymorphism in the signal sequence of apolipoprotein B has no effect on lipid parameters	
Gaffney D., Freeman D.J., Shepherd J., Packard C.J.	131
Fluorescence-based diagnosis of lipid storage diseases by analysis of the culture medium of skin fibroblasts	
Vered A., Eugenio M., Arie D., Augusto P., Sergio M., Shimon G.	139
Two-dimensional gel electrophoresis of four serum samples from patients with IgD myeloma	
Stulik J., Tichy M., Kovarova H.	149
Rapid determination of glutamine in biological samples by high-performance liquid chromatography	
Teerlink T., Hennekes M.W.T., Van Leeuwen P.A.M., Houdijk A.	159

Urinary excretion of modified nucleosides as biological marker of RNA turnover in patients with cancer and AIDS	
Nakano K., Nakao T., Schram K.H., Hammargren W.M., McClure T.D., Katz M., Petersen E.	169
Increased excretion of endogenous urinary leukotriene E ₄ in extrahepatic cholestasis	
Mayatepek E., Pecher G.	185
High-performance liquid chromatographic microassay for L-glutamate:glyoxylate aminotransferase activity in human liver. Application in primary hyperoxaluria type 1	
Petrarulo M., Pellegrino S., Marangella M., Cosseddu D., Linari F.	193
Simultaneous determination of lead, bismuth and thallium in plasma and urine by inductively coupled plasma mass spectrometry	
Mauras Y., Premel-Cabic A., Berre S., Allain P.	201
Rapid automated enzymatic measurement of phenylalanine in plasma and blood spots	
Taylor R.P., Smith I.C., Standing S.J.	207
Compact gas chromatography used with thin-layer chromatography for assessment of abnormalities of organic acids	
Lee B.Y., Thurmon T.F.	215
Effects of gemfibrozil on the activities of plasma lipolytic enzymes in normolipidemic subjects	
Horsmans Y., Desager J.P., Harvengt C.	223
<i>Letter to the Editor</i>	
Variation in the amount of stimulatory GTP-binding protein (Gs) in human erythrocyte membranes in relation to alcohol consumption	
Nakamura J.	229

Vol. 219/1-2, 1993

A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 1 (interstitial collagenase) using monoclonal antibodies	
Zhang J., Fujimoto N., Iwata K., Sakai T., Okada Y., Hayakawa T.	1
Development and application of an enzyme immunoassay for tenascin	
Washizu K., Kimura S., Hiraiwa H., Matsunaga K., Kuwabara M., Ariyoshi Y., Kato K., Takeuchi K.	15
Evaluation of thyroid function in patients with thyroid hormone autoantibodies	
Sakata S., Komaki T., Ogawa T., Takuno H., Matsui I., Sarui H., Kojima N., Takamatsu J., Miura K.	23
Blood and liver lipid peroxide status after chronic ethanol administration in rats	
Coudray C., Richard M.J., Faure H., Favier A.	35
Evidence for a multifactorial control of serum erythropoietin concentration in liver disease	
Pirisi M., Fabris C., Falletti E., Soardo G., Toniutto P., Gonano F., Bartoli E.	47
Levels and physicochemical properties of lipoprotein subclasses in moderate hypertriglyceridemia	
Manzato E., Zambon S., Zambon A., Cortella A., Sartore G., Crepaldi G.	57
Postoperative studies on parathyroid hormone secretion in patients operated on for primary hyperparathyroidism	
Bergenfels A., Norden N.E., Ahren B.	67
Bismuth in human serum: Reference interval and concentrations after intake of a therapeutic dose of colloidal bismuth subcitrate	
Vanhoe H., Versieck J., Vanballenberghe L., Dams R.	79
Studies on the extraction of phospholipids from erythrocyte membranes in the rat	
Eder K., Reichlmayr-Lais A.M., Kirchgessner M.	93
Neonatal screening for maple syrup urine disease by an enzyme-mediated colorimetric method	
Wendel U., Gonzales J., Hummel W.	105
Dissipation of the calcium gradient in human erythrocytes results in increased heat production	
Engstrom I., Waldenstrom A., Nilsson-Ehle P., Ronquist G.	113
Determination of changes in serum lathosterol during treatment with simvastatin to evaluate the role of lathosterol as a parameter for whole body cholesterol synthesis	
De Cuyper I., Wolthers B.G., Van Doormaal J.J., Wijnandts P.N.	123

Serum sialic acid and acute phase proteins in type 1 and type 2 diabetes mellitus	
Crook M.A., Tutt P., Simpson H., Pickup J.C.	131
Radioimmunoassay of glutathione peroxidase in human serum	
Huang W., Akesson B.	139
Lipoprotein(a) phenotyping using a computerized micro scale and phenotype frequencies in a healthy Japanese population	
Abe A., Noma A., Itakura H.	149
Changes in glutathione metabolic enzymes in erythrocytes of <i>Plasmodium vivax</i> infected patients	
Mathews S.T., Selvam R.	159
The vitamin A fat-loading test in young normolipidemic subjects	
Heller F.R., Vandenplas C., Desager J.P., Harvenge C.	167
Abnormally fucosylated haptoglobin as a marker for alcoholic liver disease but not excessive alcohol consumption or non-alcoholic liver disease	
Chambers W., Thompson S., Skillen A.W., Record C.O., Turner G.A.	177
Differences in the time course of creatine kinase-MB activity and mass concentration after acute myocardial infarction	
Baum H.H.E., Schwab I., Boekstegers P., Steinbeck G., Neumeier D.	183
Simple mixing technique for the preparation of blood samples of known oxygen saturation	
Clark D.J.	189
<i>Letter to the Editor</i>	
Effect of ethanol on prolidase I and prolidase activity in the human hepatoma cell line Hep G2	
Myara I., Cosson C., Pourci M.L., Moatti N., Lemonnier A.	195

Vol. 220/1, 1993

HPLC determination of oxidized and reduced pyridine coenzymes in human erythrocytes	
Micheli V., Simmonds H.A., Bari M., Pompucci G.	1
Urinary growth hormone measurements as a marker of renal tubular function in diabetes mellitus	
Turner G., Coates P., Porter S., Peters J.R., Woodhead J.S.	19
A one-step sandwich enzyme immunoassay for tissue inhibitor of metalloproteinases-2 using monoclonal antibodies	
Fujimoto N., Zhang J., Iwata K., Shinya T., Okada Y., Hayakawa T.	31
Regulation of transferrin receptor expression and distribution in vitro cultured human cytotrophoblasts	
Starreveld J.S., Van Dijk H.P., Kroos M.J., Van Eijk H.G.	47
Scintillation proximity assay for calcitriol in serum without high pressure liquid chromatography	
Wildermuth S., Dittrich K., Schmidt-Gayk H., Zahn I., O'Riordan J.L.H.	61
A fibroblast glutaryl-CoA dehydrogenase assay using detritiation of ³ H-labelled glutaryl-CoA: Application in the genotyping of the glutaryl-CoA dehydrogenase locus	
Christensen E.	71
The hepatobiliary disease marker serum lanine aminopeptidase predominantly comprises an isoform of the haematological myeloid differentiation antigen and leukaemia marker CD-13/gp150	
Favaloro E.J., Browning T., Nandurkar H.	81
An improved Bradford protein assay for collagen proteins	
Lopez J.M., Imperial S., Valderramad R., Navarro S.	91
False rise in C-reactive protein in a pateint with monoclonal IgM immunoglobulin	
Ponge T.D., Le Carrer D.L., Sagniez M.M., Pontet M., Cottin S.L.	101
The glycosylation of haptoglobin in rheumatoid arthritis	
Thompson S., Dargan E., Griffiths I.D., Kelly C.A., Turner G.A.	107
<i>Letters to the Editor</i>	
Senile cataractic lenses do not accumulate galactitol in either lactose tolerant or intolerant subjects	
Wacker H., Soldati L., Simonelli F., Richter C., Gazzaniga A., Auricchio S., Semenza G.	115
Non-enzymatic glycosylation of mouse monoclonal antibody reduces its binding activity to antigen	
Sasaki Y., Mori T., Shiiki H., Dohi K., Ishikawa H.	119

- Measurement of seminal plasma α -amylase that is not inhibited by monoclonal antibodies against the salivary isoenzyme
Huguet J., Cortes J.D., Arranz B., Fuentes-Arderiu X. 123

Vol. 220/2, 1993

- Urinary macromolecules are promoters of calcium oxalate nucleation in human urine: Turbidimetric studies
Rodgers A.L., Ball D., Harper W. 125
- Increase in plasma concentration of ubiquitin in dialysis patients: Possible involvement in β_2 -microglobulin amyloidosis
Okada M., Miyazaki S., Hirasawa Y. 135
- Determination of human immunoglobulin A and secretory immunoglobulin A in bronchoalveolar lavage fluids by solid phase enzyme immunoassay
Sato T., Kitajima A., Ohmoto S., Chikuma M., Kado M., Nagai S., Izumi T., Takeyama M., Masada M. 145
- Quantification of 3-methylglutaconic acid in urine, plasma, and amniotic fluid by isotope-dilution gas chromatography/mass spectrometry
Kelley R.I. 157
- Improved PCR/NcoI method for the molecular diagnosis of medium chain acyl-CoA dehydrogenase deficiency using dried blood samples: Two-stage amplification using two different sets of primers improves accuracy and sensitivity
Nagao M., Raymond D., Kim J., Tanaka K. 165
- Alterations in the concentration of an apolipoprotein E-containing subfraction of plasma high density lipoprotein in coronary heart disease
Wilson H.M., Patel J.C., Russell D., Skinner E.R. 175
- 2,4-Diamino-7-hydroxy-pteridines in biological fluids of patients on high-dose methotrexate
Dhondt J.-L., Millot F., Hayte J.-M., Mazingue F., Farriaux J.-P. 189
- Sensitive chemiluminescent immunoassay for human granulocyte colony-stimulating factor (G-CSF) in clinical applications
Kiriya R., Chichibu K., Matsuno T., Ohsawa N. 201
- Anion exchange protein (band 3) in human erythrocytes characterized by different abnormalities
Giuliani A., Marini S., Ferroni L., Condo S.G., Ramacci M.T., Giardina B. 211
- Investigation of β -oxidation in human liver biopsies
Eaton S., Record C.O., Bartlett K. 219
- Letters to the Editor*
- Enzyme-linked immunoglobulin which binds both lactate dehydrogenase and hydroxybutyric dehydrogenase
Taguchi H., Nishiya K., Yamanaka S., Saika K., Morishita N., Sasaki M., Miyoshi I. 225
- A simple coloured-paper thin-layer chromatography test as a backup procedure for vanillylmandelic acid and homovanillic acid evaluation in a neuroblastoma screening program
Mathieu P., Philip T., Greffe J., Montague A., Lacroix C. 229
- A comparative study of thyroid function in African neonates: A reference thyroid profile
Das S.C., Isichei U.P. 233

Vol. 221/1-2, 1993

- An investigation of argininosuccinic acid anhydrides in argininosuccinic acid lyase deficiency by $^1\text{H-NMR}$ spectroscopy
Burns S.P., Iles R.A. 1
- Is an increase of deoxyribonucleases activity in uraemic lymphocytes caused by their augmented synthesis in these cells?
Adamiec R., Szweczyk Z., Szmidzinski R., Szopa J. 15

Microwave oven digestion procedure for atomic absorption spectrometry analysis of bone and teeth	
Gil F., Perez M.L., Facio A., Villanueva E., Tojo R., Gil A.	23
Insulin receptor binding to erythrocytes in the first half of pregnancy is increased in healthy pregnant women as compared with non-pregnant or gestational diabetic women	
Schmon B., Desoye G., Friedl H., Hofmann H., Weiss P.A.M., Hagmuller K.	33
The lysosomal N-acetyl- β -D-glucosaminidase (NAG) isoenzymes in plasma: Study of distribution in a general population by a simple routine chromatofocusing procedure	
Goi G., Bairati C., Roggi C., Maccarini L., Tettamanti G., Meloni Lombardo C.A.	47
Changes in the glycoforms of rat alpha-1-acid glycoprotein during experimental polyarthritis	
Venembre P.C., Cong H.N., Biou D.R., Durand G.M.	59
Cholesteryl ester transfer activity in lipoprotein lipase deficiency and other primary hypertriglyceridemias	
Iglesias A., Contreras J.A., Martinez-Pardo M., Entrala A., Herrera E., Lasuncion M.A.	73
A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 2 (72-kDa gelatinase/type IV collagenase) using monoclonal antibodies	
Fujimoto N., Mouri N., Iwata K., Ohuchi E., Okada Y., Hayakawa T.	91
Expression and nature of the alkaline phosphatase gene in cultured osteosarcoma cells	
Stinson R.A., Thacker J.D., Lin C.C.	105
A novel HPLC method for the simultaneous quantification of monosaccharides and disaccharides used in tests of intestinal function and permeability	
Sorensen S.H., Proud F.J., Adam A., Rutgers H.C., Batt R.M.	115
LDL sialic acid content in patients with coronary artery disease	
Ruelland A., Gallou G., Legras B., Paillard F., Cloarec L.	127
The investigation of UDPGlucose and UDPGalactose concentration in red blood cells of patients with classical galactosaemia	
Keavill N.J., Holton J.B., Allen J.T.	135
High performance liquid chromatographic profiling of tryptophan and related indoles in body fluids and tissues of carcinoid patients	
Kema I.P., Schellings A.M.J., Hoppenbrouwers C.J.M., Rutgers H.M., De Vries E.G.E., Muskiet F.A.J.	143
Characterization of apo(a) polymorphism by a modified immunoblotting technique in an Italian population sample	
Geroldi D., Bellotti V., Buacaglia P., Bonetti G., Gazzaruso C., Caprioli A., Fratino P.	159
Diagnosis of mucopolysaccharidoses in a clinically selected population by urinary glycosaminoglycan analysis: A study of 2,000 urine samples	
Piraud M., Boyer S., Mathieu M., Maire I.	171
Serum bile acid concentrations in mild liver cirrhosis	
Greco A.V., Mingrone G.	183
An increase of serum remnant-like particles in non-insulin-dependent diabetic patients with microalbuminuria	
Shimizu H., Mori M., Saito T.	191
Study of chronic granulomatous disease by a nitroblue tetrazolium densitometric kinetic test: A new research method	
Iacobini M., Torre A., Palumbo G., Del Principe D., Digilio G., Balducci L.	197
β -N-Acetylhexosaminidase in spinal muscular atrophy fibroblasts	
Tassi C., Beccari T., Costanzi E., Orlacchio A.	203
Enzyme immunoassay of serum type IV collagen in anti-HCV positive chronic liver diseases	
Matsuzaki Y., Yoshiga S., Sugitani T., Shoda J., Tanaka N., Osuga T.	209
The validity of the ethanol precipitation method for the measurement of mucin content in human gastric juices and its possible relationship to gastroduodenal diseases	
Azuumi Y., Ichikawa T., Ishihara K., Hotta K.	219

Vol. 222/1-2, 1993

Introduction — The OpenLabs Project	
O'Moore R.	1
Object-oriented methods	
Solberg H.E.	3
Practical experience with graphical user interfaces and object-oriented design in the clinical laboratory	
Wells I.G., Cartwright R.Y., Farnan L.P.	13
RefVal for Windows — migrating to the Windows environment with the help of Visual Basic	
Kenny D., Solberg H.E.	19
Supporting interoperability between medical knowledge-based systems: Experience from pilot implementations	
Boran G.P., Given P.G., Grimson J.B., O'Moore R.R.	23
Interpretative reporting and alarming based on laboratory data	
Nykanen P., Boran G., Pince H., Clarke K., Yearworth M., Willems J.L., O'Moore R.	37
To test or not to test, that is the question	
Hasman A., Pop P., Winkens R.A.G., Blom J.L.	49
Problem-oriented management of laboratory work through dynamic test scheduling	
Brender J., Magdal U., Wiegell B., Schioler T., McNair P.	57
Knowledge based lipid management system for general practitioners	
Sinnott M.M., Carr B., Markey J., Brosnan P., Boran G., Grimson J., McGill T., Feely J., O'Moore R.	71
Knowledge-based approach to clinical decision-support system, with an application in tetanus serology	
Schroder J.P., Kuhlmann W.D., Trendelenburg C.	78
Competence reasoning — Handling ambiguous and imprecise data in the Pro.M.D. expert system shell	
Fohl B.	85
Integrating knowledge-based systems and databases	
Grimson J.B.	101
Clinical utility of an international thyroid database	
Nolan J., Schioler T., Brosnan P., Irjala K., Nuutila P., McNair P., O'Moore R.	117
Arden Syntax as a standard for knowledge bases in the clinical chemistry laboratory	
Johansson B., Bergvin L.	123
Analytical quality goals and assessment to ensure transferability of laboratory results	
Groth T., De Verdier C.-H.	129
Distribution of laboratory test results to primary health care centres with the EDIFACT standard	
Bergvin L., Johansson B., Borjesson U.	141
Guidelines for hospital laboratory computer systems	
Kenny D.	147
The Ninth International Conference on Computing in Clinical Laboratories: Foreword	
Kenny D., Solberg H.E.	vii

Vol. 223/1-2, 1993

Comparative drug influence on peroxide mediated increase of cytosolic calcium in human endothelial cells	
David M., Horvath G., Schimke I., Nagy I., Mueller M.M.	1
Metabolites of dietary (soya) isoflavones in human urine	
Kelly G.E., Nelson C., Waring M.A., Joannou G.E., Reeder A.Y.	9
Excretion of 3,6-epoxydicarboxylic acids in peroxisomal disorders	
Pitt J.J., Poulos A.	23
Immunonephelometric/turbidimetric apolipoprotein B assays for the clinical laboratory	
Levinson S.S., Wagner S.G.	31

Effect of tamoxifen on serum cholesterol and lipoproteins during chemohormonal therapy Dnistrian A.M., Schwartz M.K., Greenberg E.J., Smith C.A., Schwartz D.C.	43
Cholesterol in remnant-like lipoproteins in human serum using monoclonal anti apo B-100 and anti apo A-I immunoaffinity mixed gels Nakajima K., Saito T., Tamura A., Suzuki M., Nakano T., Adachi M., Tanaka A., Tada N., Nakamura H., Campos E., Havel R.J.	53
The relationship between serum levels of lipoprotein(a) and proteins associated with the acute phase response Ledue T.B., Neveux L.M., Palomaki G.E., Ritchie R.F., Craig W.Y.	73
Lactate dehydrogenase isoenzyme patterns in blood cells from histiocytosis X children Perlino E., Marra E., Maenza S., De Terlizzi M., Coppola B.C., Santostasi T., Quagliariello E.	83
High adenosine content in human uterine smooth muscle compared with striated skeletal muscle Ronquist G., Wedenberg K., Waldenstrom A., Ericson A., Ulmsten U.	93
Bilirubin inhibits protein kinase C activity and protein kinase C-mediated phosphorylation of endogenous substrates in human skin fibroblasts Amit Y., Boneh A.	103
Lipoprotein composition in the insulin-deficient non-acidotic phase of type I diabetic patients and early evolution after the start of insulin therapy Reverter J.L., Senti M., Rubies-Prat J., Lucas A., Salinas I., Pizarro E., Pedro-Botet J., Sanmarti A.	113
Lipoprotein(a) serum concentration in familial combined hyperlipidemia Fonda M., Da Col P.G., La Verde R., Battello C., Fisicaro M., Tonizzo M., Cattin L.	121
Effect of bone resorbing factors on u-PA and its specific receptor in osteosarcoma cell line Nonaka T., Matsumoto H., Shimada W., Okada K., Fukao H., Ueshima S., Kikuchi H., Tanaka S., Matsuo O.	129
Extremely high HDL levels in a patient with multiple symmetric lipomatosis Deiana L., Pes G.M., Carru C., Campus G.V., Tidore M.G.B., Cherchi G.M.	143
An enzymatic assay of inorganic phosphate in serum using nucleoside phosphorylase and xanthine oxidase Ungerer J.P.J., Oosthuizen H.M., Bissbort S.H.	149
Free thyroxine measured in undiluted serum by dialysis and ultrafiltration: Effects of non-thyroidal illness, and an acute load of salicylate or heparin Faber J., Waetjen I., Siersbaek-Nielsen K.	159
Total pancreatic and salivary serum iso-amylase activities in alcohol misusers in relapse and remission and in alcoholic liver disease Deenmamode J.M.J.M., Sherwood R.A., Sherman D.I.N., Peters T.J.	169
Prenatal screening for Down's syndrome and human chorionic gonadotropin isoforms Defasque P., Pigny P., Racadot A., Farriaux J.-P., Boersma A.	173
Stability of human plasma atrial natriuretic peptide during storage at -80°C Bhaggoe U.M., Boomsma F., Admiraal P.J.J., Man in 't Veld A.J., Schalekamp M.A.D.H.	179
<i>Letters to the Editor</i>	
Pedigree of serum extracellular-superoxide dismutase level Adachi T., Nakamura M., Yamada H., Kitano M., Futenma A., Kato K., Hirano K.	185
The molecular mass and concentrations of protein 1 or Clara cell protein in biological fluids: A reappraisal Bernard A., Dumont X., Roels H., Lauwerys R., Dierynck I., De Ley M., Stroobant V., De Hoffmann E.	189

Vol. 224/1, 1994

Alteration of asparagine-linked glycosylation in serum transferrin of patients with hepatocellular carcinoma Matsumoto K., Maeda Y., Kato S., Yuki H.	1
Non-specific steroidal esterase activity and distribution in human and other mammalian tissues Lund-Pero M., Jeppson B., Arneklo-Nobin B., Sjogren H.-O., Holmgren K., Pero R.W.	9

Very-fast ultracentrifugation of human plasma lipoproteins: Influence of the centrifugal field on lipoprotein composition Leonhardt W., Pietzsch J., Nitzsche S.	21
Glomerular filtration rate and creatinine production in adult icteric patients Delanghe J.R., Louagie H.K., De Buyzere M.L., Leroux-Roels G.G.	33
The allopurinol loading test for identification of carriers for ornithine carbamoyl transferase deficiency: Studies in a healthy control population and females at risk Sebesta I., Fairbanks L.D., Davies P.M., Simmonds H.A., Leonard J.V.	45
A simplified method for the determination of phosphoribosylpyrophosphate synthetase activity in hemolysates Torres R.J., Mateos F.A., Puig J.G., Becker M.A.	55
Effects of alpha-1 acid glycoprotein on human polymorphonuclear neutrophils: Influence of glycan microheterogeneity Vasson M.P., Roch-Arveiller M., Couderc R., Baguet J.C., Raichvarg D.	65
Detection of viral DNA in hybridization reactions using two chemiluminescent substrates Girotti S., Ferri E., Musiani M., Gibellini D., Zerbini M., Roda A., Gentilomi G., Venturoli S.	73
Application of ¹ H-NMR spectroscopy for qualitative measurement of muscle carnitine levels Nakajima H., Sugie H., Tsurui S., Ito M.	81
Relationship of plasma malondialdehyde, vitamin E and antioxidant micronutrients to human immunodeficiency virus-1 seropositivity Malvy D.J.-M., Richard M.-J., Arnaud J., Favier A., Amedee-Manesme O.	89
Development of a radioimmunoassay for fetal antigen 2 Price K.M., Silman R., Armstrong P., Grudzinskas J.G.	95
<i>Letters to the Editor</i>	
Improvement of ELISA sensitivity by allogenic adsorption of polyclonal antibodies — A technical note for non-experts Abe Y., Kimura K., Horiuchi A., Miyake M., Kimura S.	103
Effect of short-term and long-term alcohol intake on plasma cholesterol in non-alcoholic humans Nakamura J.	107

Vol. 224/2, 1994

A new case of apo C-II deficiency with a nonsense mutation in the apo C-II gene Zanelli T., Catapano A.L., Aversa M.R., Barbagallo C.M., Liotta A., Giardina F.C., Notarbartolo A.	111
Analytical goals for rectilinear calibration functions Fuentes-Arderiu X., Alia P., Machuca I.	119
Relationship between lipoprotein(a) levels in serum and some indices of protein synthesis in liver cirrhosis Alessandri C., Basili S., Maurelli M., Andreozzi P., Violi F., Cordova C.	125
Genotyping compared with protein phenotyping the common apolipoprotein E polymorphism Hansen P.S., Gerdes L.U., Klausen I.C., Gregersen N., Faergeman O.	131
Serum octadeca-9,11 dienoic acid — An assay of free radical activity or a result of bacterial production? Jack C.I.A., Ridgway E., Jackson M.J., Hind C.R.K.	139
The esters of p-hydroxy-benzoate (parabens) inhibit the release of lysosomal enzymes by mitogen-stimulated peripheral human lymphocytes in culture Bairati C., Goi G., Lombardo A., Tettamanti G.	147
Therapeutic whole-blood levels of cyclosporin A in heart and lung transplantation Andert S.E., Grimm M., Schreiner W., Muller M.M.	159
Erythrocyte Na ⁺ ,K ⁺ -ATPase properties and adenylate energy charge in normotensives and in essential hypertensives Canestrari F., Galli F., Boschi S., Albertini M.C., Gheller G., De Crescentini S., Bosu M.	167
HPLC assay of conjugated bile acids in human fluids using on-line sample pretreatment on a standard isocratic chromatograph	

Scalia S., Pazzi P.	181
Light chains of immunoglobulins in human secretions	
Mole C.M., Bene M.C., Montagne P.M., Seilles E., Faure G.C.	191
Acute phase proteins and lipoprotein(a) in patients with severe hypercholesterolaemia and normal subjects	
Crook M.A., Haq S., Chusney G., Haq M., Tutt P.	199
Creatine kinase isoenzyme MB mass concentrations in patients undergoing aortocoronary bypass surgery	
Mair P., Mair J., Seibth I., Antretter H., Balogh D., Puschendorf B.	203
Tissue polypeptide specific antigen (TPS) in liver disease	
Moraglio D., Pagano M., Galeazzi D., Arnelli A., Bertero L.	209
<i>Letters to the Editor</i>	
The effect of silicon as silicic acid on superoxide dismutase activity	
Bellia J.P., Birchall J.D., Roberts N.B.	215
Concanavalin A affinity pattern of CA 19-9: An in vitro and in vivo study	
Basso D., Panozzo M.P., Del Giudice G., Carbone M.D., Meggiato T., Del Favero G., Piebani M.	219
The clinical laboratory is in the information business	
Weilert M.	51
Interpreting the results is the expertise of the laboratory	
Winkel P.	89
The role of suppliers and laboratories in the new information environment	
Korpman R.A.	S23

Vol. 225/1, 1994

Glycosylation of gamma-glutamyltransferase is modified by ethanol in H5-6 hepatoma cell line	
Odoul M., Bagrel D., Peyrieras N., Bodaud H., Siest G.	1
A solid phase enzyme linked immunofiltration assay for secretory leucocyte-proteinase inhibitor	
Melrose J., Ghosh P., Jacoby A.	17
GC-MS determination of ratios of stable-isotope labelled to natural urea using [$^{13}\text{C}^{15}\text{N}_2$]urea for studying urea kinetics in serum and as a means to validate routine methods for the quantitative assay of urea in dialysate	
Wolthers B.G., Tepper T., Withag A., Nagel G.T., de Haan T.H.Y., van Leeuwen J.J., Stegeman C.A., Huisman R.M.	29
ELISA for the core protein of the cartilage large aggregating proteoglycan, aggrecan: Comparison with the concentrations of immunogenic keratan sulphate in synovial fluid, serum and urine	
Moller H.J., Larsen F.S., Ingemann-Hansen T., Poulsen J.H.	43
Serum carnosinase activities in central nervous system disorders	
Wassif W.S., Sherwood R.A., Amir A., Idowu B., Summers B., Leigh N., Peters T.J.	57
Increased proteolysis of oxidatively damaged hemoglobin in erythrocyte lysates in diabetes mellitus	
Raghothama C., Rao P.	65
Increases in serum $\beta 3$ integrin in patients with chronic liver disease	
Yamauchi M., Mizuhara Y., Takahashi A., Toda G.	71
Phenotyping is an accurate means of analysing the principal apolipoprotein E isoforms	
James R.W., Ruiz J., Blanche H., Pometta D., Passa P., Froguel P.	77
<i>Letters to the Editor</i>	
The topography of thiol distribution on sperm surface: A sensitive marker for human fertility diagnosis	
Sinha S., Kumar G.P., Laloraya M.	83
Polymorphism at the glutathione s-transferase GSTM1 locus: A study of the frequencies of the GSTM1 a, b, A/B and null phenotypes in nigerians	
Zhao L., Aldersea J., Fryer A., Tighe A., Ollier B., Thomson W., Jones P., Strange R.	85

Enzyme reference materials: Their place in diagnostic enzymology	
Mois D.W.	S3
Systematic top-down approach to clinical chemistry	
Pardue H.L.	S13
Molecular biology techniques in the diagnosis of monogenic diseases	
Wagener C., Epplen J.T., Erlich H., Peretz H., Vihko P.	S35

Vol. 225/2, 1994

Elevated serum and urinary thrombomodulin levels in patients with non-insulin-dependent diabetes mellitus	
Shimano H., Takahashi K., Kawakami M., Gotoda T., Harada K., Shimada M., Yazaki Y., Yamada N.	89
Vitamin E and the hypercoagulability of neonatal blood	
Jain S.K., McCoy B., Wise R.	97
Fully mechanized latex immunoassay for serum lipoprotein(a)	
Abe A., Yoshimura Y., Sekine T., Maeda S., Yamashita S., Noma A.	105
Sex hormone binding globulin phenotypes: Their detection and distribution in healthy adults and in different clinical conditions	
Cornelisse M.M., Bennett P.E., Christiansen M., Blaakaer J., Gluud C., Andersen J.R., Homann C., Garred P.	115
Anti-thyroid peroxidase (anti-TPO) antibodies in thyroid diseases, non-thyroidal illness and controls. Clinical validity of a new commercial method for detection of anti-TPO (thyroid microsomal) autoantibodies	
Engler H., Riesen W.F., Keller B.	123
Monitoring of osteoblast activity with an immunoradiometric assay for determination of bone alkaline phosphatase mass concentration in patients receiving renal transplants	
Withold W., Degenhardt S., Castelli D., Heins M., Grabensee B.	137
Changes in purine nucleotide content in the lymphocyte subpopulations of patients infected with HIV	
Tabucchi A., Carlucci F., Consolmagno E., Monari P., Re M.C., Marinello E., Pizzichini M., Pagani R., Lai C.Y.	147
A rapid and sensitive sandwich enzyme immunoassay for low concentration of albumin in human urine	
Chiabrando G.A., Zalazar F.E., Aldao M.A.J., Vides M.A.	155
Feto-maternal plasma phenylalanine concentration gradient from 19 weeks gestation to term	
Schoonheydt W.E., Clarke J.T.R., Hanley A.B., Johnson J.M., Lehotay D.C.	165
Stability of human atrial natriuretic peptide in blood samples	
Tsuji T., Masuda H., Imagawa K., Haraikawa M., Shibata K., Kono M., Inouye K., Uchida K.	171
The glycosaminoglycan composition of human tracheas and the changes observed during aging and in disease	
Binette J.P., Burgi W., Ohishi H., Grundboeck-Jusko J., Burki R., Maekawa Y., Tachopp T.A., Kimura A., Schmid K.	179
1,25-Dihydroxyvitamin D radioreceptor assay using bovine mammary gland receptor non-high performance liquid chromatographic purification	
Watanabe Y., Kubota T., Suzumura E., Suzuki T., Yonezawa M., Ishigamia T., Ichikawa M., Seino Y.	187
Effect of storage of sera and apolipoprotein(a) phenotypes on detection of lipoprotein(a) by a new agarose gel	
Roberta B., Grandhomme M., Mainard F., Madec Y.	195

Vol. 226/1, 1994

Rapid and accurate random urinary porphyrin quantitation	
Boynton S.B., Roth K.S.	1

Serum levels of steroid sulfatase protein in gynecologic carcinomas	
Sugawara T., Honke K., Gasa S., Tanaka T., Fujimoto S., Makita A.	13
Quantitative analysis of Lewis antigens on erythrocytes by flow cytometry	
Murai J., Naka K., Shimojo N., Katakami T., Nakagishi M., Kuroki T., Okuda K.	21
Glycosaminoglycan content is increased in dissecting aneurysms of human thoracic aorta	
Cattell M.A., Hasleton P.S., Anderson J.C.	29
Clinical evaluation of biotin-binding immunoglobulin in patients with Graves' disease	
Nagamine T., Takehara K., Fukui T., Mori M.	47
Demonstration and some properties of N-acetyl- β -D-hexosaminidase (HEX) C isoenzyme in human renal tissues: Relative increase of HEX C activity in renal cell carcinoma	
Yoshida K.-I.	55
Sensitive colorimetric assay of serum diamine oxidase	
Takagi K., Nakao M., Ogura Y., Nabeshima T., Kunii A.	67
Rapid detection of common metachromatic leukodystrophy mutations by restriction analysis of arylsulfatase A gene amplimers	
Ben-Yoseph Y., Mitchell D.A.	77
Isolation of fetal antigen 2 and assay standardisation	
Price K.M., Silman R., Grudzinskas J.G.	83
Rapid assay for furancarboxylic acid accumulated in uremic serum using high-performance liquid chromatography and on-line mass spectrometry	
Niwa T., Kawagishi I., Ohya N.	89
A rapid semi-micro method for the separation of lipoprotein fractions that uses a benchtop ultracentrifuge	
Fletcher C.D., Barnes J.F., Farish E.	95
<i>Letters to the Editor</i>	
Suitability of copper chloride as a reaction terminator for superoxide dismutase activity assay	
Sun Y., Oberley L.W.	101
Lipoprotein(a) concentration in various hyperlipidemias	
Alberty R.	105
Automated kinetic assay for endotoxin and (1 \rightarrow 3)- β -D-glucan in human blood	
Tamura H., Arimoto Y., Tanaka S., Yoshida M., Obayashi T., Kawai T.	109
Increased levels of plasma endothelin-1 in patients with psoriasis	
Cecchi R., Giomi A., Ciuti M., Gironi A., Seghieri G.	113
Increasing the biosafety of analytical systems in the clinical laboratory	
Truchaud A., Schnipelsky P., Pardue H.L., Place J., Ozawa K.	55

Vol. 226/2, 1994

Human prostatic acid phosphatase: Selected properties and practical applications	
Ostrowski W.S., Kuciel R.	121
Release of membrane-bound enzymes from cells and the generation of isoforms	
Moss D.W.	131
Bone-origin alkaline phosphatase in plasma by wheat-germ lectin methods in bone disease	
Rosalki S.B.	143
Monitoring skeletal cancer metastases with the bone isoenzyme of tissue unspecific alkaline phosphatase	
Burlina A., Rubin D., Secchiario S., Sciacovelli L., Zaninotto M., Plebani M.	151
Antenatal diagnosis of tetrahydrobiopterin deficiency by quantification of pterins in amniotic fluid and enzyme activity in fetal and extrafetal tissue	
Blau N., Kierat L., Matasovic A., Leimbacher W., Heizmann C.W., Guardamagna O., Ponzone A.	159
Drug metabolizing enzymes related to laboratory medicine: Cytochromes P-450 and UDP-glucuronosyltransferases	
Batt A.M., Magdalou J., Vincent-Viry M., Ouzzine M., Fournel-Gigleux S., Galteau M.M., Siest G.	171

Enzymes and circulating proteins as markers of alcohol abuse	
Goldberg D.M., Kapur B.M.	191
The enzymes of mitochondrial fatty acid oxidation	
Bennett M.J.	211
Detection of point mutations in human genes by the solid-phase minisequencing method	
Syvanen A.-C.	225
Viral diagnosis using hybridization assays with digoxigenin labeled probes	
Musiani M., Zerbini M., Gibellini D., Venturoli S., Gentilomi G., Gallinella G., La Placa M.	237
Screening for cystic fibrosis mutations in Spanish patients	
Molano J., Ezquieta B., Granell R.	247

Vol. 227/1-2, 1994

Inflammation-associated changes in the cellular availability of tryptophan and kynurenine in renal transplant recipients	
Holmes E.W., Russell P.M., Kinzler G.J., Bermes Jr. E.W.	1
Initiation codon mutation of the tyrosinase gene as a cause of human albinism	
Breimer L.H., Winder A.F., Jay B., Jay M.	17
Neurotrophin-3 (NT-3) levels in the developing rat nervous system and in human samples	
Murase K., Igarashi K., Hayashi K.	23
Role of the measurement of serum procollagen type III N-terminal peptide in the evaluation of liver diseases	
Collazos J., Diaz F.	37
Factors affecting the thiobarbituric acid test as index of red blood cell susceptibility to lipid peroxidation: A multivariate analysis	
Girelli D., Olivieri O., Stanzial A.M., Guarini P., Trevisan M.T., Bassi A., Corocher R.	45
Serum beta-hexosaminidase isoenzymes are related to risk factors for atherosclerosis in a large population of postmenopausal women	
Hultberg B., Isaksson A., Nilsson J.A., Lindgarde F.	59
Monosaccharide composition of haptoglobin in liver diseases and alcohol abuse: Large changes in glycosylation associated with alcoholic liver disease	
Mann A.C., Record C.O., Self C.H., Turner G.A.	69
An HPLC-linked assay of phosphoribosylpyrophosphate synthetase activity in the erythrocytes of adults and children with neurological disorders	
Micheli V., Rocchigiani M., Pompucci G.	79
Biological markers and ovarian carcinomas: Galactosyltransferase, CA 125, isoenzymes of amylase and alkaline phosphatase	
Sichel F., Salaun V., Bara E., Gauduchon P., Malas J.-P., Goussard J., Le Talaer J.-Y.	87
Acidification reveals a greater number of epidermal growth factor receptors in human placental and breast cancer membranes	
Hatzoglou A., Romain S., Dalapascha H., Castanas E.	97
Separation of alkaline phosphatase isoforms with and without intact glycan-phosphatidylinositol anchors in aqueous polymer phase systems	
Raymond F.D., Moss D.W., Fisher D.	111
[¹⁴ C]Methylamine accumulation in cultured human skin fibroblasts a biochemical test for lysosomal storage and lysosomal diseases	
Kopitz J., Gerhard C., Hofler P., Cantz M.	121
Plasma fibronectin and angiotensin-converting enzyme: Markers of primary pulmonary injury in burn patients	
Plassart F., Cynober L., Baudin B., Lioret N., Beneteau-Burnat B., Nolland X.B., Saizy R., Giboudeau J.	135
Optimized determination of angiotensin I-converting enzyme activity with hippuryl-L-histidyl-L-leucine as substrate	
Schnaith E., Beyrau R., Buckner B., Klein R.M., Rick W.	145

A rapid and simple method for measuring the susceptibility of low-density-lipoprotein and very-low-density-lipoprotein to copper-catalyzed oxidation Zhang A., Vertommen J., van Gaal L., De Leeuw I.	159
Comparison of enhanced chemiluminescence and colorimetric techniques for the immune-detection of α -antitrypsin Venembre P., Seta N., Boutten A., Dehoux M., Aubier M., Durand G.	175
Use of HPLC to measure circulating adenosine levels in migrainous patients Guieu R., Sampieri F., Bechis G., Rochat H.	185
Relationship between ulinastatin and alpha-1-microglobulin in human urine Shikimi T., Himeno Y., Shigeno K., Gonda T., Ishibe T., Hattori K., Takaori S.	195
Measurement of the oxidisable polyamine content of seminal plasma Fernandez C., Sharrard R.M., Monks N., Barrett C.L.R.	201
<i>Letters to the Editor</i>	
Detection of modified β_2 -microglobulin (β_2 m) from amyloid deposits in tenosynovial tissue of carpal tunnel syndrome (CTS) Odani H., Mizutani A., Hidaka H., Fujisawa K., Shinzato T., Maeda K.	209
Examination of bronchoalveolar wash fluid for ras, myc and fos oncoproteins and glutathione S-transferase isoenzymes: An attempt to improve the accuracy of lung cancer diagnosis Pinkham J.M., Strange R.C., Pantin C.	211

Vol. 228/1, 1994

ClinBio 92' and 93': Introduction Salvatore F., Pane F.	1
Greater differences in forensic DNA profile frequencies estimated from racial groups than from ethnic subgroups Budowle B., Monson K.L.	3
Magnetic resonance imaging and spectroscopy of MCF7 human breast cancer: Pathophysiology and monitoring of treatment Degani H., Furman E., Fields S.	19
Biochemical and molecular investigations in respiratory chain deficiencies Rustin P., Chretien D., Bourgeron T., Gerard B., Rotig A., Saudubray J.M., Munnich A.	35
Reference charts for respiratory chain activities in human tissues Chretien D., Rustin P., Bourgeron T., Rotig A., Saudubray J.M., Munnich A.	53

Vol. 228/2, 1994

Estimation of pS2 protein level in human body fluids by a sensitive two-site enzyme immunoassay Miyashita S., Nomoto H., Konishi H., Hayashi K.	71
A detection method for point mutation in the precore region of human hepatitis B virus (HBV)-DNA using mutation-site-specific assay Kinoshita M., Seno T., Fukui T., Shin S., Tsubota A., Kumada H.	83
Elevation of serum group II phospholipase A_2 levels in patients with advanced cancer Yamashita S., Ogawa M., Sakamoto K., Abe T., Arakawa H., Yamashita J.	91
Evaluation of tumor-associated antigen (2H6 antigen) in detecting early stages of gastric cancer Harada H., Tsukada Y., Karasawa Y.	101
Thyroid hormone autoantibodies in patients with Graves' disease: Effect of anti-thyroid drug treatment Ogawa T., Sakata S., Nakamura S., Takuno H., Matsui I., Sarui H., Yasuda K.	113
Tubular nephrotoxicity after cardiac surgery utilising cardiopulmonary bypass Westhuyzen J., McGiffin D.C., McCarthy J., Fleming S.J.	123
A new precipitation method with magnetic separation for high-density-lipoprotein cholesterol assay Iida S., Osawa S., Yonemitsu H.	133

Cathepsin B and N-acetyl- β -D-glucosaminidase in human synovial cells in culture: Effects of interleukin-1	
Lecomte V., Knott I., Burton M., Remacle J., Raes M.	143
Modulation defect of sodium pump evidenced in diabetic patients by a microcalorimetric study	
Issautier T., Kovacic H., Gallice P., Raccach D., Vague P., Crevat A.	161
Lipoprotein profiling by high performance gel chromatography	
Tallis G.A., Shephard M.D.S., Whiting M.J.	171
Interaction of β -cyclodextrin with bile acids and their competition with vitamins A and D3 as determined by $^1\text{H-NMR}$ spectrometry	
Comini S., Olivier P., Riottot M., Duhamel D.	181
Characterization of tumor-associated antigen (2H6 Antigen) defined by monoclonal antibody 2H6	
Harada H., Kubota T., Tsukada Y., Karasawa Y.	195
Serum type IV collagen and prolyl hydroxylase levels: Effect of gender and age	
Tamaro G., Moretti M., Marchi P., Mangiarotti M., Pozzato G.	205

Vol. 229/1-2, 1994

The correlation coefficient in the interpretation of laboratory data	
Chinchilli V.M., Gruemer H.-D.	1
Early changes of serum N-acetyl- β -glucosaminidase, tissue plasminogen activator and erythrocyte superoxide dismutase in relation to retinopathy in type 1 diabetes mellitus	
Skrha J., Hodinar A., Kvasnicka J., Stibor V., Sperl M., Stolba P., Hilgertova J.	5
Nicotinic acid phosphoribosyltransferase activity in human erythrocytes: Studies using a new HPLC method	
Pescaglini M., Micheli V., Simmonds H.A., Rocchigiani M., Pompucci G.	15
The utilization of intravenously infused pyridoxine in humans	
Zempleni J., Kubler W.	27
Determination of trace amounts of albumin in human bronchoalveolar lavage fluids by fluorometry with chromazurol S	
Sato T., Oh'ue M., Nakarai T., Ichikawa K., Kitagishi K., Murakami N., Chikuma M., Saito Y., Kado M., Nagai S., Izumi T.	37
A processing-independent assay for human procholecystokinin and its products	
Paloheimo L.L., Rehfeld J.F.	49
Quantitative computer image analysis of a human chorionic gonadotropin colloidal carbon dipstick assay	
Van Amerongen A., Van Loon D., Berendsen L.B.J.M., Wichers J.H.	67
Precipitation procedures used to isolate high density lipoprotein with particular reference to effects on apo A-I-only particles and lipoprotein(a)	
Vaillance D.T., Byrne D.J., Winder A.F.	77
Differential processing of type I and type III procollagens in the tumour cysts and peritoneal ascitic fluid of patients with benign and malignant ovarian tumours	
Zhu G.-G., Melkko J., Risteli J., Kauppila A., Risteli L.	87
Measurement of short-chain acyl-CoA dehydrogenase (SCAD) in cultured skin fibroblasts with hexanoyl-CoA as a competitive inhibitor to eliminate the contribution of medium-chain acyl-CoA dehydrogenase	
Niezen-Koning K.E., Wanders R.J.A., Nagel G.T., Sewell A.C., Heymans H.S.A.	99
Estradiol receptors in combination with neu or myc oncogene amplifications might define new subtypes of breast cancer	
Bolufer P., Molina R., Ruiz A., Hernandez M., Vazquez C., Lluch A.	107
Quantitative and qualitative changes of extracellular-superoxide dismutase in patients with various diseases	
Adachia T., Nakamura M., Yamada H., Futenma A., Kato K., Hirano K.	123
L-Homoserine hydroxamic acid as an antitumor agent	
Natelson S., Pantazis P., Natelson E.A.	133

Semi-quantitative latex method for lipoprotein(a) assay	
Lippi G., Ruzzenente O., Facchinetti R., Guidi G.	147
Cardiac myosin light chain-1 release in acute myocardial infarction is associated with scintigraphic estimates of myocardial scar	
Mair J., Wagner I., Fridrich L., Lechleitner P., Diestl F., Puschendorf B., Michel G.	153
A combination assay of MB and MM isoforms of serum creatine kinase in acute myocardial infarction	
Kanemitsu F., Okigaki T.	161
Determination of amyloid type by ELISA using milligram amounts of tissue	
Kaplan B., German G., Ravid M., Pras M.	171
Simultaneous detection of $\Delta F508$, G542X, N1303K and 1717-1G—A mutations in cystic fibrosis by capillary electrophoresis in polymer networks	
Gelfi C., Righetti P.G., Magnani C., Cremonesi L., Ferrari M.	181
Serum hyaluronate levels predict responsiveness to interferon alpha therapy in patients with chronic hepatitis C	
Matsunaga T., Kojima J., Inoue A., Matsumiya K.	191
A simple non-radioactive assay for hemoglobin E gene in prenatal diagnosis	
Fucharoen S., Fucharoen G., Ratanasiri T., Jetsrisuparb A., Fukumaki Y.	197
<i>Letter to the Editor</i>	
Serum serotonin, lactate and pyruvate levels in infantile autistic children	
Laszlo A., Horvath E., Eck E., Fekete M.	205

Vol. 230/1, 1994

The detection of 3-methylglutaryl carnitine and a new dicarboxylic conjugate, 3-methylglutacylcarnitine, in 3-methylglutaconic aciduria	
Jocste S., Erasmus E., Mienie L.J., De Wet W.J., Gibson K.M.	1
Carbohydrate-mediated recognition of a circulating placental alkaline phosphatase-immunoglobulin M complex	
Koyama I., Sato M., Sakai H., Nagata A., Miura M., Sakagishi Y., Komoda T.	9
Determinants of serum 1,25-dihydroxyvitamin D concentration in healthy premenopausal subjects	
Ho S.C., Mac Donald D., Chan C., Fan Y.K., Chan S.S.G., Swaminathan R.	21
Vitamin, trace element and peroxide status in HIV seropositive patients: Asymptomatic patients present a severe β -carotene deficiency	
Sappey C., Leclercq P., Coudray C., Faure P., Micoud M., Favier A.	35
A screening method for abnormally high lipoprotein(a) concentrations by agarose lipoprotein electrophoresis	
Campos E., Fievet P., Caces E., Fruchart J.-C., Fievet C.	43
Copper,zinc-superoxide dismutase and hydrogen peroxide: A hydroxyl radical generating system	
Koningsberger J.C., Van Asbeck B.S., Van Faassen E., Wiegman L.J.J.M., Van Hattum J., Van Berge Henegouwen G.P., Marx J.J.M.	51
The use of HPLC to evaluate the variations of blood coronary adenosine levels during percutaneous transluminal angioplasty	
Guieu R., Paganelli F., Sampieri F., Bechis G., Levy S., Rochat H.	63
Abnormal glycine betaine content of the blood and urine of diabetic and renal patients	
Lever M., Sizeland P.C.B., Bason L.M., Hayman C.M., Robson R.A., Chambers S.T.	69
A competitive dual-label time-resolved immunofluorometric assay for simultaneous detection of carbonic anhydrase I and II in cerebrospinal fluid	
Parkkila A.-K., Parkkila S., Serlo W., Reunanen M., Vierjoki T., Rajenieni H.	81
Urinary excretion of homocitric acid and methylhomocitric acid in propionic acidemia: Minor metabolic products of the citrate synthase aldol condensation reaction	
Van Rooyen J.P.G., Mienie L.J., Erasmus E., De Wet W.J., Duran M., Wadman S.K.	91
Incidence of the apolipoprotein B-3500 mutation in Denmark	
Hansen P.S., Norgaard-Petersen B., Meinertz H., Jensen H.K., Hansen A.B.B., Klausen I.C., Gerdes L.U., Faergeman O.	101

Letter to the Editor

- Effect of a 6-month storage on human serum fructosamine concentration [1]
 Balland M., Schiele F., Henny J. 105

Vol. 230/2, 1994

- Cystic fibrosis and phosphatidylcholine biosynthesis
 Ulane M.M., Butler DeB. J., Peri A., Miele L., Ulane R.E., Hubbard V.S. 109
- Serum neurone-specific enolase concentrations in patients with neurological disorders
 Cunningham R.T., Morrow J.I., Johnston C.F., Buchanan K.D. 117
- Radioimmunoassay of vasopressin during pregnancy. Use and removal of cystylaminopeptidase inhibitors
 Van der Post J.A., Van Heerikhuize J.J., Boer K., Van Boxtel C.E., Swaab D.F. 125
- Binding of aluminium to plasma proteins: Comparative effect of desferrioxamine and deferiprone (L1)
 Fernandez-Martin J.L., Menendez-Fraga P., Canteros M.A., Diaz-Lopez J.B., Cannata-Andia J.B. 137
- Longitudinal changes in serum zinc concentration and distribution after acute myocardial infarction
 Arnaud J., Faure H., Bourlard P., Denis B., Favier A.E. 147
- Naftidrofuryl inhibits the release of 5-hydroxytryptamine and platelet-derived growth factor from human platelets
 Barradas M.A., Jagroop I.A., Mikhailidis D.P. 157
- Simultaneous determination of betaine and N,N-dimethylglycine in urine
 Laryea M.D., Zass R., Ritgen J., Wendel U. 169
- Development and evaluation of a spectrophotometric assay for complex III in isolated mitochondria, tissues and fibroblasts from rats and humans
 Krahenbuhl S., Talos C., Wiesmann U., Hoppel C.L. 177
- Heightened free radical activity in blacks with chronic pancreatitis at Johannesburg, South Africa
 Gut A., Shiel N., Kay P.M., Segal I., Braganza J.M. 189
- A microassay for urinary phenol using capillary gas chromatography and optimised enzymic hydrolysis
 Ahmed N., Hale K. 201
- Is there a need for creatinine adjustment of urinary steroid hormone levels in studies of early fetal loss?
 Hakim R.B., Gray R.H., Zacur H.A. 209
- Separation of lipoprotein fractions by ultracentrifugation: Investigation of analytical recovery with sequential flotation and density gradient procedures
 Potts J.L., Fisher R.M., Humphreys S.M., Gibbons G.F., Frayn K.N. 215
- Letter to the Editor*
- Sulfadiazine/N⁶-acetylsulfadiazine crystalluria in a patient with the acquired immune deficiency syndrome (AIDS)
 Potter J.L., Kofron W.G. 221

Vol. 231/1, 1994

- Characterization of human creatine kinase BE and MB isoforms by means of isoelectric focusing
 Kanemitsu F., Okigaki T. 1
- Crystallization of urinary calcium oxalate at standardized osmolality and pH in the frozen state
 Gohel M.D.I., Shum D.K.Y., Li M.K. 11
- Trace elements in the sweat of acclimatized persons
 Omokhodion F.O., Howard J.M. 23
- Plasma cholesteryl ester transfer in patients with non-insulin dependent diabetes mellitus
 Sutherland W.H., Walker R.J., Lewis-Barned N.J., Pratt H., Tillman H.C. 29

Urinary selenium concentrations in heroin abusers	
Rodriquez Rodriguez E.M., Sanz Alaejos M., Diaz Romero C.	39
Different time courses of cardiac contractile proteins after acute myocardial infarction	
Mair J., Wagner I., Jakob G., Lechleitner P., Dienstl F., Puschendorf B., Michel G.	47
Regulation of the mitochondrial ATP-synthase in human fibroblasts	
Das A.M., Byrd D.J., Brodehl J.	61
Excess matrix accumulation in scleroderma is caused partly by differential regulation of stromelysin and TIMP-1 synthesis	
Bou-Gharios G., Osman J., Black C., Olsen I.	69
A one-step sandwich enzyme immunoassay for inactive precursor and complexed forms of human matrix metalloproteinase 9 (92 kDa gelatinase/type IV collagenase, gelatinase B) using monoclonal antibodies	
Fujimoto N., Hosokawa N., Iwata K., Shinya T., Okada Y., Hayakawa T.	79
Effect of storage at -70°C on immunoturbidimetric assays for apolipoproteins A-I and B	
Myron Johnson A., Ledue T.B., Palomaki G.E., Craig W.Y.	89
Red blood cell glutathione status in patients with gastrointestinal malignancies treated with 5-fluorouracil	
Walach N., Evans S., Zaidman J.L., Cotariu D.	95
Increased angiotensin-converting enzyme activity in cerebrospinal fluid of treated patients with Parkinson's disease	
Konings C.H., Kuiper M.A., Bergmans P.L.M., Grijpma A.M., van Kamp G.J., Wolters Ch. E.	101
Measurement of steroid hormones in plasma by isocratic high performance liquid chromatography coupled to radioimmunoassay	
Boschi S., De Iasio R., Mesini P., Bolelli G.F., Sciajno R., Pasquali R., Capelli M.	107
<i>Letter to the Editor</i>	
Usefulness of the kappa/lambda ratio for detecting a new monoclonal component in a patient with a monoclonal gammopathy	
Myara I., Asso M., Tenenhaus D., Moatti N.	115

Vol. 231/2, 1994

An improved bromide assay for the estimation of extracellular water volume by capillary gas chromatography autovan	
Kreel B.K.	117
Routine high-performance liquid chromatographic determination of urinary unconjugated cortisol using solid-phase extraction and ultraviolet detection	
Dolezalova M.	129
Immunoheterogeneity of parathyroid hormone after surgery for primary hyperparathyroidism	
Bergenfels A., Valdermarsson S., Ahren B.	139
Competitive enzyme-linked immunosorbent assay of the human cholesteryl ester transfer protein (CETP)	
Guyard-Dangremont V., Lagrost L., Gambert P., Lallemand C.	147
Hemoglobins S and C: Reference values for glycohemoglobin in heterozygous, double-heterozygous and homozygous subjects, as established by 13 methods	
Weykamp C.W., Martina W.V., Van der Dijs F.P.L., Penders T.J., Van der Slik W., Muskiet F.A.J.	161
Gamma-type gliadins cause secretion of prostaglandin E ₂ in patients with coeliac disease	
Friis S., Anthonen D., Noren O., Sjoström H.	173
Evaluation of enzyme inhibitors of cystinyl aminopeptidase and application to the measurement of immunoreactive atrial natriuretic peptide in human pregnancy	
Irons D.W., Davison J.M., Baylis P.H.	185
Improved method for quantitative analysis of lung surfactant phospholipids in bronchoalveolar lavage fluids by high-performance liquid chromatography	
Bruch J., Gono E., Malkusch W., Rehn B.	193

Vol. 232/1-2, 1994

Clinical chemistry as scientific discipline: Historical perspectives Buttner J.	1
Training and continuous education of clinical laboratory technologists and technicians Mocarelli P.	11
The impact of clinical biochemistry on the pre-doctoral medical curriculum: An Italian viewpoint Burlina A.	23
Post-doctoral training of the non-physician doctoral scientist Schwartz M.K.	33
Teaching and training of clinical biochemists: Sub-specialization Rock R.C.	39
Continuing clinical chemistry education in the United States Conn R.B.	47
Clinical chemistry education in the United States Tetrault G.A., Gruemer H.-D.	53
Education in clinical biochemistry: The Canadian scene Bunting P.S., Goldberg D.M.	63
Training and education in clinical biochemistry in the United Kingdom Price C.P., Dryburgh F.J., Elder G.H.	85
Education and clinical biochemistry in Australia O'Leary T.D., Guerin M.D.	97
Education in clinical chemistry in Germany Delbruck A.F.	109
Education in clinical chemistry and laboratory sciences in Austria Muller M.M.	119
Teaching clinical chemistry in central European countries — past and present Naskalski J.W., Palicka V.	125
Clinical biochemistry training in the Netherlands Sanders G.T.B., Brombacher P.J.	133
The impact of clinical biochemistry on university education in France Masseyeff R.F., Dreux C., Goussault Y.	143
Clinical biochemistry education in Spain Queralto J.M.	153
Education in clinical chemistry in Japan Kanno T.	159
Training of clinical laboratory professionals in Russia Morozova V.T., Dolgov V.V., Malakhov V.N.	167

Vol. 233/1-2, 1995

Appropriate utilization and cost control of the hospital laboratory: Panel testing and repeat orders Werner M.	1
Multi-enzyme reference material from established human cell lines and human sources Marui Y., Hayashi C., Matsuda Y., Matsuura S., Eto A., Ohba Y., Okuda K.	19
Biotinylation of biotinidase following incubation with biocytin Hymes J., Fleischauer K., Wolf B.	39
The influence of the blood levels of lead, aluminum and vanadium upon the arterial hypertension Granadillo V.A., Tahan J.E., Salgado O., Elejalde L.E., Rodriguez-Iturbe B., Romero G.B., Romero R.A.	47
Role of monocyte/macrophage derived matrix-metalloproteinases (gelatinases) in prolonged skin inflammation Maillard J.L., Favreau C., Reboud-Ravaux M.	61
Hungarian hereditary acatalasemia and hypocatalasemia are not associated with chronic hemolysis Goth L., Vitai M.	75

Dimethylamine in human urine	
Zhang A.Q., Mitchell S.C., Smith R.L.	81
A preliminary report on urinary BT-PABA/PAS excretion index, serum pancreatic isoamylase and faecal chymotrypsin tests of pancreatic dysfunction in Sowetan Africans	
Chaloner C., Segal I., Hassan H., McIntosh J., Gut A., Rahme M., Braganza J.M.	89
Estimation of the value of serum β_2 -microglobulin concentration in the diagnosis of Hashimoto's disease	
Kochanska-Dziurawicz A., Starzewski J.J., Dunal W.	101
Electrolyte balanced heparin may produce a bias in the measurement of ionized calcium concentration in specimens with abnormally low protein concentration	
Lyon M.E., Guajardo M., Laha T., Malik S., Henderson P.J., Kenny M.A.	105
Transient hyperphosphatasemia in infancy associated with an increased urinary hydroxyproline excretion	
Stepan J.J., Kutilek S., Bayer M.	115
Cholestyramine orally administered to patients with erythropoietic protoporphyria results in urinary excretion of protoporphyrin: To be explained by the Herbst-Volkheimer effect?	
Beukeveld G.J.J., Wolthers B.G.	119
Sialyl salivary-type amylasemia associated with immunoglobulin D-type multiple myeloma	
Moriyama T., Tozawa T., Nobuoka M., Ikeda H.	127

Vol. 234/1-2, 1995

Detection of the three common mutations of adenosinephosphoribosyltransferase deficiency among Japanese	
Higashimoto H., Ouchi A., Kawaguchi R.	1
Differentiation between organophosphate and carbamate poisoning	
Rotenberg M., Shefi M., Dany S., Dore I., Tirosh M., Almog S.	11
Tissue accumulation of sulfatide and G(M3) ganglioside in a patient with variant Farber disease	
Fujiwaki T., Hamanaka S., Tate S., Inagaki F., Suzuki M., Suzuki A., Mori C.	23
Immunochemical detection of urinary 5-methyl-2'-deoxycytidine as a potential biologic marker for leukemia	
Itoh K., Aida S., Ishiwata S., Yamaguchi T., Ishida N., Mizugaki M.	37
Reactivity of lecithin-cholesterol acyl transferase (LCAT) towards glycated high-density lipoproteins (HDL)	
Fournier N., Myara I., Atger V., Moatti N.	47
Measurement of serum lipid peroxidation during exercise using three different methods: Diene conjugation, thiobarbituric acid reactive material and fluorescent chromolipids	
Vasankari T., Kujala U., Heinonen O., Kapanen J., Ahotupa M.	63
An HPLC assay for sn-1,2-diacylglycerol	
Eaton S., Shmueli E., Al-Mardini H., Bartlett K., Record C.O.	71
An improved method for quantification of extra domain A-containing cellular fibronectin (EDAcFN) in different body fluids	
Ylatupa S., Mertaniemi P., Haglund C., Partanen P.	79
Comparison of erythrocyte transketolase activity with thiamine and thiamine phosphate ester levels in chronic alcoholic patients	
Herve C., Beyne P., Letteron Ph., Delacoux E.	91
Galactosyl hydroxylysine in assessment of Paget's bone disease	
Michalsky M., Stepan J.J., Wilczek H., Formankova J., Moro L.	101
Evaluation of oxidant stress in chronic hemodialysis patients: Use of different parameters	
Haklar G., Yegenaga I., Yalcin A.S.	109
One-step sandwich enzyme immunoassays for human C4b-binding protein (C4BP) and protein S-C4BP complex using monoclonal antibodies	
Tamei H., Hoshino T., Yoshida S., Hayashi T., Iwata K., Suzuki K.	115
Redox state, antioxidative activity and lipid peroxidation in erythrocytes and plasma of chronic ambulatory peritoneal dialysis patients	

Canestrari F., Buoncristiani U., Galli F., Giorgini A., Albertini M.C., Carobi C., Pascucci M., Bossu M.	127
Effect of double-blind cross-over selenium supplementation on lipid peroxidation markers in cystic fibrosis patients	
Portal B., Richard M.-J., Coudray C., Arnaud J., Favier A.	137
Quantitative determination of faecal fat, nitrogen and water by means of a spectrophotometric technique: Near infrared reflectance analysis (NIRA). Assessment of its accuracy and reproducibility compared with chemical methods	
Picarelli A., Greco M., Di Giovambattista F., Ramazzotti A., Cedrone C., Corazziari E., Torsoli A.	147
Ion-selective electrode for serum bromide assay in patients with epilepsy	
Katsu T., Furuno K., Yamashita S., Kawasaki H., Gomita Y., Ohtsuka Y., Ohtahara S.	157
Total renin in pre-eclampsia	
Matinlauri I., Ekblad U., Maenpaa K., Koskinen P., Aalto M., Peltola O., Irjala K.	163
Clinical usefulness of cimetidine treatment for acute relapse in intermittent porphyria	
Horie Y., Norimoto M., Tajima F., Sasaki H., Nanba E., Kawasaki H.	171
<i>Letter to the Editor</i>	
Determination of α -amylase using a new blocked substrate (2-chloro-4-nitrophenyl 4 ⁴ -O- β -D-galactopyranosyl- β -maltotetraoside)	
Majima K., Teshima S., Hamada Y., Kikuchi T., Kawamura Y., Kitahata S.	177

Vol. 235/1, 1995

A scheme for the interpretation of primary and secondary disturbances of plasma and urinary amino acid profiles. A possible way to an expert system	
Parvy P., Bardet J., Rabier D., Kamoun P.	1
Active and acid-activatable TGF- β in human sera, platelets and plasma	
Grainger D.J., Mosedale D.E., Metcalfe J.C., Weissberg P.L., Kemp P.R.	11
Serum pyridinolines as specific markers of bone resorption in hemodialyzed patients	
Niwa T., Shiobara K., Hamada T., Miyazaki T., Tsukushi S., Uema K., Tsuzuki T.	33
Genetic basis of the silent phenotype of serum butyrylcholinesterase in three compound heterozygotes	
Maekawa M., Sudo K., Kanno T., Kotani K., Dey D.C., Ishikawa J., Izumi M., Etoh K.	41
Immunoturbidimetric determination of lipoprotein(a): Improvement in the measurement of turbid and triglyceride-rich samples	
Molinari E.A., Pichler P.F., Grillhofer H., Kostner G.M.	59
Alpha-fetoprotein and the acute phase response. A study using acute pelvic inflammatory disease as a model system	
Christiansen M., Hogdall C.K., Brihmer C.	71
A simple and automated HPLC method for determination of total hydroxyproline in urine. Comparison with excretion of pyridinolines	
Castelain S., Kamel S., Picard C., Desmet G., Sebert J.L., Brazier M.	81
Comparison of micro-enzymatic and high-performance liquid chromatographic methods for the assay of serum 1,5-anhydroglucitol	
Chusney G.D., Philippa M., Pickup J.C.	91
CSF acetylcholinesterase in Parkinson disease: Decreased enzyme activity and immunoreactivity in demented patients	
Konings C.H., Kuiper M.A., Mulder C., Calliauw J., Wolters Ch. E.	101
Cholinesterase activities in cerebrospinal fluid of patients with idiopathic convulsive disorders	
Basu P.S., Databyal S.K., Bhattacharya A., Datta T.K.	107

Vol. 235/2, 1995

Radiochemical assay of minute quantities of galactose-1-phosphate uridylyltransferase activity in

erythrocytes and leukocytes of galactosemia patients Xu Y.-K., Kaufman F.R., Donnell G.N., Ng W.G.	125
A sandwich enzyme immunoassay for the determination of neutrophil lipocalin in body fluids Blaser J., Triebel S., Tachesche H.	137
Blood antioxidants (vitamin E and β -carotene) in long-term low density lipoprotein apheresis Assogha U., Lepage S., Bruckert E., Bonnefont-Rousselot D., Dairou F., De Gennes J.L., Delattre J.	147
Changed transferrin sialylation in Parkinson's disease Van Kamp G.J., Mulder K., Kuiper M., Wolters Ch. E.	159
Enzymatic determination of bicarbonate: In serum by flow injection analysis Quiles R., Fernandez-Romero J.M., De Castro M.D.L.	169
Oxidative stress and erythrocyte membrane fluidity in patients undergoing regular dialysis McGrath L.T., Douglas A.F., McClean E., Brown J.H., Doherty C.C., Johnston G.D., Archbold G.P.R.	179
Serum lipoprotein profile in Algerian patients with celiac disease Mediene S., Hakem S., Bard J.M., Medjaoui I., Benhamamouch S., Lebel P., Fruchart J.C., Clavey V.	189
A specific immunoradiometric assay of cationic trypsin(ogen) that does not recognize trypsin-alpha-1-proteinase inhibitor complex Lafont P., Guy-Crotte O., Paulin C., Galvain D., Mertani S., Figarella C., Barbier Y.	197
The red wine phenolics trans-resveratrol and quercetin block human platelet aggregation and eicosanoid synthesis: Implications for protection against coronary heart disease Pace-Asciak C.R., Hahn S., Diamandis E.P., Soleas G., Goldberg D.M.	207
Assay of serum pyridinoline: A potential marker for bone resorption Hata K., Miura M., Fukumoto S., Matsumoto T.	221
Surfactant associated protein A determination using a chemiluminescence system — Application to tracheal aspirates from newborns Taieb J., Francoual J., Magny J.-F., Fraslon d C., Messaoudi C., Lindenbaum A., Bourbon J.	229

Vol. 236/1, 1995

Calcium and magnesium in human toenails do not reflect bone mineral density Vecht-Hart C.M., Bode P., Trouerbach Th. W., Collette H.J.A.	1
Human triacylglycerol-rich lipoprotein subfractions as substrates for lipoprotein lipase Fisher R.M., Coppack S.W., Humphrey S.M., Gibbons G.F., Frayn K.N.	7
Urinary creatinine adjusted reference ranges for homovanillic and vanillylmandelic acid in children and adults Cole M., Craft A.W., Parker L., Bell S., Seviour J.A., McGill A.C., Dale G.	19
Urinary steroid profile of a newborn suffering from pseudohypoadosteronism Wolthers B.G., Kraan G.P.B., Van der Molen J.C., Nagel G.T., Rouwe C.W., Lenting F.	33
Diagnosis of Smith-Lemli-Opitz syndrome by gas chromatography/mass spectrometry of 7-dehydrocholesterol in plasma, amniotic fluid and cultured skin fibroblasts Kelley R.I.	45
A sandwich ELISA based on anti-apo(a) and anti-apo B monoclonal antibodies for lipoprotein(a) measurement Sorell L., Rojas G., Rodriguez M., Ramos C., Torres L., Torres M.B.	59
Clinical evaluation of a monoclonal antibody to serum KM01 for the diagnosis of hepatocellular carcinoma Fujii T., Horie Y., Ikuta Y., Nishimuki E., Murawaki Y., Suou T., Kawasaki H.	71
Genetic analysis of Japanese patients with myophosphorylase deficiency (McArdle's disease): Single-codon deletion in exon 17 is the predominant mutation Sugie H., Sugie Y., Ito M., Fukuda T., Nonaka I., Igarashi Y.	81
Within- and between-subject variation in, and associations between, serum concentrations and urinary excretion of testosterone and estradiol in Chinese men Ooi L.S.M., Panesar N.S., Masarei J.R.L.	87

Enzyme-linked immunosorbent assay for. Extracellular glutathione peroxidase in serum of normal individuals and patients with renal failure on hemodialysis Yamamoto Y., Takekoshi Y., Itami N., Honjo T., Kojima H., Yano S., Takahashi H., Saito I., Takahashi K.	93
Neutrophil activation and collagen metabolism in lung cancer Gabazza E.C., Taguchi O., Yoshida M., Yamakami T., Kobayashi H., Ibata H., Shima T.	101
Erratum: The correlation in the interpretation of laboratory data (Clinica Chimica Acta 229 (1994) (1-3)) Chinchilli V.M., Gruemer H.-D.	113

Vol. 236/2, 1995

The value of polymorphonuclear elastase in adult respiratory distress syndrome Diaz J., Tornel P.L., Jara P., Canizares F., Egea J.M., Martinez P.	119
Polymorphonuclear leukocyte elastase levels in patients with Behcet's disease Deger O., Orem A., Akyol N., Bahadir S., Yildirmis S.	129
Amino acid sequence of a modified β_2 -microglobulin in renal failure patient urine and long-term dialysis patient blood Momoi T., Suzuki M., Titani K., Hisanaga S., Ogawa H., Saito A.	135
Evidence of toxic metabolite stress in black South Africans with chronic pancreatitis Gut A., Chaloner C., Schofield D., Sandle L.R., Purmasir M., Segal I., Braganza J.M.	145
Immunoglobulin cleavage by hypochlorous acid treatment Drozd R., Guevara I., Naskalski J.W.	155
Decreased branching, increased fucosylation and changed sialylation of alpha-1-proteinase inhibitor in breast and ovarian cancer Goodarzi M.T., Turner G.A.	161
Selenium dependent glutathione peroxidase activity in human follicular fluid Paszkowski T., Traub A.I., Robinson S.Y., McMaster D.	173
The typing of fetal antigen 2 in human amniotic fluid Price K.M., Silman R., Armstrong P., Teisner B., Grudzinskas J.G.	181
Demonstration of immunogenic keratan Sulphate in commercial chondroitin 6-sulphate from shark cartilage. Implications for ELISA assays Moller H.J., Moller-Pedersen T., Damsgaard T.E., Poulsen J.H.	195
Maximization of determination of basement membrane collagen complex in the sera from patients with chronic liver diseases by a sandwich ELISA Amano S., Miyake K., Shiba K., Yoshida T., Hayashi T.	205

Vol. 237/1-2, 1995

Novel enzymes as reagents Price C.P., Campbell R.S., Hammond P.M.	3
The effect of temperature on enzymes used in diagnostics Scopes R.K.	17
^{13}C NMR investigation of cholesterol esterification rate in human whole blood Nouri-Sorkhabi M.H., Sullivan D.R., Kuchel P.W.	25
Ligand western blotting for specific detection of active forms of proteases Kido T., Yook H.-Y., Ueda K.	31
Recombinant M-, B- and MB-type isozymes of human phosphoglyceric acid mutase: Their large-scale production and preparation of polyclonal antibodies specific to M- and B-type isozymes Uchida K., Kondoh K., Matuo Y.	43

Cardiac troponin, CK-MB and myoglobin for the early detection of acute myocardial infarction and monitoring of reperfusion following thrombolytic therapy Apple F.S., Voss E., Lund L., Preese L., Berger C.R., Henry T.D.	59
Tissue cathepsins as tumor markers Schwartz M.K.	67
Clinical applications of the p53 tumor suppressor gene Diamandis E.P.	79
Clinical and analytical considerations in the utilization of cholinesterase measurements McQueen M.J.	91
Nephrotoxicity of xenobiotics Werner M., Costa M.J., Mitchell L.G., Nayar R.	107
Beyond alcohol: Beverage consumption and cardiovascular mortality Goldberg D.M., Hahn S.E., Parkes J.G.	155

Vol. 238/1, 1995

Detection of autoantibodies against oxidized low-density lipoproteins and of IgG-bound low density lipoproteins in patients with coronary artery disease Boullier A., Hamon M., Walters-Laporte E., Martin-Nizart F., Mackereel R., Fruchart J.-C., Bertrand M., Duriez P.	1
Whether, to what degree, and why lipoprotein(a) levels change over time Glueck C.J., Tracy T., Sieve-Smith L., Wang P.	11
Quantitation of procholecystokinin and its products in plasma by processing-independent analysis Paloheimo L.I., Rehfeld J.F.	21
Glycaemia and body mass as determinants of plasma lecithin: Cholesterol acyltransferase activity in Nigerian patients with non-insulin dependent diabetes mellitus Akanji A.O., Agbedana E.O.	35
Compositions of very low density lipoprotein subfractions from patients with polydisperse low density lipoproteins Saheki S., Hitsumoto Y., Bando S., Murase M., Takeuchi N., Uchida K.	43
Relationships between triglycerides, lipoproteins, glucose and coronary artery disease Hobbs G.A., Wagner S.G., Levinson S.S.	59
Study of the chromium binding in plasma of patients on continuous ambulatory peritoneal dialysis Borguet F., Cornelis R., Delanghe J., Lambert M.-C., Lameire N.	71
Serum concentrations of the pyridinoline crosslinked carboxy-terminal telopeptide of type I collagen in chronic liver disease Hayasaka A., Iida S., Ohto M.	85
Low serum alkaline phosphatase activity associated with severe Wilson's disease. Is the breakdown of alkaline phosphatase molecules caused by reactive oxygen species? Hoshino T., Kumasaka K., Kawano K., Yamagishi F., Koyama I., Fujimori-Arai Y., Nakajima T., Komoda T.	91
Hyaluronic acid and dermatan sulfate in nonlesional pseudoxanthoma elasticum skin Lebwohl M., Longas M.O., Konstadt J., Koren R., Grabowski G.A., Phelps R., Lemlich G., Schwartz E.	101

Vol. 238/2, 1995

Identification of the D-enantiomer of 2-hydroxyglutaric acid in glutaric aciduria type II Watanabe H., Yamaguchi S., Saiki K., Shimizu N., Fukao T., Kondo N., Orii T.	115
Effect of neurotoxic metal ions in vitro on proteolytic enzyme activities in human cerebral cortex Falkous G., Harris J.B., Mantle D.	125
N-Methyladrenaline: Age-dependent urinary excretion, perinatal organ content and relation with 'classical' catecholamines Kema I.P., Schoots C.J.F., Gidding C.E.M., Okken A., Aarnoudse J.G., Muskiet F.A.J.	137

Maternal serum screening for fetal Down's syndrome, a retrospective study Lemay C., Roussel-Mizon N., Thepot F., Desmet G.	151
Variegate porphyria: Diagnostic value of fluorometric scanning of plasma porphyrins Da Silva V., Simonin S., Deybach J.C., Puy H., Nordmann Y.	163
Demonstration of induction of erythrocyte inosine monophosphate dehydrogenase activity in Ribavirin-treated patients using a high performance liquid chromatography linked method Montero C., Duley J.A., Fairbanks L.D., McBride M.B., Micheli V., Cant A.J., Morgan G.	169
Serotonin content in human blood platelets is not stable Karas S.I., Yazykov K.G., Makarova E.V.	179
Actions of arachidonic acid on erythrocyte membrane Rb permeability Dwight St. J.F.J., Hendry B.M.	187
Fasting plasma glucose as a diagnostic indicator of diabetes mellitus Wiener K.	199
Rapid degradation of 3,4-dihydroxyphenylalanine in plasma samples containing ethylenedi- aminetetraacetic acid Eriksson T., Holmgren B.	209
Erratum: Primary focal segmental glomerulosclerosis is associated with increased intraglomerular type IV collagen synthesis (<i>Clinica Chimica Acta</i> 235 (1995) (121-124)) Razzaque M.S., Koji T., Harada T., Nakane P., Taguchi T.	217

Vol. 239/1, 1995

Effects of urinary macromolecules on the nucleation of calcium oxalate in idiopathic stone formers and healthy controls Borghi L., Meschi T., Guerra A., Bergamaschi E., Mutti A., Novarini A.	1
Free fatty acid analysis in ascitic fluid improves diagnosis in malignant abdominal tumors Greco A.V., Mingrone G., Gasbarrini G.	13
Components of biological variation in plasma haptoglobin: Relationships to plasma fibrinogen and immune variables, including interleukin-6 and its receptor Maes M., Cooreman W., Delanghe J., Scharpe S., Wauters A., Neels H., D'Hondt P., Peeters D., Cosyns P., Ranjan R., Bosmans E., DeMeyer F., Elks H., Desnyder R.	23
A method for the separation of delta bilirubin using Cibacron Blue affinity chromatography Burnett J.R., Lim C.W., Mahoney G.N., Crooke M.J.	37
Interaction of human pregnancy-associated plasma protein-A with serine proteinases Zorin N.A., Zhabin S.G., Semenov N.N.	47
Sensitivity and specificity of a new ELISA method for determination of chromogranin A in the diagnosis of pheochromocytoma and neuroblastoma Boomsma F., Bhaggoe U.M., Man in 't Veld A.J., Schalekamp M.A.D.H.	57
Determination of ammonia in ear-lobe capillary blood is an alternative to arterial blood ammonia Huizenga J.R., Gips C.H., Conn H.O., Jansen P.L.M.	65
Micronutrient antioxidant status in black South Africans with chronic pancreatitis: Opportunity for prophylaxis Segal I., Gut A., Schofield D., Shiel N., Braganza J.M.	71
Plasma non-esterified fatty acid response to dialysate acetate concentration during haemodialysis in humans Jackson J.M., Talbot S.T., Lee H.A.	81
Quantitative cytochemistry of human leukocyte elastase compared with plasma elastase and acute phase proteins in inflammatory diseases Pradella M., Nemetz L., Bovo C., Visentin M., Clemen P., Zappala G., Ossani M., Plebani M.	91

Vol. 239/2, 1995

Daily variation of serum lipids in relation to the circadian rhythm of platelet aggregation in healthy male persons Knöfler R., Urano T., Taminato T., Yoshimi T., Nakano T., Nakajima K., Takada Y., Takada A.	109
--	-----

- New aspects concerning the regulation of the postoperative acute phase reaction during cardiac surgery
Berger D., Bolke E., Huegel H., Seidelmann M., Hannekum A., Begerb H.G. 121
- Enhanced susceptibility of low-density lipoprotein to in vitro oxidation in type 1 and type 2 diabetic patients
Beaudeau J.-L., Guillausseau P.-J., Peynet J., Flourie F., Assayag M., Tielmans D., Warnet A., Rousselet F. 131
- Comparison between serum levels of bone alkaline phosphatase and the carboxy-terminal propeptide of type I procollagen as markers of bone formation in patients following renal transplantation
Withold W., Degenhardt S., Grabensee B., Reinauer H. 143
- Bi-enzyme reactor for electrochemical detection of low concentrations of uric acid and glucose
Elekes O., Moscone D., Venema K., Korf J. 153
- Solid-phase competitive luminescence immunoassay for lysozyme in faeces
Gao P., John M.R., Schmidt-Gayk H., Arndt B., Scheida M., Theuer D. 167
- Osteocalcin metabolism in late fetal life: Fetal and maternal osteocalcin levels
Alatasa O., Colak O., Alatas E., Tekin B., Inal M., Ozalp S. 179
- The effect of ascorbic acid on the measurement of total cholesterol and triglycerides: Possible artefactual lowering in individuals with high plasma concentration of ascorbic acid
Benzie I.F.F., Strain J.J. 185
- Phenotyping of glutathione S-transferase m1 in the estonian population by ELISA using GSTM1a and GSTM1b specific monoclonal antibodies
Tasa G., Juronen E., Uuskula M., Viikmaa M., Mikelsaar A.-V. 191
- Creatine kinase isoenzymes in human trichinellosis
Tassia C., Materazzi L., Pozio E., Bruachi F. 197

Vol. 240/1, 1995

- Enzyme-linked immunosorbent assay for cholesteryl ester transfer protein in human serum
Sato T., Fukasawa M., Kinoshita M., Arai H., Saeki T., Naraki T., Iwasaki Y., Teramoto T., Takahashi K., Saito Y., Inoue K. 1
- New transcutaneous sampling of glucose for patients with type II diabetes using an ion-sensitive field-effect transistor
Kayashima S., Arai T., Noritake M., Nagata N., Kikuchi M., Ito N., Matsumoto Y., Kaneyoshi A., Kimura J., Kuriyama T. 11
- HPLC analysis of uridine diphosphate sugars: Decreased concentrations of uridine diphosphate galactose in erythrocytes and cultured skin fibroblasts from classical galactosemia patients
Xu Y.-K., Kaufman F.R., Donnell G.N., Giudici T., Alfi O., Ng W.G. 21
- 3-Hydroxyisovalerylcarnitine in patients with deficiency of 3-methylcrotonyl CoA carboxylase
Roachinger W., Millington D.S., Gage D.A., Huang Z.-H., Iwamoto T., Yano S., Packman S., Johnston K., Berry S.A., Sweetman L. 35
- Susceptibility to ulcerative colitis and Crohn's disease: Interactions between glutathione S-transferase GSTM1 and GSTT1 genotypes
Duncan H., Swan C., Green J., Jones P., Brannigan K., Aldersea J., Fryer A.A., Strange R.C. 53
- The total apolipoprotein B/LDL-cholesterol ratio does not predict LDL particle size
Tallis G.A., Shephard M.D.S., Sobocki S., Whiting M.J. 63
- Characterization of commercial antibodies for use in high resolution apo(a) phenotyping by immunoblot analysis
Gaw A., Chiesa G. 75
- Screening urine protein abnormalities by a novel test paper
Shiba K., Kanamori K., Sugiura K. 83
- Changes in red blood cell glutathione and glutathione-dependent enzymes on long-term treatment with captopril and enalapril
Golik A., Weisgarten J., Evans S., Cohen N., Zaidenstein R., Modal D., Cotariu D. 89
- Letters to the Editor*
- The stability of pyridinium crosslinks in urine and serum
Walne A.J., James I.T., Perrett D. 95

- No evidence for abnormalities in kinetics of platelet monoamine oxidase in Alzheimer's disease
Königs C.H., Scheltens P., Kuiper M.A., Wolters E.C. 99

Vol. 240/2, 1995

- Evaluation of the decarbamylation process of cholinesterase during assay of enzyme activity
Rotenberg M., Almog S. 107
- Effect of encapsulation on the anti-inflammatory properties of superoxide dismutase after oral administration
Regnault C., Roch-Arveiller M., Tissot M., Sarfati G., Giroud J.P., Postaire E., Hazebroucq G. 117
- An improved spectrophotometric assay of pyruvate dehydrogenase in lactate dehydrogenase contaminated mitochondrial preparations from human skeletal muscle
Chretien D., Pourrier M., Bourgeron T., Sene M., Rotig A., Munnich A., Rustin P. 129
- Development of an enzyme-linked immunosorbent assay to measure total TIMP-1 (free TIMP-1 and TIMP-1 in combination with matrix-metalloproteinases) and measurement of TIMP 1 and CRP in serum
Plumpton T.A., Clark I.M., Plumpton C., Calvin J., Cawston T.E. 137
- Comparison of 5-hydroxyindole-acetic acid and para-amino hippurate clearances in newborn rabbits
Sulyok E., Nagy L., Baranyai Z., Thonney M., Guignard J.-P. 155
- Plasma protein leakage and local secretion of proteins assessed in sputum in asthma and COPD. The effect of inhaled corticosteroids
Schoonbrood D.F.M., Out T.A., Lutter R., Reimert C.M., Van Overveld F.J., Jansen H.M. 163
- Elastase binding capacity of α_2 -macroglobulin and its association with glucocorticoid concentration in southern African black patients with hepatocellular carcinoma
Gaillard M.C., Song E., Nogueira C.M., Kilroe-Smith T.A. 179
- Separation of T and B lymphocytes from human peripheral blood mononuclear cells using density perturbation methods
Patel D., Rubbi C.P., Rickwood D. 187
- An improved GLC method for a rapid, simultaneous analysis of both medium chain fatty acids and medium chain triglycerides in plasma
Mingrone G., Greco A.V., Capristo E., Benedetti G., Castagneto M., Gasbarrini G. 195
- Preferential depletion of selenoprotein P in hypercholesterolaemic patients treated by LDL-apheresis
Persson-Moschos M., Bonnefont-Rousselot D., Assogba U., Bruckert E., Jaudon M.C., Delattre J., Akesson B. 209
- Tartrate resistant acid phosphatase in the serum of patients with larynx carcinoma — A marker of osteoclast activation
Jarkowska-Kozłowska J., Kozłowski Z. 213

Vols. 241-242, 1995

- Clinical mass spectrometry
Niwa T. 3

Vol. 243/1, 1995

- Enzyme-linked immunosorbent assay (ELISA) for *Aleuria aurantia* lectin-reactive serum cholinesterase to differentiate liver cirrhosis and chronic hepatitis
Kondo M., Hada T., Fukui K., Iwasaki A., Higashino K., Yasukawa K. 1
- Urinary C4 excretion in systemic lupus erythematosus
Ueda Y., Nagasawa K., Tsukamoto H., Horiuchi T., Yoshizawa S., Tsuru T., Furugo I., Niho Y. 11

Non-specific increase of serum carbohydrate antigen 19-9 in patients with liver disease associated with increased circulating levels of adhesion molecules Fabris C., Falletti E., Pirisi M., Soardo G., Toniutto P., Vitulli D., Bortolotti N., Gonano F., Bartoli E.	25
A trinucleotide deletion together with a base duplication event at codon 439 in the human tyrosinase gene identifies a mutational hotspot Breimer L.H., Winder A.F., Panayiotidis P., Jay M., Moore A., Jay B.	35
Direct double antibody sandwich immunoassay of mucin M ₁ epitopes in human mucus secreting pancreatic cell lines Montserrat C., Hollande E., Guy-Crotte O., Figarella C.	43
Type I collagen synthesis by skin fibroblasts from 17 patients with osteogenesis imperfecta type III Pocock A.E., Francis M.J.O., Smith R.	53
Effects of membrane incorporation of short-chain phospholipids on sodium pump function in human erythrocytes Dwight J.F.S.J., Hendry B.M.	73
Plasma carnitine concentrations in pregnancy, cord blood, and neonates and children Winter S.C., Linn L.S., Helton E.	87
Intra-individual variation of overnight urinary albumin excretion in clinically healthy individual middle-aged individuals Jensen J.S.	95
Erratum: Nephrotoxicity of xenobiotics (<i>Clinica Chimica Acta</i> 237 (1995) 107-154) Werner M., Costa M.J., Mitchell L.G., Nayar R.	101

Vol. 243/2, 1995

Aminooxy acetic acid: A selective inhibitor of alanine:glyoxylate aminotransferase and its use in the diagnosis of primary hyperoxaluria type I Horvath V.A.P., Wanders R.J.A.	105
Acetylcholinesterase of human intestinal tissue affected by Hirschsprung's disease: Effect of magnesium chloride on isoforms Johnson G., Moore S.W., Purves L.R.	115
Measurement of mevalonic acid in human urine by bench top gas chromatography-mass spectrometry Siavoshian S., Simoneau C., Maugeais P., Marks L., Rodary L., Gardette J., Krempf M.	129
Enhanced susceptibility of low-density lipoproteins to oxidation in coronary bypass patients with progression of atherosclerosis De Rijke Y.B., Verwey H.F., Vogelezang C.J.M., Van Der Velde E.A., Princen H.M.G., Van Der Laarse A., Bruschke A.V.G., van Berkel T.J.C.	137
Purine and carnitine metabolism in muscle of patients with Duchenne muscular dystrophy Camina F., Novo-Rodriguez M.I., Rodriguez-Segade S., Castro-Gago M.	151
Determination of serum total lipid and free N-acetylneuraminic acid in genitourinary malignancies by fluorimetric high performance liquid chromatography. Relevance of free N-acetylneuraminic acid as tumour marker Lagana A., Pardo-Martinez B., Marino A., Fago G., Bizzarri M.	165
Characterization of mitochondrial DNA in primary cardiomyopathies Bobbà A., Giannattasio S., Pucci A., Lippolis R., Camaschella C., Marra E.	181
Evaluation of antibody- and antigen-coated enzymeimmunoassays for measuring oestrone-3-glucuronide concentrations in urine Henderson K.M., Camberis M., Hardie A.H.	191
Does plasma neuropeptide-Y immunoreactivity in patients with pheochromocytoma depend on hormonal activity of the tumor? Pruszczyk P., Wocial B., Ignatowska-Switalska H., Feltynowski T., Ellafi M., Januszewicz A., Lapinski M., Zukowska-Grojec Z., Januszewicz W.	205

Vol. 244/1, 1996

Characterization of lipoproteins during human cholestasis Tallet F., Vaasson M.-P., Couderc R., Lefevre G., Raichvarg D.	1
Determination of metalloproteinases, plasminogen-activators and their inhibitors in the synovial fluids of patients with rheumatoid arthritis during chemical synoviorthesis Blaser J., Triebel S., Maasjosthusmann U., Romisch J., Krahl-Mateblowski U., Freudenberg W., Fricke R., Tachesche H.	17
Plasma β -hexosaminidase isoenzymes A and B in patients with cerebral infarction Hultberg B., Isaksson A., Lindgren A., Israelsson B., Brattstrom L.	35
Phosphates precipitating from artificial urine and fine structure of phosphate renal calculi Grases F., Sohnel O., Vilacampa A.I., March J.G.	45
High plasma level of plasmin- α 2-plasmin inhibitor complex is predictor of poor prognosis in patients with lung cancer Taguchi O., Gabazza E.C., Yoshida M., Yamakami T., Kobayashi H., Shima T.	69
Diagnostic discrimination and cost effective assay strategy for leukocyte alkaline phosphatase DePalma L., Delgado P., Werner M.	83
Detection of iron-deficiency anemia in hospitalized patients by zinc protoporphyrin Wong S.S., Qutishat A.S., Lange J., Gornet T.G., Buja L.M.	91
Simultaneous extraction of sufentanil and midazolam from human plasma Rossi S.S., Dyck J.B., Yaksh T.L.	103
A sensitive microplate-based continuous-monitoring (kinetic) assay for serum neutral endopeptidase (EC 3.4.24.11) activity Selmeci L., Szokodi I., Horvat-Karajz K.	111

Vol. 244/2, 1996

International Federation of Clinical Chemistry. A strategy to promote the rational use of laboratory tests Pannall P., Marshall W., Jabor A., Magid E.	121
A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 8 (neutrophil collagenase) using monoclonal antibodies Matsuki H., Fujimoto N., Iwata K., Knauper V., Okada Y., Hayakawa T.	129
Use of β -maltosides (p-nitrophenyl- β -D-maltoside, 2-chloro-4-nitrophenyl- β -D-maltoside and 4-methylumbelliferyl- β -D-maltoside) as substrates for the assay of neutral α -glucosidase from human kidney and urine Lukomska I.S., Voznyi Y.V., Lanska I.M., Podkidisheva E.I.	145
Evaluation of endotoxiuria for diagnosis of urinary tract infection after major surgical procedures Berger D., Boelke E., Seidelmann M., Beger H.G.	155
Plasma parathyroid hormone related-protein levels in patients with cancer, normocalcemic and hypercalcemic Segura Dominguez A., Andrade Olivie M.A., Rodriguez Sousa T., Terron Alvarez M.L., Rodriguez Perez D., Alvarez Novoa R., Garcia-Mayor R.V.G.	163
Investigation of erythrocyte membrane lipid peroxidation and antioxidant defense systems of patients with coronary artery disease (CAD) documented by angiography Akkus I., Saglam N.I., Caglayan O., Vural H., Kalak S., Saglam M.	173
A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 7 (matrilysin) using monoclonal antibodies Ohuchi E., Azumano I., Yoshida S., Iwata K., Okada Y.	181
Increased serum levels of neuron-specific enolase in epileptic patients and after electroconvulsive therapy — A preliminary report Greffé J., Lemoine P., Lacroix C., Brunon A.-M., Terra J.-L., Dalery J., Bernier E., Soares-Bouchaud I., Rochet T., Leduc T.A., Balvay G., Mathieu P.	199
Radioimmunoassay of free thyroxine (T4) using 125 I-labeled T4-IgG complex with very large molecular weight Georgiou S., Christofidis I.	209

Leukocyte lipid peroxidation, superoxide dismutase, glutathione peroxidase and serum and leukocyte vitamin C levels of patients with type II diabetes mellitus	
Akkus I., Kalak S., Vural H., Caglayan O., Menekse E., Can G., Durmus B.	221
On the stability of human lysosomal enzymes at room temperature in normal and acidified plasma and serum	
Den Tandt W.R.	229

Vol. 245/1, 1996

Human serum hyaluronidase: Characterization of a clinical assay	
Natowicz M.R., Wang Y.	1
Stabilization of plasmin by lysine derivatives	
Ueshima S., Okada K., Matsuo O.	7
Cardiac troponin I in the diagnosis of myocardial injury and infarction	
Mair J., Genser N., Morandell D., Maier J., Mair P., Lechleitner P., Calzolari C., Larue C., Ambach E., Dienstl F., Pau B., Puschendorf B.	19
Immunochemiluminometric assay with two monoclonal antibodies against the N-terminal sequence of human parathyroid hormone	
Gao P., Schmidt-Gayk H., Ditttrich K., Nolting B., Maier A., Roth H.J., Soemann O., Reichel H., Ritz E., Schilling T.	39
A simple method for screening for Farber disease on cultured skin fibroblasts	
Chatelut M., Feunteun J., Harzer K., Fensom A.H., Basile J.-P., Salvayre R., Levade T.	61
Age-related changes in amounts and concentrations of collagen and elastin in normotensive human thoracic aorta	
Cattell M.A., Anderson J.C., Hasleton P.S.	73
A fast and simple screening method for detection of 2,8-dihydroxyadenine urolithiasis by capillary zone electrophoresis	
Sevcik J., Adam T., Mazacova H.	85
Perilymph detection by β 2-transferrin immunoblotting assay. Application to the diagnosis of perilymphatic fistulae	
Delaroche O., Bordure P., Lippert E., Sagniez M.	93
Determination of phosphoribosylpyrophosphate synthetase activity in human cells by a non-isotopic, one step method	
Torres R.J., Mateos F.A., Puig J.G., Becker M.A.	105
Problems and perspectives of clinical biochemistry training, and the example of Italy	
Salvatore F., Sacchetti L.	113
Letters to the Editor	
Concentration of mRNA for von Willebrand factor in platelets of type I von Willebrand disease	
Takenaka T., Kuribayashi K., Tsukiyama M., Nakamine H., Fukuhara Y., Kuno T.	125
Helicobacter pylori infection and serum IgG avidity	
Basso D., Brigato L., Di Mario F., Plebani M.	129

Vol. 245/2, 1996

Hereditary hemochromatosis	
Witte D.L., Crosby W.H., Edwards C.Q., Fairbanks V.F., Mitros F.A.	139
Degradation of glycated hemoglobin by erythrocytic proteolytic enzymes	
Raghothama C., Rao P.	201
Prevalence of biotin-binding immunoglobulin in patients with allergic and autoimmune diseases	
Nagamine T., Takagi H., Sugimoto H., Takehara K., Fukui T., Mori M.	209
Properties and units in the clinical laboratory sciences. I. Syntax and semantic rules. IUPAC-IFCC recommendations 1995	
Olesen H.	S5

Properties and units in the clinical laboratory sciences. V. Properties and units in thrombosis and haemostasis. ISTH-IUPAC-IFCC Recommendations 1995

Blomback M., Dybkaer R., Jorgensen K., Olesen H., Thorsen S. \$23

Vol. 246/1-2, 1996

Assessing moderate alcohol consumption as a personal risk factor	
Werner M.	5
The pathogenesis of atherosclerosis	
Hegele R.A.	21
Effects of alcohol on plasma lipoprotein metabolism	
Frohlich J.J.	39
Alcohol and coronary heart disease: Consistent relationship and public health implications	
Criqui M.H.	51
Alcohol and coronary heart disease: The evidence for a protective effect	
Kannel W.B., Ellison R.C.	59
Effects of alcohol on platelet functions	
Renaud S.C., Ruf J.-C.	77
Alcohol and hypertension	
Klatsky A.L.	91
Case-control and cohort studies of moderate alcohol consumption and stroke	
Camargo Jr. C.A.	107
Epidemiologic data on alcoholic beverage consumption and risk of cancer	
Longnecker M.P., Enger S.M.	121
The questionable link between alcohol intake and cancer	
Rubin E.	143
'Moderate' drinking during pregnancy: Cause for concern?	
Abel E.L.	149
Mild maternal drinking and pregnancy outcome: Perceived versus true risks	
Koren G., Koren T., Gladstone J.	155
Wines and grape juices as modulators of platelet aggregation in healthy human subjects	
Pace-Asciak C.R., Rounova O., Hahn S.E., Diamandis E.P., Goldberg D.M.	163
Wine: Does the colour count?	
Goldberg D.M., Garovic-Kocic V., Diamandis E.P., Pace-Asciak C.R.	183

Vol. 247/1,2, 1996

An improved technique for the assay of red blood cell superoxide dismutase (SOD) activity	
L.E. Ilouno, E.N. Shu, G.E. Igboke	1
The activity of a blood type B specific exoglycosidase from <i>Glycine max</i>	
L. Hobbs, M. Mitra, R. Phillips, D. Smith	7
Immunoluminometric assay (ILMA) for intact human proinsulin and its conversion intermediates	
T.M. Zilkens, A.M. Eberle, H. Schmidt-Gayk	23
Plasma lysosomal glycohydrolases in a general population	
A. Lombardo, C. Bairati, G. Goi, C. Roggi, L. Maccarini, D. Bollini, A. Burlina	39
Plasma 1,5-anhydro-D-glucitol concentration and its relation to other plasma components in renal failure and renal transplant recipients	
G. Phillipov, V.T. Ninan, T.H. Mathew	51
Peptic-tryptic digests of gliadin: contaminating trypsin but not pepsin interferes with gastrointestinal protein binding characteristics	
G. Bolte, A. Osman, T. Mothes, M. Stern	59
Serum paraoxonase activity is decreased in uremic patients	
R. Schiavon, E. De Fanti, D. Giavarina, S. Biasioli, G. Cavalcanti, G. Guidi	71

Ketone body ratios of the superior and inferior vena cava and of pulmonary arterial blood compared to that of arterial blood: central venous ketone body ratio as a substitute for the arterial ketone body ratio	
Y. Terada, Y. Eguchi, Y.-J. Chang, R. Tabata, H. Sakamoto, O. Takehiro, S. Uno, K. Ozawa	81
Specific oncological contribution of cathepsin D and pS2 in human breast cancer: their relationship with TNM status, estradiol receptors, epidermal growth factor receptor and <i>neu</i> amplification	
P. Bolufer, M.D. Torregrosa, L. Gomez, B. Munarriz, J.A. López, E. Asins, S. Espinoza, F. Vera, C. Vazquez, V. Guillem	89
Intracellular degradation of sulforhodamine-G _{M1} : use for a fluorescence-based characterization of G _{M2} -gangliosidosis variants in fibroblasts and white blood cells	
V. Agmon, R. Khosravi, S. Marchesini, T. Dinur, A. Dagan, S. Gatt, R. Navon	105
Analysis of plasma isoflavones by reversed-phase HPLC-multiple reaction ion monitoring-mass spectrometry	
L. Coward, M. Kirk, N. Albin, S. Barnes	121
Impairment of ganglioside metabolism in cultured fibroblasts from Salla patients	
M. Pitto, V. Chigorno, M. Renlund, G. Tettamanti	143
Characterization of low-density-lipoprotein in apolipoprotein E deficiency in a patient without coronary atherosclerosis	
T. Teramoto, M. Kinoshita, M. Kawamura, Y.-Q. Liang, K. Kaneko, N. Shimazu, S. Suga, N. Akatuka, M. Yamanaka	159
Characterization of clinical assays for leukocyte and fibroblast alpha-N-acetylgalactosaminidase activities for the diagnosis of alpha-N-acetylgalactosaminidase deficiency	
E.M. Prencz, J. Gleason, M.R. Natowicz	167
Influence of apolipoprotein E phenotypes on total serum cholesterol in Japanese children	
M. Tozuka, H. Ohta, H. Hidaka, N. Okumura, K. Furihata, T. Katsuyama	175
Increased plasma thiobarbituric acid-reactive substances (TBARS) before opportunistic infection symptoms in HIV infected individuals	
O. Lopez, D. Bonnefont-Rousselot, M. Mollereau, R. Olivier, L. Montagnier, J. Emerit, M. Gentilini, J. Delattre	181
Specific carbohydrate antigen LeY expression on lymphocytes in diabetes mellitus	
T. Nagai, T. Tomizawa, K. Sudou, C. Shiozawa, M. Mori	189
<i>Letter to the Editor</i>	
Simultaneous metabolic profile studies of three patients with fatal infantile mitochondrial myopathy-de Toni-Fanconi-Debré syndrome by GC/MS	
C. Ning, T. Kuhara, I. Matsumoto	197

Vol. 248/1, 1996

Computing in Clinical Laboratories

CCL 94—The 10th International Conference, Jerusalem, Israel, 29 May–3 June 1994

Foreword	1
The CCL conferences revisited	
H.E. Solberg (Norway)	3
The clinical interface — a British Physician's view	
J. Neuberger, M. Peters	11
Computer architecture for the clinical laboratory	
A new clinical laboratory information system architecture from the OpenLabs project offering advanced services for laboratory staff and users	
G. Boran, R. O'Moore, W. Grimson, M. Peters, A. Hasman, T. Groth, F. Van Merode	19
Client/Server computing: is this the future direction for the clinical laboratory?	
I.G. Wells, L.P. Farnan, M.W. Rayment	31
Automated technical validation — a real time expert system for decision support	
J.S. de Graeve, J.P. Cambus, A. Gruson, P.M. Valdigué	39

The clinical workstation as a means of improving laboratory use	
D.P. Connelly, B.H. Sielaff, K.E. Willard	51
Improvement of emergency and routine turnaround time by data processing and instrumentation changes	
M.D. Pelegri, L. García-Beltrán, C. Pascual	65
<i>Neural network technology</i>	
Neural networks and fuzzy logic in clinical laboratory computing with application to integrated monitoring	
A. Guez, I. Nevo	73
Self-organizing neural networks — an alternative way of cluster analysis in clinical chemistry	
G. Reibnegger, H. Wachter	91
Introduction of a new laboratory test: an econometric approach with the use of neural network analysis	
A. Jabor, T. Vlk, P. Bofil	99
<i>Applications</i>	
Computer assisted interpretation of laboratory test data with 'MDI-LabLink'	
A. Regeniter, W.H. Siede, U.B. Seiffert	107
Learning clinical biochemistry using multimedia interactive clinical cases	
J. Hooper, J. O'Connor, R. Cheesmar	119

Vol. 248/2, 1996

4-Pentafluoroethylumbelliferyl- β -D-glucoside as a new fluorogenic substrate for acid β -D-glucosidase	
I.V. Tsvetkova, E.A. Karpova, T.V. Dudukina, Y.V. Voznyi	125
Evaluation of nonisotopic binding assays for measuring vitamin B ₁₂ and folate in serum	
L.S.W. Steijns, J.K. Braams-Wiatrowska, H.-J. Luiting, J. van der Weide	135
Amniotic fluid for screening of lysosomal storage diseases presenting in utero (mainly as non-immune hydrops fetalis)	
M. Piraud, R. Froissart, G. Mandon, A. Bernard, I. Maire	143
Cholesteryl ester transfer activity in liver disease and cholestasis, and its relation with fatty acid composition of lipoprotein lipids	
A. Iglesias, M. Arranz, J.J. Alvarez, J. Perales, J. Villar, E. Herrera, M.A. Lasunción	157
Bile acid conjugation in early stage cholestatic liver disease before and during treatment with ursodeoxycholic acid	
M. Fracchia, K.D.R. Setchell, A. Crosignani, M. Podda, N. O'Connell, Roberto Ferraris, A.F. Hofmann, G. Galatola	175
The carboxy-terminal cross-linked telopeptide of type I collagen (ICTP) is a potential serum marker of ongoing liver fibrosis	
S. Ricard-Blum, P. Chossegros, S. Guerret, C. Trepo, J.-A. Grimaud, M. Chevallier	187
Analysis of the deletion/insertion polymorphism of the angiotensin I-converting enzyme gene by capillary electrophoresis	
M. Koch, R. Lehmann, M. Pfohl, W. Voelter, H. Liebich	197
<i>Letters to the Editor</i>	
Ketone body ratios of capillary blood compared with that of arterial blood	
H. Sakumoto, Y. Terada, R. Tabata, Y. Eguchi, O. Takehiro, S. Uno, T. Banba, K. Ozawa	205
Effect of ascorbic acid deficiency on plasma prothrombin time and partial thromboplastin time in guinea pigs	
M.W. Roomi, C.S. Tsao	209
ISCE Newsletter	51

Vol. 249/1,2, 1996

Conference report

Medical, economic, and regulatory factors affecting point-of-care testing. A Report of the Conference on Factors Affecting Point-of-Care Testing	
B. Seamonds	1

Evaluation of multicomponent fluorimetric analytical data by use of partial least-squares calibration. Application to the determination of codeine and ephedrine in urine	
R.D. Bautista Jiménez, A.I. Jiménez Abizanda, F. Jiménez Moreno, J.M. Arias León	21
Defective hepatobiliary leukotriene elimination in patients with the Dubin-Johnson syndrome	
E. Mayatepek, W.D. Lehmann	37
Effect of prednisone on protease activities and structural protein levels in rat muscles in vivo	
J.W. Haycock, G. Falkous, C.A. Maltin, M.I. Delday, D. Mantle	47
Lithogenicity of human bile is reduced by freezing and thawing	
F.M. Konikoff, H. Laufer, T. Gilat	59
Reference intervals for some serum biochemical markers of bone metabolism in Kuwait	
K. Prabha, A. Al-Suliman, S.O. Olusi, T.N. Sugathan	67
Serum neuron specific enolase (S-NSE) reference interval evaluation by time-resolved immunofluorometry compared with a radioimmunoassay	
L.G.M. Jørgensen, J. Løber, N.L.T. Carlsen, G. Mømsen, F.R. Hirsch	77
Production and characterization of monoclonal antibodies against L-carnitine: radioimmunologic assays for L-carnitine determination	
S. Marini, G.F. Fasciglione, B. Giardina	93
Serum alkaline phosphatase activity during zinc deficiency and long-term inflammatory stress	
T.H.J. Naber, H. Baadenhuysen, J.B.M.J. Jansen, C.J.A. van den Hamer, W. van den Broek	109
Comparison of proline endopeptidase activity in brain tissue from normal cases and cases with Alzheimer's disease, Lewy body dementia, Parkinson's disease and Huntington's disease	
D. Mantle, G. Falkous, S. Ishiura, P.J. Blanchard, E.K. Perry	129
Kinetic nephelometric determination of albumin produced by rat hepatocytes in culture	
P. Pernet, M. Vaubourdolle, S.-Kyung Lim, L. Cynober, J. Giboudeau	141
Alterations of erythrocyte morphology and lipid composition by hyperbilirubinemia	
M.A. Brito, R.M. Silva, D.C. Matos, A.T. da Silva, D.T. Brites	149
The effects of <i>tert</i> -butyl hydroperoxide on human erythrocyte membrane ion transport and the protective actions of antioxidants	
J.F.St.J. Dwight, B.M. Hendry	167
Plasma thiols, cognitive function and ageing	
S.W. Welsh, F.M. Corrigan, W.E. Smith, D.F. Horrobin	183
The use of calcein acetomethylester (AM)-labelled polymorphonuclear cells in a polycarbonate filter chemotaxis assay	
C.M. De Gendt, L.S. De Clerck, C.H. Bridts, W.J. Stevens	189
<i>Letter to the Editor</i>	
Adenylosuccinate lyase deficiency and disorders of saccharide metabolism; experience with combined screening test	
D. Valika, J.D. Jones	197

THE HISTORY OF THE

REIGN OF THE MOST CHRISTIAN KING OF FRANCE

CHARLES THE NINTH

BY M. DE LAURENCE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

OF THE ACADEMY OF SCIENCES

AND OF THE ACADEMY OF HISTORY

AND OF THE ACADEMY OF MEDICINE

